

Complexes with free actions by the group of automorphisms of a free group

Jeff Kiralis

1. INTRODUCTION

We define two finite-dimensional complexes on which the group $\text{Aut}(F_n)$ acts freely, and study their homotopy groups. The definitions of these complexes were motivated by low-dimensional pseudo-isotopy theory. So not surprisingly, the complexes are related to a number of spaces arising in algebraic K-theory, and to another in low-dimensional topology.

The complexes are denoted by X_n and X_n^{alg} . The cells of X_n correspond bijectively to certain pointed graphs which are marked by homotopy equivalences with $\bigvee_{i=1}^n S^1$, the wedge of n circles. Each of these marked graphs is required to have some additional structure which we call a coloring. The action of $\text{Aut}(F_n)$ on X_n is defined by precomposing the markings with maps $\bigvee_{i=1}^n S^1 \rightarrow \bigvee_{i=1}^n S^1$ corresponding to elements of $\text{Aut}(F_n)$. The complex X_n^{alg} is defined by algebraically mimicing the definition of X_n . There is an equivariant map $X_n \rightarrow X_n^{alg}$ which, after stabilizing, may be a homotopy equivalence.

Forgetting the colorings defines an equivariant map from X_n to the sphere complex $\mathcal{S}_n$. This last complex was introduced by Hatcher in [3]. Forgetting basepoints as well gives a map from X_n to Outer space. Outer space was defined by Culer and Vogtmann in [1], and denoted by X_n there. The colorings provide just enough structure to eliminate the fixed points of $\mathcal{S}_n$, which $\text{Aut}(F_n)$ acts on, but not freely. From the point of view of pseudo-isotopy theory, the cells of X_n may be thought of as parametrized families of certain handle-body structures of the 4-disk, or equivalently, of smooth functions on D^4 . The colorings of the graphs prevent certain higher singularities from occurring among these functions so that the homotopy groups of X_n resemble a K -theory. There is no such restriction on the singularities with $\mathcal{S}_n$, which may be thought of as a sort of low-dimensional Whitehead space, with a little added structure.

The space $X = \lim_{n \rightarrow \infty} X_n$ (and probably $X^{alg} = \lim_{n \rightarrow \infty} X_n^{alg}$ as well) should perhaps be thought of as a pre- K -theory, with the corresponding K -theory indicated in the remarks at the end of §2.4. The homotopy groups $\pi_i(X_n)$ depend on n , even for $i \gg n$, but seem to be more amenable to computation using elementary techniques than the probably stable homotopy groups of the corresponding K -theory.

The homotopy groups of X and X^{alg} map to the algebraic K -groups of the ring $\mathbb{Z}$, essentially by abelianizing F_n , or, with pseudo-isotopy theory in mind, by suspending the families of smooth functions mentioned above. More specifically, there is a map $h: X^{alg} \rightarrow V$, where $V \equiv V(\mathbb{Z})$ is the space with $\pi_{n-1}(V) = K_n(\mathbb{Z})$ defined by Volodin, as in [11].

The complex X_n^{alg} is, in some ways, similar to the unstable Volodin space V_n , but with $\text{Aut}(F_n)$ in place of $\text{GL}_n(\mathbb{Z})$. A difference is that, while $\tilde{V}/\text{St}(\mathbb{Z})$ is acyclic, $\tilde{X}^{alg}/\text{StN}$ is not. (Here the tildes denote universal covers of a component of V and X . Also, the group StN , introduced by Gersten in [2], and also studied in [7], is a nonabelian version of $\text{St}(\mathbb{Z})$, the Steinberg group of $\mathbb{Z}$.) Indeed, the homology group $H_2(\tilde{X}^{alg}/\text{StN})$ is nontrivial, as is $H_2(\tilde{X}/\text{StN})$. The elements of these groups detect elements of $\pi_2(X)$ which map onto the cokernel of the map $\pi_3^S \rightarrow K_3(\mathbb{Z})$.

There is also a map f which makes the diagram

$$\begin{array}{ccc} \Omega B\text{StN}^+ & \xrightarrow{f} & \tilde{X}^{\text{alg}} \\ \downarrow g & & \downarrow \tilde{h} \\ \Omega B\text{St}(\mathbb{Z})^+ & \xrightarrow{\sim} & \tilde{V} \end{array}$$

homotopy commute, where $\tilde{h}$ is induced by $h: X_{\text{alg}} \rightarrow V$. The lower horizontal map is a homotopy equivalence coming from the one between V and $\Omega B\text{GL}(\mathbb{Z})^+$ mentioned above.

The main results, summarized in Theorem 12.2, are the exactness of the sequence

$$0 \longrightarrow H_3(\text{StN}) \longrightarrow H_2(\tilde{X})_{\text{StN}} \xrightarrow{q_*} H_2(\tilde{X}/\text{StN}) \longrightarrow 0$$

and a computation of its terms. In particular, the group $H_3(\text{StN})$ is cyclic of order 24. While this is analogous to Lee and Szarbo's result in [8] that the group $H_3(\text{St}(\mathbb{Z})) \approx K_3(\mathbb{Z})$ is cyclic of order 48, our methods are completely different from theirs. We show in §3 and the following two that the group $H_3(\tilde{X}_n/\text{StN}_n)$ is trivial, which gives the injectivity of the map $H_3(\text{StN}) \longrightarrow H_2(\tilde{X})_{\text{StN}}$. The homomorphism q_* in the sequence is easily computed using the isomorphism Θ defined in §6, where $H_2(\tilde{X}_n/\text{StN}_n)$ is also computed. The contractibility of the sphere complex is used in §11 to show that every 2-cycle in $\tilde{X}$ is homologous to one of a relatively simple type. Such cycles are worked with in §10 and §12 to compute $H_2(\tilde{X})_{\text{StN}}$. Along the way, we give many explicit descriptions of generators of $K_3(\mathbb{Z})$ as 2-cycles in $\tilde{X}$ (or $\tilde{V}$), and show in an elementary way that they have order 48. We also point out in Theorem 12.3 that these generators appear as nontrivial elements of the homology of certain upper triangular subgroups of $\text{Aut}(F_n)$.

My main motivation for writing this paper was to compute $H_3(\text{StN}) = H_3(\text{Aut}(F_n)) = \pi_3(B\text{Aut}(F_n)^+)$, for large n . But Galatius has recently announced that the homology groups of $\text{Aut}(F_n)$ and the symmetric group are the same stably in *all* dimensions. The spaces $\tilde{X}$ and $\tilde{X}/\text{StN}$ may still have some interest.

2. PRELIMINARIES

2.1. Partitioned graphs. In this paper, the word graph means a connected 1-dimensional CW-complex with finitely many vertices. A graph G will often be specified combinatorially by two sets $E(G)$ and $V(G)$, the edges and vertices of G , respectively, along with two functions $d_0, d_1: E(G) \rightarrow V(G)$ which determine the attaching maps of the 1-cells $E(G)$ of G . The rank of a graph is, by definition, the rank of its fundamental group. We often identify cycles in a graph G , as well as subgraphs of G (with each component larger than a vertex), with the subset of $E(G)$ they comprise.

A partition of a graph G of rank n is a partition of the set $E(G)$ of edges of G into n nonempty sets $h_1, \dots, h_n$ such that any nontrivial cycle of G contains one of the h_i . The sets h_i are called the edge sets of the partitioned graph G . If e is the only element of an h_i , then e is called a singleton edge, and h_i a singleton edge set. We often write $(G, \{h_i\}, n)$, or just $(G, \{h_i\})$, for the partitioned graph G with edge sets $h_i, \dots, h_n$.

If G is a graph with edge e , let G_e denote the graph obtained from G by collapsing e to a point. That is, G_e is the graph with $E(G_e) = E(G) - \{e\}$ and $V(G_e) = V(G)/d_0(e) \sim d_1(e)$, and with attaching maps those of G , followed by the quotient map $V(G) \rightarrow V(G_e)$.

Lemma 2.1. *Let $(G, \{h_i\}, n)$ be a partitioned graph. If $e \in h_k$ where $|h_k| > 1$, then the graph G_e is partitioned by the edge sets $h_1, \dots, h_{k-1}, h'_k, h_{k+1}, \dots, h_n$ where $h'_k = h_k - \{e\}$, and the quotient map $f: G \rightarrow G_e$ is a homotopy equivalence.*

We say that the partition of G_e in the lemma is induced from the partition of $(G, \{h_i\})$.

Proof of 2.1. Since $|h_k| > 1$, the edge e cannot itself form a cycle of length one. For such a cycle would not contain any of the edgesets. Thus the collapsing map $f: G \rightarrow G_e$ is a homotopy equivalence.

Any cycle C_e of G_e corresponds, via f , to a unique cycle C of G . Since $h_i \subset C$ for some i we have either $h_i \subset C_e$, if $i \neq k$, or $h'_k \subset C_e$ if $i = k$. $\square$

Quotient maps $f: G \rightarrow G_e$ between partitioned graphs where the cardinality of the edge set containing e is > 1 , and where G_e has the induced partition, are called blowdowns, as are compositions of such maps. If $f: G \rightarrow G'$ is a blowdown, we say that G' is obtained from G by blowing down the edges of G which are collapsed by f .

An edge basis of a partitioned graph $(G, \{h_i\}, n)$ is, by definition, any set of n edges $\{e_1, \dots, e_n\}$ where $e_i \in h_i$ for each i . If $\{e_i\}$ is an edge basis for $(G, \{h_i\})$, we always assume, without comment, that $e_i \in h_i$ for each i .

The complement of any edge basis is a maximal tree. More precisely we have:

Lemma 2.2. *Assume that $E = \{e_1, \dots, e_n\}$ is an edge basis for the partitioned graph $(G, \{h_i\})$. Then the subgraph T^E of G consisting of all the edges of G except for those in E is a maximal tree in G .*

Proof. The subgraph $T = T^E$ has no cycles since any cycle in G contains an edge set h_i for some i .

By Lemma 2.1, all the edges of T can be blown down giving a graph we call R . Since R and G are homotopy equivalent, we have

$$\chi(R) = \chi(G) = \text{rank of } H_0(G) - \text{rank of } H_1(G) = 1 - n.$$

Since R has n edges, it follows that it has just one vertex. Therefore T is connected and contains all the vertices of G . So T is maximal. $\square$

Assume that $E = \{e_i\}$ is an edge basis for the partitioned graph $(G, \{h_i\})$, and let T^E be the corresponding maximal tree as defined in the lemma. Let G_i be the subgraph of G formed by adding the single edge e_i to T^E . Denote by C_i^E , or just C_i , the unique (shortest) cycle in G_i . Removing any edge in h_i from G_i gives, by Lemma 2.2, a tree. It follows that $h_i \subset C_i$. Note that C_i contains none of the other edge sets h_j for $j \neq i$ since C_i contains none of the edges e_j for $j \neq i$.

The C_i are called basic cycles (associated with the edge basis E). If a vertex $*$ of G is chosen as a basepoint, then the basic cycles determine a basis for the free group $\pi_1(G, *)$.

Some basic facts about basic cycles are given in the next two lemmas.

Lemma 2.3. *If C_j and C_k are two basic cycles associated with the edge basis $E = \{e_i\}$, then $C_j \cap C_k$ is connected.*

Proof. Let T be the maximal tree associated with the edge basis E and let $x, y \in T$. Denote by $[x, y]$ the smallest connected set in T which contains both x and y . Note that if $u, v \in [x, y]$, then $[u, v] \subset [x, y]$. So if $u, v \in C_j \cap C_k = (C_j \cap T) \cap (C_k \cap T)$, then $[u, v] \subset C_j \cap C_k$ since each set $C_i \cap T$ is of the form $[x_i, y_i]$ (where x_i and y_i are the vertices incident with the edge e_i). $\square$

Lemma 2.4. *If C_i and C_j are basic cycles associated to the edge basis E of the partitioned graph $(G, \{h_i\})$, then at least one of the sets of edges $C_i \cap h_j$ or $C_j \cap h_i$ is empty.*

Proof. Suppose that $e'_j \in C_i \cap h_j$ and $e'_i \in C_j \cap h_i$. Then both of e'_i and e'_j are contained in the connected set $C_i \cap C_j$. So the complement of the edge basis $E \cup \{e'_i, e'_j\} - \{e_i, e_j\}$ contains a cycle, namely $C_i \cup C_j - C_i \cap C_j$. This contradicts Lemma 2.2. $\square$

2.2. Oriented, Partitioned Graphs. An edge basis is said to be oriented if each edge of it is oriented. If $E = \{e_i\}$ is an oriented edge basis of a partitioned graph $(G, \{h_i\})$, then the orientation of each e_i determines an orientation of the corresponding basic cycle C_i^E , and hence of all the edges in $\{h_i\}$. Thus, since $E(G) = \coprod h_i$, E induces an orientation of every edge of G . A partitioned graph is said to be oriented if all of its edges are oriented by some oriented edge basis in this way.

If a partitioned graph G has rank n then there are just 2^n possible orientations of G as the next lemma shows.

Lemma 2.5. *Assume that G is a partitioned graph which is oriented by the oriented edge basis E . If the orientation of G is used to orient any edge basis F of G , then the orientations of G induced by F and E agree.*

Proof. If the basic cycles C_k^E and C_k^F are not equal for some k , then $E \cap h_j \neq F \cap h_j$ for some $j \neq k$. For such a j , replace $e_j \in E \cap h_j$ with $e'_j \in F \cap h_j$ to obtain a new edge basis E' . If $C_k^E \neq C_k^{E'}$, then e'_j is contained in $C_k^E \cap C_k^{E'}$. In this case $C_k^E \cap C_k^{E'} = C_k^E - (C_k^E \cap C_j^E)$ is connected since $C_k^E \cap C_j^E$ is by Lemma 2.3. Since $h_k \subset C_k^E \cap C_k^{E'}$, the orientations of the edges in h_1 determined by E and E' agree. Continuing in this way shows that the orientations determined by E and F of all the edges in h_k , and also in the other h_i , agree. $\square$

A consequence of the previous lemma is that if $f: G \rightarrow G'$ is a blowdown of partitioned graphs and G is oriented, then the orientation which each edge of G' inherits from the corresponding edge of G together give an orientation of G' . When G' has this orientation, we say that f is orientation preserving.

2.3. Colored Graphs. A pointed graph is a graph with one of its vertices designated as a basepoint. Basepoints are denoted by $*$. A colored graph is a pointed, oriented, partitioned graph G together with a bijection between the set $\mathbf{n} = \{1, \dots, n\}$ and the set $\{h_1, \dots, h_n\}$ of edge sets of G . Unless otherwise mentioned, we will always assume that, for each $i \in \mathbf{n}$, this bijection takes i to h_i . We often think of the elements of the set $\mathbf{n}$ as distinct colors and of drawing the colored graph G so that all the edges of G in h_i have color i . Thus the set $\mathbf{n}$ is sometimes called the set of colors of G , and the bijection from $\mathbf{n}$ to the h_i , a coloring.

Colored graphs (really just partitioned graphs) are often drawn using the smooth edge convention. Namely, two edges are in the same edgeset if and only if they meet in a smooth curve at some vertex. For example, partitions of the graphs in Figure 2, except for the last two, are indicated in this way. Alternatively, edges in the same edgeset of a colored graph are drawn with the same type of line, eg. solid, dotted, or finely dotted. Both conventions are often used on the same graph.

For convenience, we require that each vertex of a colored graph has valence at least three, except for the basepoint, which may be bivalent.

Two colored graphs G and G' are said to be equivalent if there is a pointed cellular homeomorphism $f: G \rightarrow G'$ which preserves colors and orientations. Such a homeomorphism is called an equivalence of colored graphs.

Proposition 2.6. *Any two equivalences $G \rightarrow G'$ induce the same bijections $E(G) \rightarrow E(G')$ and $V(G) \rightarrow V(G')$.*

The proof uses the next two lemmas. These are preceded by a definition.

If G is a graph, the number of half-edges incident with a vertex v of G is the integer $|\{e \in E(G) : d_0(e) = v\}| + |\{e \in E(G) : d_1(e) = v\}|$.

Lemma 2.7. *The number of half-edges of the same color incident with any vertex v of a colored graph G is at most two.*

Proof. If three half-edges of the same color k were incident with v , then some subcycle of the basic cycle C_k (with respect to any edge basis) would not contain any edge set of G . $\square$

Lemma 2.8. *If two equivalences $G \rightarrow G'$ of colored graphs agree on a vertex v of G , they agree on all edges of G incident with v .*

This follows from Lemma 2.7 since equivalences preserve colorings and orientations.

Proof of 2.6. Let $v \in V(G)$. Choose an edge path p in G from $*$ to v . Since equivalences preserve $*$, it follows from repeated use of Lemma 2.8 that equivalences agree on the edges and, by continuity, on the (nonbasepoint) vertices of p . Thus any two equivalences induce the same bijection $V(G) \rightarrow V(G')$, and so by Lemma 2.8 the same bijection $E(G) \rightarrow E(G')$. $\square$

2.4. The space X_n . Let R_n be a fixed colored graph of rank n having just one vertex. A marked, colored graph (G, μ) , or just G , is, by definition, a colored graph G together with a pointed homotopy equivalence $\mu : R_n \rightarrow G$. The map μ is called a marking (of G). Two marked, colored graphs (G, μ) and (G', μ') are equivalent if there is an equivalence $f : G \rightarrow G'$ of colored graphs and a homotopy between $f \circ \mu$ and μ' which keeps the basepoint of R_n fixed at all times. We say that (G', μ') is a blowdown of (G, μ) if there is a blowdown $h : G \rightarrow G'$ of the underlying partitioned graphs which preserves basepoints, colors and orientations, and if (G', μ') is equivalent to $(G', h \circ \mu)$.

Marked, colored graphs might be called decorated graphs, so we use $\mathbf{bD}_n$ to denote the poset of equivalence classes of marked, colored graphs of rank n ordered by blowdowns. Specifically, two elements G_1 and G_2 of $\mathbf{bD}_n$ satisfy $G_1 \geq G_2$ if G_2 is a blowdown of G_1 . Here and in the sequel we use the same symbol to denote both a marked, colored graph and its equivalence class in $\mathbf{bD}_n$. A poset map $\mathbf{bD}_n \rightarrow \mathbf{bD}_{n+1}$ is defined by sending $\mu : R_n \rightarrow G$ to $\mu \vee id : R_n \vee R_1 \rightarrow G \vee R_1$ where the color of R_1 is $n + 1$. The limit of the posets $\mathbf{bD}_n$ using these maps is itself a poset which we denote by $\mathbf{bD}$.

Definition 2.9. The space X_n is the geometric realization of the poset $\mathbf{bD}_n$, and X is the direct limit of the X_n as $n \rightarrow \infty$.

We now work towards another definition of X_n which is ultimately given in Proposition 2.14 below.

Lemma 2.10. *The minimal elements of the poset $\mathbf{bD}_n$ are in bijective correspondence with the elements of the group $\text{Aut}(\pi_1(R_n))$.*

Proof. Let (R, μ) be any minimal element of $\mathbf{bD}_n$ so that the graph R has just one vertex. Using the unique (up to homotopy) equivalence $R_n \rightarrow R$ of colored graphs, the marking μ of R may be viewed as a self homotopy equivalence of R_n . These are in bijective correspondence

with the elements of $\text{Aut}(\pi_1(R_n))$ since R_n is an Eilenburg-MacClane space of type $K(F_n, 1)$. $\square$

We now fix an isomorphism from the free group F_n with basis $\{x_1, \dots, x_n\}$ to $\pi_1(R_n) \equiv \pi_1(R_n, *)$. It takes x_i to the (class of the) loop which wraps once around the single edge in h_i in an orientation preserving manner. This isomorphism, along with the previous lemma, gives a bijection

$$(1) \quad \{\text{minimal elements of } \text{b}\mathcal{D}_n\} \longleftrightarrow \text{Aut}(F_n).$$

Each marked, colored graph G determines a full poset $\text{b}G$ of $\text{b}\mathcal{D}$ containing G and all of its blowdowns. As an example, if G has just two vertices, with, say, $h_1 = \{a, b\}$ the only edgeset with > 1 element, then $\text{b}G$ has three elements: G , G_a and G_b . The realization of $\text{b}G$ is the barycentric subdivision of a 1-simplex, with the barycenter corresponding to G . The other two vertices G_a and G_b , when identified with elements of $\text{Aut}(F_n)$, differ by a Whitehead automorphism as we now explain.

We first review the definition of Whitehead automorphisms of the free group F_n with basis $\{x_1, \dots, x_n\}$. If $a \in L = \{x_1^{\pm 1}, \dots, x_n^{\pm 1}\}$ and $a \notin \{x_k, x_k^{-1}\}$, then the Whitehead automorphism $\langle a, x_k \rangle$ takes a to ax_k and fixes those x_i different from a and a^{-1} . In general, if $A \subset L$ with $A \cap \{x_k, x_k^{-1}\} = \emptyset$, then the Whitehead automorphism $\langle A, x_k \rangle = \prod_{a \in A} \langle a, x_k \rangle$. We often write $\langle A, x_k^{-1} \rangle$ for $\langle A, x_k \rangle^{-1}$.

Now assume that the two vertices of G are the basepoint $*$ and v , and that G has edge sets $h_1, \dots, h_n$. Also assume that the basic cycle C_1 is orientated so that $\text{init}(b) = *$. Define a subset A of the letters L of F_n by

$$\begin{aligned} A = \{ & x_i : i \neq 1 \text{ and } \text{term}(e_i) = v \text{ where } e_i \in h_i \} \\ & \cup \{ x_i^{-1} : i \neq 1 \text{ and } \text{init}(e_i) = v \text{ where } e_i \in h_i \}. \end{aligned}$$

Writing fg for the composite first f , then g , we have the following.

Lemma 2.11. *If G_a and G_b are identified with automorphisms of F_n as in (1) above, then $G_b \langle A, x_1^{-1} \rangle = G_a$.*

The omitted proof is not difficult, cf. 3.1.1 of [1].

We will denote the edgepath in $|\text{b}G| \subset X$ from G_b to G_a corresponding to the automorphism $\langle A, x_1 \rangle$ by $b \wedge a$. The symbol $b \wedge a$ could be read as (blow)up b , (blow)down a .

Given $G \in \text{b}\mathcal{D}_n$ with edgesets $h_1, \dots, h_n$, let P_i denote the poset of all nonempty subsets of h_i ordered by inclusion and let $P_1 \times \dots \times P_n$ be the product poset where ordered pairs (s_i) and (s'_i) satisfy $(s_i) \geq (s'_i)$ if $s_i \supseteq s'_i$ for all i . Let $f: P_1 \times \dots \times P_n \rightarrow \text{b}G$ be the poset map which takes $s = (s_i) \in P_1 \times \dots \times P_n$ to the marked colored graph G^s obtained from G by collapsing the edges $\cup_i (h_i - s_i)$ of G . Thus the edges of G^s correspond to the elements of the set $\cup_i s_i$.

Lemma 2.12. *The map $f: P_1 \times \dots \times P_n \rightarrow \text{b}G$ is a poset isomorphism.*

Proof. Surjectivity is clear. To show that f is injective, we first assume that s and t are distinct minimal elements of $P_1 \times \dots \times P_n$ with say $s = (\{e_i\})$ and $t = (\{e'_i\})$. Let $j_1, \dots, j_m$ be the elements of the set $\{i \in \mathbf{n} : e_i \neq e'_i\}$ listed in any order. Then the path in $\text{b}G$ from $f(s)$ to $f(t)$ formed by the sequence of length two edgepaths $e'_{j_1} \wedge e_{j_1}, e'_{j_2} \wedge e_{j_2}, \dots, e'_{j_m} \wedge e_{j_m}$ gives, by repeated use of Lemma 8.3, that $f(t) = f(s) \prod_{i=1}^m \langle A_{j_i}, x_{j_i}^{\pm 1} \rangle$. Here the sets A_{j_i} are

as in the Lemma, and the choices of the powers of each x_{j_i} are dictated by the orientation of G .

Now $\prod_{i=1}^{m-1} \langle A_{j_i}, x_{j_i}^{\pm 1} \rangle \neq \langle A_{j_m}, x_{j_m}^{\mp 1} \rangle$ since, if $x_k^{\pm 1} \in A_{j_m}$, the automorphism $\langle A_{j_m}, x_{j_m}^{\pm 1} \rangle$ takes x_k to a word containing at least one of x_{j_m} or $x_{j_m}^{-1}$ whereas the image of x_k under $\prod_{i=1}^{m-1} \langle A_{j_i}, x_{j_i}^{\pm 1} \rangle$ contains neither x_{j_m} nor $x_{j_m}^{-1}$. Therefore $\prod_{i=1}^m \langle A_{j_i}, x_{j_i}^{\pm 1} \rangle$ is not the identity and so $f(t) \neq f(s)$.

Now assume that $f(s) = f(t)$ for distinct, nonminimal $s, t \in P_1 \times \dots \times P_n$. Choose distinct minimal elements $s' = (\{e_i\})$ and $t' = (\{e'_i\})$ such that $s \geq s'$ and $t \geq t'$, and such that the unique equivalence $f(s) \rightarrow f(t)$ of colored graphs takes each e_i to e'_i . This last condition implies that $f(s') = f(t')$ which contradicts the first part of the proof since $s' \neq t'$. $\square$

If $\mathcal{P}$ is any poset, we let $|\mathcal{P}|$ denote the geometric realization of $\mathcal{P}$. Since $|P_1 \times \dots \times P_n|$ is homeomorphic to $|P_1| \times \dots \times |P_n|$, and since $|P_i|$ is simplicially isomorphic to the first barycentric subdivision of $\sigma^{\|h_i\|}$, the standard simplex of dimension $\|h_i\| = |h_i| - 1$, the previous lemma gives the following.

Proposition 2.13. *Let $G \in \mathbf{bD}_n$. Then the space $|bG|$ is homeomorphic to the disk $\sigma^{\|h_i\|} \times \dots \times \sigma^{\|h_n\|}$ of dimension n .*

The following is a consequence of Lemma 2.12 and Proposition 2.13.

Proposition 2.14. *The space X_n is homeomorphic to the n -dimensional CW -complex having one k -cell $|bG|$ corresponding to each $G \in \mathbf{bD}_n$ with $k + 1$ vertices. The attaching maps of the cells are as indicated by 2.12 and 2.13.*

Proof. For each $G \in \mathbf{bD}_n$, there is a natural inclusion $|bG| \hookrightarrow X_n$. These inclusions take the interiors of the disks $|bG|$ to pairwise disjoint sets which cover X_n . $\square$

We will make use of both this CW -structure and the simplicial structure of X_n . Any mention of the cells of X_n or its skelata indicates that we have the CW -structure in mind. The simplicial structure is inferred by reference to simplices of X_n .

We conclude this section with some remarks concerning the space X_n . Most of these will not be used in the sequel.

According to Proposition 2.13, the cells of X_n of dimension k have different “shapes”, one for each partition of k as a sum of positive integers. It may be helpful to think of X_n as the realization of a “generalized simplicial complex”. We say generalized here since the building blocks in degree k —one for each marked colored graphs with $k + 1$ vertices—are not just the usual k -simplices, but all the shapes corresponding to the partitions of k . Of course, taking the “barycentric subdivision” of these shapes gives a simplicial set. Its realization is the description of X_n given in definition 2.9.

Since the shapes are products of simplices, perhaps X_n is the realization of a multisimplicial set in some natural way. It seems that only a subcomplex is, as we now explain. Let $\mathbf{bD}'_n$ be the full subposet of $\mathbf{bD}_n$ consisting of the marked colored graphs of rank n whose edgesets can be canonically linearly ordered. Note that every edgeset of a colored graph is cyclically ordered in a canonical way by any edgepath containing it. If, for instance, such an edgepath passes through the basepoint, then the edges of the edgeset are canonically linearly ordered as well.

Let $Y = Y_{\bullet \dots \bullet}$ be the n -simplicial set with $Y_{i_1 \dots i_n}$ the marked colored graphs in $\mathbf{bD}'_n$ with $1 + i_j$ edges of color j , for $1 \leq j \leq n$. The m^{th} face map $Y_{i_1 \dots i_j \dots i_n} \longrightarrow Y_{i_1 \dots i_j - 1 \dots i_n}$ is defined by

blowing down the m^{th} edge of the edgeset h_j of each $G \in Y_{i_1 \dots i_j \dots i_n}$. To define the m^{th} edge, we identify the linearly ordered edges in each h_j with the set $\{0, 1, \dots, i_j\}$ in the obvious way. Degeneracies can be defined by introducing bivalent vertices. The realization of Y describes a large CW-subcomplex of X_n . For instance, $|Y|$ contains the entire 2-skeleton of X_n except for those 2-cells corresponding to theta graphs. Theta graphs are those colored graphs equivalent to the colored graph shown on the left of Figure 6 for some choice of colors x and y . The image of the composite $|Y| \hookrightarrow X_n \rightarrow V$ agrees with that of the map $X_n \rightarrow V$ but representatives of certain elements of $\pi_2(V)$, namely exotic elements of $K_3(\mathbb{Z})$, lift to X_n but not to $|Y|$. Here $V = V(\text{GL}(\mathbb{Z}))$ is the space defined by Volodin in [12] and discussed in §2.9 below.

Theta graphs are also noteworthy for the following reason. Colored graphs and partition-preserving maps between them form a Waldhausen category $\mathcal{C}$ (as discussed in §1.3 of [13]) with weak equivalences those morphisms which induce bijections on the set of colors and with injections as cofibrations. If every colored graph G admitted a filtration $* \twoheadrightarrow G_1 \twoheadrightarrow \dots \twoheadrightarrow G_n = G$ by cofibrations with all subquotients G_i/G_{i-1} roses, then essentially Waldhausen's additivity theorem would give a homotopy equivalence $\text{pS}_\bullet \mathcal{C} \simeq \Omega^\infty S^\infty$, where p denotes the weak equivalences just mentioned. But theta graphs do not admit such filtrations. In fact, $\Omega^\infty S^\infty$ splits off $\text{pS}_\bullet \mathcal{C}$ with theta graphs contributing to nontrivial homology of the complementary factor.

We note that the realization of the simplicial set $[n] \mapsto Y_{0 \dots 0 n}$ is the disjoint union of simplicial copies of $\mathbb{R}^{2n-2}$. This is discussed in §4 below from a slightly different viewpoint. The free abelian group $\mathbb{Z}^{2n-2} \subset \text{Aut}(F_n)$ generated by $\langle x_1, x_n \rangle, \dots, \langle x_{n-1}, x_n \rangle$ and $\langle x_1^{-1}, x_n \rangle, \dots, \langle x_{n-1}^{-1}, x_n \rangle$ acts freely on each copy. Also, the simplicial map $|Y_{0 \dots 0 \bullet}| \hookrightarrow X_n \rightarrow S\mathbb{A}_n$ is injective on each copy. In fact, it is injective on considerably larger subcomplexes of X_n . Here $S\mathbb{A}_n$ is a version of the sphere complex defined, for instance, in [5].

Lemma 2.11 implies that each (oriented) 1-cell of X_n corresponds to a Whitehead automorphism. According to Nielson's theorem, cf. Cor. 3.6 of [7], these generate the index two subgroup $\text{Aut}_+(F_n)$ of $\text{Aut}(F_n)$ consisting of all automorphisms which abelianize to elements of $\text{GL}_n(\mathbb{Z})$ with determinant 1. Thus the space X_n has two connected components. The 1-skeleton of the component containing the identity is the Caley graph of $\text{Aut}_+(F_n)$ with the Whitehead automorphisms as generators.

Let $G \in \text{b}\mathcal{D}$. The induced CW-structure on $|bG|$ is denoted by $|G|$, and is just the product CW-structure given by Proposition 2.13 (assuming the CW-structure of the standard simplex $\sigma^{|h_i|}$ is just its simplicial structure). Elaborating on the middle paragraph of the proof of Lemma 2.12 shows that the 1-skeleton of $|G|$ is a geodesically convex subset of the Caley graph of $\text{Aut}_+(F_n)$ mentioned above. This, along with the next two sections, suggests an algebraic definition of the space X_n which does not use colored graphs.

2.5. The Partial Ordering determined by a Colored Graph. Each colored graph $(G, \{h_i\})$ of rank n defines an ordering of the set $\mathbf{n} = \{1, \dots, n\}$ of colors as follows. If $b, r \in \mathbf{n}$, then $b >_G r$ or just $b > r$ if $b \neq r$ and if there is some edge basis E of G such that $C_b^E \cap h_r \neq \emptyset$.

Note that $b > r$ if and only if there exists an edge basis $E = \{e_1, \dots, e_n\}$ and one additional edge $e'_r \in h_r$ such that the edge $e_b \in E$ joins the two vertices of the graph obtained by blowing down all the edges of G except for those in $E \cup \{e'_r\}$.

The following two lemmas will show that the ordering $>$ is antisymmetric and transitive. Thus $\geq$ is a partial ordering of the set $\mathbf{n}$.

Lemma 2.15. *If $b > r$, then $C_r^E \cap h_b = \emptyset$ for every edge basis E .*

It follows from this that both $b > r$ and $r > b$ cannot occur.

Proof of 2.15. Let $\mathcal{E}$ be the set of edgebases defined by $E \in \mathcal{E}$ if $C_b^E \cap h_r \neq \emptyset$. Note that if $E \in \mathcal{E}$, then $C_r^E \cap h_b = \emptyset$ by Lemma 2.4.

Let $F = \{f_1, \dots, f_n\}$ be any edge basis with $f_i \in h_i$. We will show that there is an $E \in \mathcal{E}$ such that $C_r^E \cap F = \{f_r\}$. Then $C_r^E = C_r^F$ and so $C_r^F \cap h_b = C_r^E \cap h_b = \emptyset$, which will complete the proof.

So assume that $f_i \in C_r^E \cap F$ where $i \neq r$ and $E \in \mathcal{E}$. If $f_i \in C_b^E$, then $C_i^E \cap h_b = \emptyset$ by Lemma 2.4. So letting E_1 be the edge basis $E \cup \{f_i\} - \{e_i\}$, we have $C_b^{E_1} = (C_b^E \cup C_i^E) - (C_b^E \cap C_i^E)$. Therefore $E_1 \in \mathcal{E}$ since $C_i^E \cap h_r = \emptyset$ by Lemma 2.4 once more. And if $f_i \notin C_b^E$, then $C_b^{E_1} = C_b^E$ again giving $E_1 \in \mathcal{E}$. Note that this shows that if $E \in \mathcal{E}$ and $|C_r^E \cap F| > 1$, then there is an $E_1 \in \mathcal{E}$ such that $|E_1 \cap F| > |E \cap F|$. So now either $|C_r^{E_1} \cap F| = 1$ which would complete the proof or, if not, there is, as just noted, an $E_2 \in \mathcal{E}$ with $|E_2 \cap F| > |E_1 \cap F|$. Repeating this as needed eventually gives an $E \in \mathcal{E}$ with $|C_r^E \cap F| = 1$ since $C_r^F \cap F = \{f_r\}$. $\square$

Lemma 2.16. *If $b > r$ and $r > g$, then $C_b^E \cap h_r \neq \emptyset$ and $C_b^E \cap h_g \neq \emptyset$ for some edge basis E .*

So the relation $>_G$ is transitive.

Proof of 2.16. Assume $r > g$ and let $F = \{f_i\}$ be an edge basis with $f_i \in h_i$ such that $C_r^F \cap h_g \neq \emptyset$. Also assume $b > r$. Then as in the proof of Lemma 2.15, there is an edge basis E with $C_b^E \cap h_r \neq \emptyset$ and $C_r^E \cap F = \{f_r\}$. If $C_b^E \cap h_g \neq \emptyset$ the lemma is proved, so assume $C_b^E \cap h_g = \emptyset$. Pick an edge $e'_r \in C_b^E \cap h_r$, and let $E' = E \cup \{e'_r\} - \{e_r\}$. Then $C_b^{E'} = (C_b^E \cup C_r^E) - (C_b^E \cap C_r^E)$, which contains an edge of h_g since $C_r^E = C_r^F$ does, and C_b^E does not. Also $e_r \in C_b^{E'}$ since $e_r \notin C_b^E$. $\square$

2.6. Triangular subgroups. Let G be a colored graph. The previous section shows that the Whitehead automorphisms corresponding to the 1-cells of $|G|$ are contained in a triangular subgroup T_n^σ of $\text{Aut}(F_n)$ which we are about to define.

Let σ be any partial ordering of the set $\mathbf{n}$. In particular, $\sigma = \sigma(G)$ could be the order determined by G . Let S_n^σ be the subgroup of $\text{Aut}(F_n)$ generated by the set $\{\langle x_i, x_j \rangle : i \overset{\sigma}{>} j\} \cup \{\langle x_i^{-1}, x_j \rangle : i \overset{\sigma}{>} j\}$, and let C_n be the subgroup generated by the set $\{\langle x_i, x_i^{-1}; x_j \rangle : i \neq j\}$. Elements of C_n include automorphisms which conjugate any one basis element by any other and leave all other basis elements fixed. The triangular subgroup T_n^σ mentioned above is, by definition, the one generated by $S_n^\sigma \cup C_n$. Elements of T_n^σ are referred to as (σ) -triangular automorphisms, while those of S_n^σ are called strictly triangular.

Lemma 2.17. *The group T_n^σ of triangular automorphisms is torsion free.*

Proof. Let $a: \text{Aut}(F_n) \rightarrow \text{GL}_n(\mathbb{Z})$ be the natural homomorphism. Let K be the kernel of a restricted to T_n^σ so that the sequence

$$1 \rightarrow K \rightarrow T_n^\sigma \rightarrow a(T_n^\sigma) \rightarrow 1$$

is exact. K is contained in the torsion free subgroup $\text{IA}(F_n)$ of $\text{Aut}(F_n)$. Also, $a(T_n^\sigma)$ is a torsion free subgroup of $\text{GL}_n(\mathbb{Z})$. If σ is the standard ordering, this subgroup consists of all upper triangular matrices with diagonal entries equal to 1, and, for other σ , is isomorphic to it. Thus both K and $a(T_n^\sigma)$ are torsion free and so T_n^σ is also. $\square$

2.7. The action of $\text{Aut}(F_n)$ on X_n . Let $f \in \text{Aut}(F_n)$ and $G = (G, \mu) \in \text{b}\mathcal{D}_n$. Assume f corresponds to the minimal element $f': R_n \rightarrow R_n$ of $\text{b}\mathcal{D}_n$. Denote by $f \cdot G$ or fG the element $(G, f'\mu)$ of $\text{b}\mathcal{D}_n$. This defines a left action of $\text{Aut}(F_n)$ on $\text{b}\mathcal{D}_n$ by poset isomorphisms. Thus there is a simplicial left action of $\text{Aut}(F_n)$ on X_n . This action is also cellular. Moreover, we have the following.

Proposition 2.18. *The action of $\text{Aut}(F_n)$ on X_n is free.*

Proof. Identify $X_n^{(0)}$, the 0-skeleton of X_n , with $\text{Aut}(F_n)$. Then the action of any $f \in \text{Aut}(F_n)$ on $X_n^{(0)}$ amounts to multiplication by f on the left. So $\text{Aut}(F_n)$ clearly acts freely on $X_n^{(0)}$. Therefore suppose $f \in \text{Aut}(F_n)$ fixes an interior point of some cell σ of X_n . Then f permutes the 0-cells of σ and so $f^n = \text{id}$ for some n . Let $v \in \text{Aut}(F_n)$ be any 0-cell of σ . Since fv is also a 0-cell of σ , there is a triangular automorphism t such that $fv = vt$. Therefore $t^n = v^{-1}f^n v = \text{id}$ which, by the preceding lemma, implies that $t = \text{id}$. So $f = \text{id}$ also. $\square$

2.8. The space $\tilde{X}_n$. Let $X_{*,n}$ be the component of X_n containing the vertex corresponding via the bijection (1) to the identity in $\text{Aut}(F_n)$. We often work with the universal cover $\tilde{X}_n$ of the space $X_{*,n}$. The covering $\tilde{X}_n$ can be defined using marked colored graphs just as X_n is, except that the markings involve elements of the group StN_n , rather than those of $\text{Aut}(F_n)$. Details follow, beginning with basic facts about the group StN_n .

Let $B^\pm = \{x_1, \dots, x_n\} \cup \{x_1^{-1}, \dots, x_n^{-1}\}$ where $\{x_1, \dots, x_n\}$ is a basis for F_n . The group StN_n is defined by the presentation having one generator $\langle\langle x, y \rangle\rangle$ for all $x, y \in B^\pm$ with $x \neq y^{\pm 1}$, and the relations

- (R0): $\langle\langle x, y^{-1} \rangle\rangle = \langle\langle x, y \rangle\rangle^{-1}$
- (R1): $[\langle\langle x, y \rangle\rangle, \langle\langle x', y' \rangle\rangle] = 1$ if $y^{\pm 1} \neq x' \neq x \neq (y')^{\pm 1}$
- (R2): $[\langle\langle x, y \rangle\rangle, \langle\langle y, z \rangle\rangle] = \langle\langle x, z \rangle\rangle$, if $x \neq z^{\pm 1}$
- (R3): $\langle\langle y, x \rangle\rangle \langle\langle x^{-1}, y \rangle\rangle \langle\langle y^{-1}, x^{-1} \rangle\rangle = \langle\langle y^{-1}, x^{-1} \rangle\rangle \langle\langle x, y^{-1} \rangle\rangle \langle\langle y, x \rangle\rangle$.

We sometimes will use the notation L_{ij} for $\langle\langle x_i^{-1}, x_j^{-1} \rangle\rangle$ and R_{ij} for $\langle\langle x_i, x_j^{-1} \rangle\rangle$.

Two elements of StN_n which we will often use are $w_{ij}^L = \langle\langle x_i, x_j \rangle\rangle \langle\langle x_j^{-1}, x_i \rangle\rangle \langle\langle x_i^{-1}, x_j^{-1} \rangle\rangle = R_{ij}^{-1} L_{ji}^{-1} L_{ij}$ and $w_{ij}^R = \langle\langle x_i^{-1}, x_j^{-1} \rangle\rangle \langle\langle x_j, x_i^{-1} \rangle\rangle \langle\langle x_i, x_j \rangle\rangle = L_{ij} R_{ji} R_{ij}^{-1}$.

There is a homomorphism $\phi: StN_n \rightarrow \text{Aut}(F_n)$ defined by $\phi(\langle\langle x, y \rangle\rangle) = \langle x, y \rangle$. Gersten showed in [2] that the kernel of ϕ , which we denote by $KN_{n,2}$, has order two.

With the following lemma, many of the facts about marked, colored graphs discussed above carry over to StN -marked, colored graphs. Such graphs are defined after its proof.

Lemma 2.19. *Let $T^\sigma(F_n)$ be any triangular subgroup of $\text{Aut}(F_n)$. Then there is an injection $T^\sigma(F_n) \rightarrow StN_n$ induced by $\langle x_i, x_j \rangle \mapsto \langle\langle x_i, x_j \rangle\rangle$, $\langle x_i^{-1}, x_j \rangle \mapsto \langle\langle x_i^{-1}, x_j \rangle\rangle$, and $\langle \{x_i, x_i^{-1}\}, x_j \rangle \mapsto \langle\langle x_i^{-1}, x_j \rangle\rangle \langle\langle x_i, x_j \rangle\rangle$.*

Proof. Let $\langle G_{TF} \rangle$ be the free group on the set $S_n^\sigma \cup C_n$ of defining generators of T_n^σ as in §2.6, and let $\langle G_{T\mathbb{Z}} \rangle$ be the free group on the image of this set in $\text{GL}_n(\mathbb{Z})$. We take the horizontal map $\langle G_{TF} \rangle \rightarrow StN_n$ in the diagram

$$\begin{array}{ccccc}
R_F & \longrightarrow & \mathbb{Z}_2 & & \\
\downarrow & \searrow & \downarrow & \searrow \approx & \\
& & R_{\mathbb{Z}} & \longrightarrow & \mathbb{Z}_2 \\
\downarrow & & \downarrow & & \downarrow \\
\langle G_{TF} \rangle & \longrightarrow & StN_n & & \\
\downarrow & \searrow & \downarrow \phi & \searrow & \downarrow \\
& & \langle G_{T\mathbb{Z}} \rangle & \longrightarrow & St_n(\mathbb{Z}) \\
\downarrow & & \downarrow & & \downarrow \\
T_n^\sigma & \longrightarrow & \text{Aut}(F_n) & & \\
\downarrow & \searrow & \downarrow & \searrow & \downarrow \\
& & T_n^\sigma(\mathbb{Z}) & \longrightarrow & \text{GL}_n(\mathbb{Z})
\end{array}$$

to be induced as in the statement of the lemma. The horizontal map $\langle G_{T\mathbb{Z}} \rangle \rightarrow St_n(\mathbb{Z})$ is induced similarly. Each of the four columns of the diagram is part of a short exact sequence. The groups R_F and $R_{\mathbb{Z}}$ are defined as the kernels of the relevant maps, while one of the $\mathbb{Z}_2$'s follows from the computation of $K_2(\mathbb{Z})$, the other from the nonabelian analog. It follows from a Lemma in [12] that the homomorphism $R_{\mathbb{Z}} \rightarrow \mathbb{Z}_2$ in the top horizontal square of the diagram is trivial. Since this square commutes, the homomorphism $R_F \rightarrow \mathbb{Z}_2$ in it is also trivial. Thus the injective map $T^\sigma(F_n) \approx \langle G_{TF} \rangle / R_F \rightarrow \text{Aut}(F_n)$ factors through the group StN_n . $\square$

A StN-marked, colored graph (of rank n) is, by definition, a marked, colored graph G (of rank n) together with a compatible choice of lifts of the 0-cells of $|G|$ from $\text{Aut}(F_n)$ to StN_n . Recall that $|G|$ denotes the CW -structure of $|bG|$ discussed just after the bijection (1) near the beginning of section 2.4. As usual, we are identifying the elements of $|G|^{(0)}$ with elements of $\text{Aut}(F_n)$ using this bijection. A lift of $v \in |G|^{(0)} \subset \text{Aut}(F_n)$ then means one of the two preimages of v under the map $\phi: StN_n \rightarrow \text{Aut}(F_n)$. Lifts are compatible if whenever v and w , respectively, are in $|G|^{(0)}$, their lifts $\tilde{v}$ and $\tilde{w}$, respectively, satisfy $\tilde{v} \prod_i \langle A_i, a_i \rangle^{\pm 1} = \tilde{w}$ in the group StN_n , where $\prod_i \langle A_i, a_i \rangle^{\pm 1}$ corresponds to the shortest path in $|G|^{(1)}$ from v to w . Here we are identifying (oriented) 1-cells of $|G|$ with Whitehead automorphisms in $T_n^{\sigma(G)}$ and then, using Lemma 2.19, with elements of StN_n . This lemma and section 2.5 shows that a compatible choice of lifts of the elements of $|G|^{(0)}$ always exists and is determined by a choice of lift of any one of the elements.

Two StN-marked, colored graphs are equivalent if, when their StN-markings are converted to just markings using the homomorphism $\phi: StN_n \rightarrow \text{Aut}(F_n)$, they are equal as elements of $b\mathcal{D}$, and if their StN-markings agree. Blowdowns of StN-marked, colored graphs inherit StN-markings in a natural way. Let $\widetilde{b\mathcal{D}}_n$ be the poset of equivalence classes of StN-marked, colored graphs ordered by blowdowns. Note that since the kernel of the map $\phi: StN_n \rightarrow \text{Aut}(F_n)$ has order two, there is a two-to-one poset map $\phi_{\mathcal{P}}: \widetilde{b\mathcal{D}}_n \rightarrow b\mathcal{D}_n$.

Let $\tilde{X}_n$ be the geometric realization of the poset $\widetilde{b\mathcal{D}}_n$. The space $\tilde{X}_n$ has a CW -structure analogous to that of X_n . Specifically, each $\tilde{G} \in \widetilde{b\mathcal{D}}_n$ determines a full subposet $b\tilde{G}$ of $\widetilde{b\mathcal{D}}_n$ consisting of $\tilde{G}$ and all of its blowdowns. The proof of Lemma 2.12, along with Lemma 2.19, implies that $\phi_{\mathcal{P}}|_{b\tilde{G}}: b\tilde{G} \rightarrow bG$ is a poset isomorphism. Thus, the proof of Proposition 2.14 essentially shows that the indicated CW -structure of $\tilde{X}_n$ consists of one k -cell $|b\tilde{G}|$ for each

$\tilde{G} \in \widetilde{\mathbf{bD}}_n$ with $k + 1$ vertices. Just as with the space X_n , mention of the cells of $\tilde{X}_n$ or any of its skelta indicates that we are working with the *CW*-structure of $\tilde{X}_n$, while mention of its simplices or vertices indicates the simplicial structure.

The poset map $\phi_{\mathcal{P}}: \widetilde{\mathbf{bD}}_n \rightarrow \mathbf{bD}_n$ induces a simplicial and cellular map $\phi_X: \tilde{X}_n \rightarrow X_n$ of spaces with image $X_{*,n}$. These maps are both denoted by ϕ , with subscripts that are usually omitted, since, when suitably restricted, they both agree with the homomorphism $\phi: StN_n \rightarrow \text{Aut}(F_n)$ via the bijections

$$\begin{aligned} \{\text{minimal elements of } \widetilde{\mathbf{bD}}_n\} &\leftrightarrow \tilde{X}_n^{(0)} \leftrightarrow StN_n \quad \text{and} \\ \{\text{minimal elements of } \mathbf{bD}_n\} &\leftrightarrow X_n^{(0)} \leftrightarrow \text{Aut}(F_n). \end{aligned}$$

If $\tilde{v}_1$ and $\tilde{v}_2$ are vertices of $\tilde{X}_n$ which satisfy $\phi(\tilde{v}_1) = v = \phi(\tilde{v}_2)$ then the closed stars of $\tilde{v}_1$ and $\tilde{v}_2$ in $\tilde{X}_n$ are disjoint since their vertices are. Also, each closed star is mapped homeomorphically by $\phi: \tilde{X}_n \rightarrow X_n$ to the closed star of v in X_n . Thus, since X_n is covered by the open stars of its vertices, the map $\phi: \tilde{X}_n \rightarrow X_n$ is a covering of its image X_n^{id} . The next lemma shows that it is the universal covering.

Lemma 2.20. *The space $\tilde{X}_n$ is simply-connected for $n \geq 3$.*

Proof. For each (oriented) 1-cell e of X_n which corresponds to a Whitehead automorphism of the form $\langle A, x_i \rangle$ where $|A| = 1$, identify both (oriented) lifts of e in $\tilde{X}_n$ with the generator $\langle\langle A, x_i \rangle\rangle$ of the group StN_n . Using these identifications, the defining relations of StN_n can be realized by 2-cells of $\tilde{X}_n$. Figure 1 shows how to do this for the relations (R3). Each of the three 2-cells in the figure correspond to the colored graph drawn within it. Edges of color j are dotted, and those of color i solid. Arrows near the boundary of each 2-cell indicate the boundaries of these cells. For instance, the left-hand 1-cell of the boundary of the square cell in the middle corresponds to the θ -shaped graph with its edge c blown down.

The various relations (R1) can be realized similarly using the colored graphs numbered 3, 4, 5, and 6 in Figure 35 below. In case $y = y'$ in relation (R1), then a pair of 2-cells, both corresponding to either the colored graph numbered 5 or the one numbered 6, are needed. Otherwise, a single 2-cell corresponding to one of the colored graphs 3 or 4 will do. Finally, the relations (R2) are realized by pairs of 2-cells, with one cell of each pair corresponding to the colored graph numbered 2 in Figure 35, and the other to the one numbered 5.

To complete the proof, we show that any edgeloop of $\tilde{X}_n$ can be deformed to one whose edges were identified in the first paragraph with the defining generators of StN_n . For this let $\tilde{e}$ be any 1-cell of $\tilde{X}_n$. Assume that $\tilde{e}$ maps to the edge of X_n which corresponds to the Whitehead automorphism $\langle A, x_i \rangle$. If $|A| = m > 1$, then the edgepath $\tilde{e}$ can be deformed, rel endpoints, to an edgepath whose edges correspond to the generators $\langle\langle a, x_i \rangle\rangle$, $a \in A$ of StN_n . In case $|A| = m = 2$ a cell corresponding to one of the colored graphs numbered 5 or 6 in Figure 35 facilitate such a deformation. In general, $m - 1$ such cells are needed. $\square$

We mention that, since the kernel of the homomorphism $\phi: StN_n \rightarrow \text{Aut}(F_n)$ is $KN_{2,n}$, the component X_n^{id} is homeomorphic to the quotient of $\tilde{X}_n$ by the action of the $KN_{2,n}$. So it follows from 2.20 that $\pi_1(X_n^{id}) \approx KN_{2,n}$.

2.9. The space X_n^{alg} . In this section we define a space X_n^{alg} which is, to some (undetermined) extent, an algebraic version of the space X_n . We will also define maps $X_n \rightarrow X_n^{\text{alg}}$ and $X_n^{\text{alg}} \rightarrow V(GL_n(\mathbb{Z}))$. Here $V(GL_n(\mathbb{Z}))$ is a space (denoted by $V(GL_n(\mathbb{Z}), \{T_n^\sigma(\mathbb{Z})\})$ in




FIGURE 1. Realizing the relation $w_{ij}^R = w_{ij}^L$ in $\tilde{X}_n$.

[11]) whose limit $V(GL(\mathbb{Z})) = \lim_{n \rightarrow \infty} V(GL_n(\mathbb{Z}))$ has homotopy groups the K -groups of $\mathbb{Z}$. The space $V(GL(\mathbb{Z}))$ was first defined by Volodin in [12] and Hatcher, but we follow Suslin's description in [11] of Volodin's work. Indeed much of this section is taken from or is similar to §1 of [11].

The definition of X_n^{alg} is similar to that of $V(GL_n(\mathbb{Z}))$, and both are essentially special instances of the following definitions. Let G be a group and $\{G_i\}$ a collection of subgroups of G . The simplices of the simplicial complex $V(G, \{G_i\})$ are, by definition, certain subsets of G . Specifically, $\sigma = \{g_0, \dots, g_p\}$ is a p -simplex of $V(G, \{G_i\})$ if the corresponding set $\{g_i g_j^{-1} : g_i, g_j \in \sigma\}$ of products is contained in one of the subgroups G_i . We also use $V(G, \{G_i\})$ to denote the geometric realization of the simplicial complex $V(G, \{G_i\})$. A space and simplicial complex $V_{\text{op}}(G, \{G_i\})$ is defined just as $V(G, \{G_i\})$ is, but with $g_i^{-1} g_j$ in place of $g_i g_j^{-1}$.

Now let T_n^σ be a triangular subgroup of $\text{Aut}(F_n)$. Such subgroups were defined in section 2.6. As in [11], let $T_n^\sigma(\mathbb{Z})$ denote the image of T_n^σ under the abelianizing homomorphism $S: \text{Aut}(F_n) \rightarrow \text{GL}_n(\mathbb{Z})$. Thus, if σ is the standard ordering of the set $\mathbf{n}$, the subgroup $T_n^\sigma(\mathbb{Z})$ of $\text{GL}_n(\mathbb{Z})$ consists of all upper triangular matrices with ones on the diagonal. Volodin defined $V(GL_n(\mathbb{Z}))$ as $V(GL_n(\mathbb{Z}), \{T_n^\sigma(\mathbb{Z})\})$ where σ ranges over all partial orders of the set $\mathbf{n}$. Similarly, we define the space X_n^{alg} by

$$X_n^{\text{alg}} = V_{\text{op}}(\text{Aut}(F_n), \{T_n^\sigma\}),$$

again with σ ranging over all partial orders of $\mathbf{n}$. The reason for the subscript op is that we are using the convention that the composite fg means first f , then g , if $f, g \in \text{Aut}(F_n)$ and, the opposite order, first g , then f , if $f, g \in GL_n(\mathbb{Z})$.

The abelianizing map S induces a simplicial map $X_n^{\text{alg}} \rightarrow V(GL_n(\mathbb{Z}))$. Also, letting

$$V(St_n(\mathbb{Z})) = V(St_n(\mathbb{Z}), \{T_n^\sigma(\mathbb{Z})\}) \quad \text{and} \quad \tilde{X}_n^{\text{alg}} = V_{\text{op}}(StN_n, \{T_n^\sigma\}),$$

a simplicial map $\tilde{X}_n^{\text{alg}} \rightarrow V(St_n(\mathbb{Z}))$ is induced by the homomorphism $StN_n \rightarrow St_n(\mathbb{Z})$. Here we are using Lemma 2.19 to identify T_n^σ with its image in StN_n .

The group StN_n acts freely on $\tilde{X}_n^{\text{alg}}$ by left multiplication, cf. Proposition 2.18. The quotient $\tilde{X}_n^{\text{alg}}/KN_{2,n}$ is homeomorphic to X_n^{alg} . Since the defining relations for the group

StN are triangular, the space $\tilde{X}_n^{\text{alg}}$ is simply connected. So it is the universal cover of $X_{*,n}^{\text{alg}}$. Here $X_{*,n}^{\text{alg}} = V_{op}(\text{Aut}_+(F_n), \{T_n^\sigma\})$ is the component of X_n^{alg} containing the 0-simplex $\{id\}$.

Letting $n \rightarrow \infty$ we obtain stable versions of the above spaces, complexes, groups, actions and maps. Stable objects are typically denoted by deleting the subscript n from the notation for the corresponding unstable object.

The following Proposition is the main reason for introducing the complex X_n^{alg} . We conjecture that it's true with $\tilde{X}$ in place of $\tilde{X}^{\text{alg}}$.

Proposition 2.21. *The group StN acts trivially on both the homology and the homotopy groups of $\tilde{X}^{\text{alg}}$.*

The proof will make use of cartesian products so, following Suslin [11], we introduce a simplicial set $\tilde{W}_n$, with limit $\tilde{W}$, whose geometric realization is homotopy equivalent to $\tilde{X}_n^{\text{alg}}$. The simplicial set $\tilde{W}_n$ has a p -simplex $(g_0, \dots, g_p)$ for each *sequence* $g_0, \dots, g_p$ of elements of StN_n such that all $g_i^{-1}g_j$ are in one of the triangular subgroups T_n^σ . The i th face, respectively degeneracy, of the simplex $(g_0, \dots, g_p)$ is defined as usual by deleting, respectively repeating, g_i .

Lemma 2.22. *If $x \in StN_{n+1}$, then the maps $u_n: \tilde{W}_n \rightarrow \tilde{W}_{n+1}$ and $x \cdot u_n$ are homotopic.*

This shows that Proposition 2.21 is true if, in its statement, X^{alg} is replaced with $\tilde{W}$. Since the obvious StN -equivariant map $\tilde{W} \rightarrow \tilde{X}^{\text{alg}}$ is a homotopy equivalence, the Proposition is true as stated.

Proof of Lemma 2.22. Because of defining relations R2, the group StN_{n+1} is generated by elements of the form $\langle\langle x_i, x_{n+1} \rangle\rangle$, $\langle\langle x_i^{-1}, x_{n+1} \rangle\rangle$, $\langle\langle x_{n+1}, x_i \rangle\rangle$, and $\langle\langle x_{n+1}^{-1}, x_i \rangle\rangle$. We first assume that $x \in StN_{n+1}$ has one of these forms and define a homotopy between u_n and $x \cdot u_n$ by taking the simplex

$$(\underbrace{0, \dots, 0}_s, \underbrace{1, \dots, 1}_t) \times (\alpha_1, \dots, \alpha_{s+t})$$

of $I \times \tilde{W}_n$ to the simplex

$$\tau = (\alpha_1, \dots, \alpha_s, x\alpha_{s+1}, \dots, x\alpha_{s+t})$$

of $\tilde{W}_{n+1}$. To see that τ is a simplex of $\tilde{W}_{n+1}$, we first mention that one can check, by using the defining relations for StN_n and considering a number of cases (or by just using marked, colored graphs) that if $x = \langle\langle x_i^{\pm 1}, x_{n+1} \rangle\rangle^{\pm 1}$ and $\alpha \in StN_n$, then $\alpha^{-1}x\alpha$ is in the subgroup of StN_{n+1} generated by

$$\{\langle\langle x_i, x_{n+1} \rangle\rangle : i \in \mathbf{n}\} \cup \{\langle\langle x_i^{-1}, x_{n+1} \rangle\rangle : i \in \mathbf{n}\} \cup \{\langle\langle x_{n+1}, x_i \rangle\rangle \langle\langle x_{n+1}^{-1}, x_i \rangle\rangle : i \in \mathbf{n}\}.$$

And if $x = \langle\langle x_{n+1}^{\pm 1}, x_i \rangle\rangle^{\pm 1}$, then $\alpha^{-1}x\alpha$ is in the subgroup generated by

$$\{\langle\langle x_{n+1}, x_i \rangle\rangle : i \in \mathbf{n}\} \cup \{\langle\langle x_{n+1}^{-1}, x_i \rangle\rangle : i \in \mathbf{n}\}.$$

Now let σ be the ordering of $\mathbf{n}$ such that all the $\alpha_i^{-1}\alpha_j$ are in T_n^σ . Writing $\alpha_i^{-1}x\alpha_j$ as $(\alpha_i^{-1}x\alpha_i)(\alpha_i^{-1}\alpha_j)$, we see that τ is a simplex of $\tilde{W}_{n+1}$ where the relevant ordering of the set $\mathbf{n} + \mathbf{1}$ is the ordering σ with $n + 1$ added as either a maximal or minimal element.

Now assume $y \in StN_{n+1}$ is of one of the forms mentioned above. Let h denote the image of the homotopy between u_n and $x \cdot u_n$. Then the subcomplex $y \cdot h$ gives a homotopy between

$y \cdot u_n$ and $y \cdot (x \cdot u_n) = yx \cdot u_n$. Since $y \cdot u_n$ is homotopic to u_n by the first part of the proof, the lemma is true for $yx \in StN_{n+1}$. Continuing in this way proves the lemma for arbitrary elements of StN_{n+1} . $\square$

2.10. The map $\mu: X \rightarrow X^{\text{alg}}$. By the (first) barycentric subdivision poset $\text{Simp}(C)$ of a simplicial complex C , we mean the poset of simplices of C ordered by inclusion. The realization of $\text{Simp}(C)$ is often called the barycentric subdivision of C and is homeomorphic to the realization of C . We define a poset map $\mu'_n: b\mathcal{D}_n \rightarrow \text{Simp}(X_n^{\text{alg}})$ by $\mu'_n(G) = |G|^{(0)}$. The poset $b\mathcal{D}_n$ was defined near the beginning of section 2.4. Recall that $|G|^{(0)}$ is the subset of $\text{Aut}(F_n)$ corresponding to all of the minimal elements of the poset bG . It follows from section 2.5, where the ordering determined by G was discussed, that $|G|^{(0)}$ is a simplex of X_n^{alg} . Also, μ'_n preserves orders because if $G_1 < G_2$ in $b\mathcal{D}_n$, then bG_1 is a subposet of bG_2 .

Taking geometric realizations gives a map $\mu_n: X_n \rightarrow X_n^{\text{alg}}$ which is $\text{Aut}(F_n)$ -equivariant. Letting $n \rightarrow \infty$ gives an AUT-equivariant map $\mu: X \rightarrow X^{\text{alg}}$. Similarly we define maps $\tilde{\mu}_n: \tilde{X}_n \rightarrow \tilde{X}_n^{\text{alg}}$ and $\tilde{\mu}: \tilde{X} \rightarrow \tilde{X}^{\text{alg}}$ which are StN_n - and StN -equivariant.

To help get a feeling for these maps, we mention that μ_n takes the 0-cells of X_n bijectively to the vertices of X_n^{alg} , and takes the set of 1-cells of X_n injectively to the set of edges of X_n^{alg} . Also, if s is a 2-cell of X_n having the shape of the square $\sigma^1 \times \sigma^1$, then $\mu_n(s)$ lies halfway between the two possible triangulations of s (which don't introduce any new vertices) in the following sense. The interior of s is taken by μ_n to the interior of a 3-simplex s' of X_n^{alg} whose four faces correspond to the two possible triangulations of s . The four 1-cells forming the boundary of s are taken to the four edges of s' different from the two diagonals of s .

2.11. Maps between various K-theories and Pre-K-theories. The following is an application of Proposition 2.21.

Theorem 2.23. *There is a map $f: \Omega BStN^+ \rightarrow \tilde{X}^{\text{alg}}$, defined up to homotopy, which makes the diagram*

$$\begin{array}{ccc} \Omega BStN^+ & \xrightarrow{f} & \tilde{X}^{\text{alg}} \\ \downarrow g & & \downarrow h \\ \Omega BSt(\mathbb{Z})^+ & \xrightarrow{\sim} & V(St(\mathbb{Z})) \end{array}$$

homotopy commute. The map g in the diagram is induced by the homomorphism $StN \rightarrow St(\mathbb{Z})$, and h was defined in section 2.9. The lower horizontal map is the homotopy equivalence of [11].

Even though maps from algebraically to geometrically defined spaces are rare, we conjecture that there is a map $\hat{f}$ such that the diagram

$$\begin{array}{ccc} & & \tilde{X} \\ & \nearrow \hat{f} & \downarrow \tilde{\mu} \\ \Omega BStN^+ & \xrightarrow{f} & \tilde{X}^{\text{alg}} \end{array}$$

homotopy commutes.

Proof of Theorem 2.23. We will show that there is a homotopy commutative diagram

$$\begin{array}{ccccc}
 & & \Omega BStN^+ & \xrightarrow{g} & \Omega BSt(\mathbb{Z})^+ \\
 & & \downarrow & & \downarrow \\
 \tilde{X}^{\text{alg}} & \xrightarrow{\sim} & F & \xrightarrow{\sim} & F_1 \\
 \downarrow h & \searrow & \downarrow & \searrow & \downarrow \\
 & & V(St(\mathbb{Z})) & \xrightarrow{\sim} & F_1 \\
 \downarrow & & \downarrow & & \downarrow \\
 \tilde{X}^{\text{alg}}/StN & \xrightarrow{\quad} & \tilde{X}^{\text{alg}}/StN^+ & \xrightarrow{\quad} & \\
 \downarrow & \searrow & \downarrow & \searrow & \downarrow \\
 & & V(St(\mathbb{Z}))/St(\mathbb{Z}) & \xrightarrow{\quad} & V(St(\mathbb{Z}))/St(\mathbb{Z})^+ \\
 \downarrow & & \downarrow & & \downarrow \\
 BStN & \xrightarrow{\quad} & BStN^+ & \xrightarrow{\quad} & \\
 \downarrow & \searrow & \downarrow & \searrow & \downarrow \\
 & & BSt(\mathbb{Z}) & \xrightarrow{\quad} & BSt(\mathbb{Z})^+
 \end{array}$$

with the top two horizontal arrows homotopy equivalences. The theorem will then follow from the square on the upper right.

The diagram consists of maps of (homotopy) fibrations, with the fibrations drawn vertically. The covering spaces $\tilde{X}^{\text{alg}} \rightarrow \tilde{X}^{\text{alg}}/StN$ and $V(St(\mathbb{Z})) \rightarrow V(St(\mathbb{Z}))/St(\mathbb{Z})$ along with their classifying maps give the two fibrations on the left. The map between these is induced by h .

Applying the Quillen plus construction to the four spaces in the square on the lower left of the diagram gives the square on the lower right. The four maps forming the edges of this last square are induced by the universal property of the plus construction so that the lower cube in the diagram commutes.

The spaces F and F_1 are the homotopy fibers of the maps below them, and the loop spaces above F and F_1 are the homotopy fibers of the maps below them in the usual way. By Proposition 2.21, the group $StN \approx \pi_1(BStN)$ acts trivially on the homology groups of the space $\tilde{X}^{\text{alg}}$. Since the plus construction does not alter homology, it follows from the spectral sequence comparison theorem that the induced map $\tilde{X}^{\text{alg}} \rightarrow F$ of fibers is a homology equivalence. Since $\pi_2(StN^+) \approx H_2(StN) \approx 0$ and $\pi_1(\tilde{X}^{\text{alg}}/StN^+) \approx 0$, the space F is simply-connected, as is $\tilde{X}^{\text{alg}}$. Thus by Whitehead's theorem the map $\tilde{X}^{\text{alg}} \rightarrow F$ is a homotopy equivalence.

Similarly, the map $V(St(\mathbb{Z})) \rightarrow F_1$ is a homotopy equivalence. For this, one needs that the group $St(\mathbb{Z})$ acts trivially on the homology groups of $V(St(\mathbb{Z}))$. This follows from (1.3) in [11]. Also, the complex $V(St(\mathbb{Z}))$ is simply-connected since the defining relations for the group $St(\mathbb{Z})$ are triangular. And F_1 is simply-connected by an argument like that, sketched above, showing F is.

Finally, the CW-complex $V(St(\mathbb{Z}))/St(\mathbb{Z})^+$ is simply-connected and, according to [11], acyclic. So it is contractible. Therefore the map $\Omega BSt(\mathbb{Z})^+ \rightarrow F_1$ is a homotopy equivalence. $\square$

Applying the homotopy group functors to the diagram in Theorem 2.23 and using standard K-theoretic identifications gives the commutative diagram in Corollary 2.24 below—once the following definitions have been made.

For each $i \geq 1$, let

$$KN_i = \pi_i(BAUT^+),$$

where $AUT = \lim_{n \rightarrow \infty} Aut(F_n)$. (Groups $KN_i(G)$ were defined in [7], where G is any group. But we will only use the groups $KN_i(1)$ below, where 1 is the trivial group, and abbreviate these by KN_i .) This definition is analogous to Quillen's definition $K_i(\mathbb{Z}) \equiv \pi_i(BGL(\mathbb{Z})^+)$ of the K-groups of $\mathbb{Z}$, but with automorphisms of free groups in place of automorphisms of free abelian groups.

The space $BAUT^+$ may also be thought of as a second approximation to Waldhausen's space $A(*) \equiv \lim_{m, n \rightarrow \infty} Aut(\vee_n S^m)$, the topological K-theory of the one-point space $*$. Here $Aut(X)$, where X is a based space, is the topological monoid of homotopy equivalences of X . Indeed, a first approximation to $A(*)$ is

$$(B \lim_{n \rightarrow \infty} Aut(\vee_n S^0))^+ = (B \lim_{n \rightarrow \infty} \Sigma_n)^+ = B\Sigma^+$$

or a component of $\Omega^\infty S^\infty$, with the second approximation being

$$B \lim_{n \rightarrow \infty} Aut(\vee_n S^1)^+ \simeq B \lim_{n \rightarrow \infty} Aut(F_n)^+ = BAUT^+.$$

For each $i \geq 1$, we also define a group KN_i^V by

$$KN_i^V = \pi_{i-1}(X^{alg}).$$

Corollary 2.24. *For each $i \geq 3$, the diagram*

$$\begin{array}{ccc} KN_i & \xrightarrow{f_*} & KN_i^V \\ & \searrow g_* & \downarrow h_* \\ & & K_i(\mathbb{Z}) \end{array}$$

induced by the one in Theorem 2.23 commutes.

Proof. We sketch the necessary K-theoretic identifications. According to Prop. 11.2.6 of [9] there is a homotopy fibration

$$BK_2(\mathbb{Z}) \rightarrow BSt(\mathbb{Z})^+ \rightarrow BE(\mathbb{Z})^+.$$

Similarly

$$BKN_2 \rightarrow BStN^+ \rightarrow BAUT_*^+$$

is a homotopy fibration. Here AUT_* is the index two subgroup of AUT consisting of all automorphisms which abelianize to elements of $GL(\mathbb{Z})$ with positive determinant.

Since $BE(\mathbb{Z})^+$ is the universal cover of $BGL(\mathbb{Z})^+$ and $BAUT_*^+$ the universal cover of $BAUT^+$, the long exact homotopy sequences of the four fibrations just mentioned give

$$\pi_i(BStN^+) \approx \pi_i(BAUT^+) \equiv KN_i, \quad \text{and} \quad \pi_i(BSt(\mathbb{Z})^+) \approx \pi_i(BGL(\mathbb{Z})^+) \equiv K_i(\mathbb{Z})$$

provided $i \geq 3$. Also,

$$\pi_i(\tilde{X}^{alg}) = \pi_i(X^{alg}) \equiv KN_{i+1}^V$$

for $i \geq 2$ since $\tilde{X}^{alg}$ is a cover of a component of X^{alg} . □

Remark 2.25. The prism

$$\begin{array}{ccccc} \Omega(BKN_2) & \longrightarrow & \mathbb{Z}_2 & & \\ \downarrow & \searrow & \downarrow & \searrow & \\ & & \Omega(BK_2(\mathbb{Z})) & & \\ \Omega(BStN^+) & \xrightarrow{f} & \tilde{X}^{alg} & \searrow & \\ & \searrow g & & \downarrow & \\ & & \Omega(BSt(\mathbb{Z})^+) & & \end{array}$$

homotopy commutes where the vertical maps come from the inclusions of the fibers in the fibrations mentioned in the proof just above. The lower triangle is the diagram in Theorem 2.23 without the space $V(St(\mathbb{Z}))$. The induced maps of the homotopy fibers of the vertical maps give the homotopy commutative diagram

$$\begin{array}{ccc} \Omega^2(BAUT_*^+) & \longrightarrow & \Omega X_*^{alg} \\ & \searrow & \downarrow \\ & & \Omega^2(BE(\mathbb{Z})^+) \end{array}$$

Applying π_0 to this gives a commutative diagram like the one in Corollary 2.24 with $i = 2$. There is also such a commutative diagram with $i = 1$.

3. ELIMINATING CUBICAL TERMS

This section and the next two are devoted to the proof of the following.

Theorem 3.1. *The homology group $H_3(\tilde{X}_n/StN_n)$ is trivial provided $n \geq 7$.*

The proof is broken up into a number of lemmas, with each showing that all the elements of the third homology group of some subcomplex of $\tilde{X}_n/StN_n$ have representatives in a smaller subcomplex. All of these lemmas, except for 3.15, are true if $n = 6$, and many are true for lesser n . Nevertheless, for convenience, we assume throughout this section and the next that $n \geq 7$.

We first introduce some terminology. Let G be a colored or partitioned graph. An edge of G is called a singleton edge if it is the only element of the edgeset containing it. The core of G , or $core(G)$, is the subgraph of G consisting of all the nonsingleton edges of G , together with either the coloring or partition of these inherited from that of G . The graph $core(G)$ may contain bivalent vertices other than the basepoint. We often identify the vertices of G with those of $core(G)$.

Let $C_k(\tilde{X}_n/StN_n)$ denote the group of cellular k -chains in $\tilde{X}_n/StN_n$ and $Z_3(\tilde{X}_n/StN_n)$ the subgroup of 3-cycles. If G is a colored graph with $k + 1$ vertices, and a choice of orientation of the corresponding cell in $\tilde{X}_n/StN_n$ has been made, then $gen(G)$ denotes the corresponding generator of $C_k(\tilde{X}_n/StN_n)$. This choice of orientation is often referred to




FIGURE 2. All reduced, partitioned, cubical graphs.

as a cell orientation for G . Cell orientations are discussed more after the proof of Lemma 3.5 below. Conversely, if x is a generator of $C_k(\tilde{X}_n/StN_n)$ corresponding to some k -cell of $\tilde{X}_n/StN_n$, then $cg(x)$ denotes the colored graph corresponding to the k -cell. The underlying partitioned graph of a colored graph G is denoted by $par(G)$. Usually $par(cg(x))$ is shortened to just $par(x)$.

A non-basepoint vertex v of a colored graph G is called an m -valent, k -color vertex if, in $core(G)$, the set of edges meeting v has a subset consisting of m edges of exactly k different colors. Let $\mathcal{F}_{4,2}$ be the subgroup of $Z_3(\tilde{X}_n/StN_n)$ generated by elements corresponding to colored graphs with no 4-valent, 2-color vertices. A d -cell of $\tilde{X}_n/StN_n$ corresponding to the colored graph G with edge sets $h_1, \dots, h_k$ is referred to as an $n_1, \dots, n_k$ -cell if there is some permutation σ of the set $\mathbf{k}$ with $|h_{\sigma(i)}| - 1 = n_i$ for each i . (In this case, $\sum_{i=1}^k n_i = d$.)

A cubical (resp. prismatic, simplicial) generator of $C_3(\tilde{X}_n/StN_n)$ is a generator which corresponds to a 1, 1, 1-cell (resp. 1, 2-cell, 0, 3-cell) of $\tilde{X}_n/StN_n$.

We say that a colored or partitioned graph G has core of type Cn if the partitioned graph $core(G)$ is isomorphic to the core of the partitioned graph labeled Cn (or $Cn(a)$ if $n = 1$ or 2) in figure 2. A generator x of $C_3(\tilde{X}_n/StN_n)$ and the corresponding colored graph $cg(x)$ are both said to be core-reduced if $cg(x)$ has core of one of the types $C1$ through $C7$. Let $\mathcal{F}_{CR}$ be the subgroup of $Z_3(\tilde{X}_n/StN_n)$ consisting of all the cycles with core-reduced cubical terms.

A cubical generator x of $C_3(\tilde{X}_n/StN_n)$ and $cg(x)$ are both said to be reduced if the partitioned graph $par(x)$ is isomorphic to one of the nine partitioned graphs shown in figure 2. Assuming x is reduced, we say that both x and $cg(x)$ are of type Cn if $par(x)$ is isomorphic to the partitioned graph labeled Cn in Figure 2. A 3-chain is said to be reduced if all its cubical terms are. Let $\mathcal{F}_R$ be the subgroup of $Z_3(\tilde{X}_n/StN_n)$ consisting of all the reduced cycles.

Let $\mathcal{F}$ be the filtration

$$Z_3(\tilde{X}_n/StN_n) \supset \mathcal{F}_{4,2} \supset \mathcal{F}_{CR} \supset \mathcal{F}_R \supset \mathcal{F}_{C7} \supset \mathcal{F}_{\theta P} \supset \mathcal{F}_{C5} \supset \mathcal{F}_{C4} \supset \dots \supset \mathcal{F}_{C1}$$

of $Z_3(\tilde{X}_n/StN_n)$ where $z \in \mathcal{F}_{Cn}$ if $z \in \mathcal{F}_R$ and none of the cubical terms of z are of type Cm for $m \geq n$. The subgroup $\mathcal{F}_{\theta P}$ is defined just before lemma 3.7 below. The lemmas in this section show that any element of a subgroup of the filtration $\mathcal{F}$ is homologous in the space $\tilde{X}_n/StN_n$ to an element of the next smaller subgroup of $\mathcal{F}$. Thus, any element of $H_3(\tilde{X}_n/StN_n)$ has a representative in $\mathcal{F}_{C1}$.

Lemma 3.2. *Each $z \in Z_3(\tilde{X}_n/StN_n)$ is homologous to a cycle in $\mathcal{F}_{4,2}$.*




FIGURE 3. All (unbased) cores with a 4-valent, 2-color vertex.

Proof. Let x be any cubical term of a cycle $z \in Z_3(\tilde{X}_n/StN_n)$. Let G be the colored graph $cg(x)$ where the edgesets are $\{a_1, a_2\}$, $\{b_1, b_2\}$ and $\{c_1, c_2\}$. Let v be a 4-valent, 2-color vertex of G meeting, say, the edges a_1, a_2, b_1 and b_2 . Since G has four vertices, at least two of these edges form a monochromatic loop, say a_1 and a_2 do. Then if b_1 and b_2 do not form a loop, $core(G)$ is as shown in figure 3(a). And if b_1 and b_2 do form a loop, then both c_1 and c_2 meet the vertex of G which is disjoint from both of the loops a_1, a_2 and b_1, b_2 . The four possibilities are shown in b(i) through b(iv) of figure 3.

In case the core of G is as in figure 3(a) and u is the basepoint of G , let G' be a colored graph with five vertices with core as shown in figure 4(a) and satisfying $G'_b = G$. Orient $cell(G')$ so that the boundary $d(gen(G'))$ contains the term $-x$. We say that $gen(G')$ is obtained from x by separating the edges a_1 and a_2 of $G = cg(x)$ along the edges b_1 and b_2 of G . Similar terminology is often used below. Note that the two cubical terms G'_{b_1} and G'_{b_2} of the chain $x + d(gen(G'))$ contain no 4-valent, 2-color vertices. Such vertices can be eliminated in similar ways in the other cases by adding suitable boundaries. For instance, in case the core of G is as in figure 3(a) with w as the basepoint, add to x the boundary of the generator of $C_4(\tilde{X}_n/StN_n)$ obtained from x by separating the edges b_1 and b_2 of G along a_1 and a_2 . If $core(G) = b(i)u$, meaning that $core(G)$ is as in b(i) of figure 3 with u as the basepoint, or if $core(G) = b(i)w$ with a similar meaning, then separate, along the monochromatic loop through v and w , the two edges of the other monochromatic loop of G . If $core(G) = b(ii)u$, then separate, along the monochromatic loop through u and v , the two edges of the other monochromatic loop through v . If $core(G) = b(ii)w$, first separate the two edges of the monochromatic loop through u and v along c_1 and c_2 , and then use the first case above twice. If $core(G) = b(iii)w$, then first separate the edges a_1 and a_2 from b_1 and b_2 along c_1 and c_2 , and then use a previous case twice. If $core(G) = b(iv)u$ or w , then separate, along the monochromatic loop through $*$, the two edges of the other monochromatic loop of $core(G)$. Finally, if $core(G) = b(iv)t$, then use a graph with core as shown in figure 4(b). This covers all the cases, and so completes the proof. $\square$

Lemma 3.3. *Each $z \in \mathcal{F}_{4,2}$ is homologous to an element of $\mathcal{F}_{CR}$.*

Proof. Let x be a cubical term of z such that $core(cg(x))$ has a 4-valent, 3-color vertex v . Since v meets all nonsingleton colors and is not a 4-valent, 2-color vertex, v lies on exactly one monochromatic loop. We claim that $core(cg(x))$ is as shown in either (i) or (ii) of figure 4. Indeed, given that a_1, b_1, c_1 and c_2 meet v , of the six ways a_2 can be attached to the four vertices of $cg(x)$, all but the two ways shown are ruled out by the fact that $cg(x)$ is colored and v is not a 4-valent, 2-color vertex. The same facts then imply that b_2 must be attached




FIGURE 4.




FIGURE 5.

as shown, or that, in figure 4(ii), b_2 joins the vertices t and u . This last possibility gives the same core as in (i).

Next we indicate, in all but one case, how the 4-valent, 3-color vertices of $cg(x)$ can be eliminated by adding suitable boundaries to x , without introducing any 4-valent, 2-color vertices. Let $G = cg(x)$. If $core(G) = 4(i)w$, meaning that the core of G is the partitioned graph shown in (i) of figure 4 with w as the basepoint, or if $core(G) = 4(ii)w$ then, in both cases, separate a_1 and b_1 along c_1 and c_2 . If $core(G) = 4(i)u$, use a graph with core as shown in figure 5(a), and if $core(G) = 4(ii)t$, first use a graph with core as shown in figure 5(b), and then use a previous case.

We may now assume that G has no 4-valent, 2-color vertices and if G has a 4-valent, 3-color vertex, then $core(G)$ is as shown in figure 4(i) with t as the basepoint, that is, $core(G)$ is of type C7. To complete the proof, we show that $core(G)$ has type C_n for some n between 1 and 7.

First assume that G has more than one monochromatic loop. Then since G has no 4-valent, 2-color vertices, these loops are either disjoint or meet at the basepoint. If they're disjoint, then two edges of the same color do not form a loop and these two edges are either disjoint or meet at $*$. In these cases $core(G)$ is either of type C6 or C4. If the monochromatic loops meet at $*$, then there are either two or three of them, and so $core(G)$ is of type C1, C2 or C3.

Now assume that G has just one monochromatic loop consisting, say, of the edges a_1 and a_2 , and let b_1 and b_2 be two other edges of G of the same color. If b_1 and b_2 form a disconnected subgraph of G , then $core(G)$ is of type C6. So assume $b_1 \cap b_2 \neq \emptyset$. If $(b_1 \cap b_2) \cap (a_1 \cup a_2) \neq \emptyset$, then this intersection consists of the basepoint of G , and so $core(G)$

is of type C4. If $(b_1 \cap b_2) \cap (a_1 \cup a_2) = \emptyset$, then $(b_1 \cup b_2) \cap (a_1 \cup a_2)$ consists of either the one or two vertices of the loop $a_1 \cup a_2$. In both cases it follows from the assumptions in effect and the fact that each non-basepoint vertex of G is 2-valent, 1-color that $\text{core}(G)$ is of type C5 or C7. $\square$

Definition 3.4. If a colored graph has a non-basepoint vertex v such that the set of edges meeting v has only one subset consisting of two edges of the same color, then, by separating the edges meeting v , we mean separate, along these two edges of the same color, the other half-edges incident with v .

By the w -vertex of a colored graph G with type C2 core, we mean the vertex which corresponds to the vertex labeled w of the graph in C2(a) of Figure 2. Similar conventions, using the labeling of the vertices in Figure 2, are used to designate the other non-basepoint vertices of G , as well as the vertices of colored graphs with other core types.

Lemma 3.5. *Each $z \in \mathcal{F}_{CR}$ is homologous to a cycle in $\mathcal{F}_R$.*

Proof. Let x be any cubical term of $z \in \mathcal{F}_{CR}$, and let $G = cg(x)$. Each of the possible seven core types for G will be considered separately.

Case C1. Assume first that G has type C1 core. If a non-basepoint vertex, say v , of G has valence > 3 , then separate the edges of G meeting v to obtain the colored graph G' . Choose the orientation of the corresponding cell so that x cancels with one of the terms of $d(\text{gen}(G'))$. Then the colored graphs corresponding to the two cubical terms of $x + d(\text{gen}(G'))$ both have type C1 cores. Also, the sum of the valences of the non-basepoint vertices of each of these is less than that of G . So by separating edges repeatedly as needed, we may assume that each cubical term of z with type C1 core is reduced.

Case C2. Assume now that G has type C2 core. By separating the edges incident with the w -vertex of G , and then possibly of resulting colored graphs with type C2 cores, we may assume that the w -vertex has valence 3. Then by separating the edges meeting the u -vertex of G , we may assume that both the u - and w -vertices of G have valence 3. Finally, let $\text{gen}(G')$ be obtained from x by separating the singleton edges meeting the v -vertex of G from the nonsingleton edge which joins the u - and v -vertices. Then one cubical term of $x + d(\text{gen}(G'))$ corresponds to a colored graph with type C1 core. The other corresponds to one with type C2 core with v -vertex having valence 3. This and case C1 give a chain c such that $x + dc$ is reduced. In fact, each cubical term has type C1 or C2.

Case C3. Next assume that G has type C3 core. By repeatedly separating the edges incident with the v -vertex of G , we may assume that it has valence 3. Then separating edges incident with the u - and w -vertices of G and possibly resulting colored graphs with type C3 cores and the previous cases give a 4-chain c such that $x + dc$ is reduced.

Case C4. If G has type C4 core, then, as usual, we may assume that the u -vertex of G has valence 3. Let G' be the partitioned graph shown in (c) of figure 5, colored and with dangling half-edges joined so that $G'_a = G$. Orient the cell of $\tilde{X}_n/StN_n$ corresponding to G' so that $d(\text{gen}(G'))$ contains the term $-x$. We say that $\text{gen}(G')$ is obtained from x by moving the singleton edges incident with the v -vertex of G toward the basepoint of G . Similar terminology is often used below. The two colored graphs corresponding to the cubical terms of $x + d(\text{gen}(G'))$ have cores of type C2 and C4. The valence of the v -vertex of the one with type C4 core is 3. So moving the singleton edges incident with its w -vertex toward the basepoint and case C2 give a chain c such that $x + dc$ is reduced.

Case C5. If G has type C5 core then, as usual, we may assume that the u -vertex of G has valence 3. Moving the singleton half-edges incident with the v -vertex of G toward the w -vertex gives a chain c' such that the two colored graphs corresponding to the cubical terms of $x + dc'$ have cores of type C2 and C5. The valence of both the u - and v -vertices of the one with type C5 core is 3. So separating edges meeting its w -vertex and using case C2 gives a chain c such that $x + dc$ is reduced.

Case C6. Assume that G has type C6 core. By moving the singleton edges of G incident with the u -vertex toward the basepoint, we may assume that the u -vertex has valence 3. For this, a 4-valent, 2-color vertex has to be eliminated as in the proof of 3.2, and case C5 is used. Then by moving the singleton edges meeting the v -vertex of G toward the w -vertex and using case C3, we may assume that the v -vertex of G has valence 3. Finally, moving the singleton edges incident with the w -vertex toward the basepoint along either edge and then using case C4 gives a chain c such that $x + dc$ is reduced. The terms of $x + dc$ can be of all possible types except for C7.

Case C7. If G has type C7 core, then first let $gen(G')$ be obtained from x by moving the singleton edges incident with the w -vertex of G toward the v -vertex of G . Then one cubical term of $x + d(gen(G'))$ corresponds to a colored graph with type C7 core with w -vertex of valence 3. The other has a 4-valent, 2-color vertex. This can be eliminated as in the proof of 3.2 at the expense of introducing a colored graph with a type C5 core, and another with a 4-valent, 3-color vertex. Eliminating this vertex, in turn, as done above, gives colored graphs with cores of type C3 and C5. So by previous cases we may assume that the w -vertex of G has valence 3.

We may also assume that the v -vertex of G also has valence 3 by moving the singleton edges incident with the v -vertex toward the basepoint of G , and then using case C5. Finally, by moving the singleton edges incident with the u -vertex of G toward the basepoint, eliminating, as in the proof of 3.3, one of the 4-valent, 3-color vertices which arise, and then using cases C4 and C5, we obtain a chain c such that $x + dc$ is reduced. The terms of $x + dc$ could be of any type other than C6.

This covers all the cases and so shows that each term of z is homologous to some element of $\mathcal{F}_R$. So z also is. $\square$

We next discuss cell orientations of colored graphs and the effect the cellular boundary map has on these. Let G be a StN-marked graph with edgesets $h_1, \dots, h_n$, and let $cm(G)$ be the corresponding cell in $\tilde{X}$. Also let $c(G)$ be the cell in $\tilde{X}/StN$ corresponding to the unmarked colored graph G . By the discussion following Lemma 2.19, the cell $cm(G)$ is homeomorphic to the product $\sigma^{\|h_1\|} \times \dots \times \sigma^{\|h_n\|}$ where $\sigma^{\|h_i\|}$ is the standard simplex of dimension $\|h_i\| = |h_i| - 1$. Also, the vertices of each $\sigma^{\|h_i\|}$ correspond, in a natural way, to the elements of the set h_i . Thus a choice, up to an even permutation, of an ordering of the edges in $h_i = \{e_i^j\}_{j=0}^{k_i}$ serves to orient $\sigma^{\|h_i\|}$. For each i , let $e_i^{\pi_i(0)}, \dots, e_i^{\pi_i(k_i)}$ be such a choice (where π_i is a bijection of the set $\{1, \dots, k_i\}$) and let $[e_i^{\pi_i(0)}, \dots, e_i^{\pi_i(k_i)}]$ denote $\sigma^{\|h_i\|}$ with the resulting orientation. The cells $cm(G) \approx \sigma^{\|h_1\|} \times \dots \times \sigma^{\|h_n\|}$ and $c(G)$ are then oriented by the product orientation. Let $[e_1^{\pi_1(0)}, \dots, e_1^{\pi_1(k_1)}] \times \dots \times [e_n^{\pi_n(0)}, \dots, e_n^{\pi_n(k_n)}]$ denote both the cell $c(G)$, so oriented, and also the generator of the cellular chain group of $\tilde{X}/StN$ it corresponds to. With these conventions the cellular boundary map $d: C_*(\tilde{X}_n/StN_n) \rightarrow C_{*-1}(\tilde{X}_n/StN_n)$ is given by




FIGURE 6.

$$\begin{aligned}
 d\left([e_1^{\pi_1(0)}, \dots, e_1^{\pi_1(k_1)}] \times \dots \times [e_n^{\pi_n(0)}, \dots, e_n^{\pi_n(k_n)}]\right) = \\
 \sum_{\substack{1 \leq i \leq n, k_i \neq 0 \\ 0 \leq j \leq k_i}} (-1)^{k_1 + \dots + k_{i-1}} (-1)^j [e_1^{\pi_1(0)}, \dots, e_1^{\pi_1(k_1)}] \times \dots \times [e_i^{\pi_i(0)}, \dots, \widehat{e_i^{\pi_i(j)}}, \dots, e_i^{\pi_i(k_i)}] \times \dots \\
 \dots \times [e_n^{\pi_n(0)}, \dots, e_n^{\pi_n(k_n)}]
 \end{aligned}$$

where the term containing $\widehat{e_i^{\pi_i(j)}}$ denotes the generator of $C_*(\tilde{X}/StN)$ corresponding to the colored graph $G_{e_i^{\pi_i(j)}}$ with orientation that inherited from G . Also the exponent $k_i + \dots + k_{i-1}$ is 0 in case $i = 1$. Just the orientation of the oriented cell $[e_1^{\pi_1(0)}, \dots, e_1^{\pi_1(k_1)}] \times \dots \times [e_n^{\pi_n(0)}, \dots, e_n^{\pi_n(k_n)}]$ is denoted by $[e_1^{\pi_1(0)}, \dots, e_1^{\pi_1(k_1)}] \dots [e_n^{\pi_n(0)}, \dots, e_n^{\pi_n(k_n)}]$. Singleton edges are often dropped from the notation. So, for instance, the orientation $[e_1^0], [e_2^0, e_2^1], [e_3^0]$ may be denoted by just $[e_2^0, e_2^1]$.

In preparation for the proof of the next lemma, we introduce notation for a number of generators of $C_*(\tilde{X}_n/StN_n)$, and also define a homomorphism involving these which counts relations of the form $w_{ij}^L = w_{ij}^R$. Let θ_{xy} and $(Lx/Ry/z)$ be the generators of $C_2(\tilde{X}_n/StN_n)$ and $C_3(\tilde{X}_n/StN_n)$ indicated in figure 6. The three other generators of $C_3(\tilde{X}_n/StN_n)$, analogous to $(Lx/Ry/z)$, but with different orientations of the edgesets x and y , are denoted by changing the prefixal L and R accordingly. We say that a colored graph G with three vertices is of type θ if the core of G is equivalent to the colored graph θ_{xy} in Fig. 6 for some choice of colors x and y .

Let $A_+(x/y/z)$ be the generator of $C_4(\tilde{X}_n/StN_n)$ indicated in figure 7. Another generator $A_-(x/y/z)$ is defined similarly, but with the orientation of the edgeset y reversed. Let $A'_+(x/y/z) = A_+(x/y/z) + c$ where c is a chain such that

$$(2) \quad d(A'_+(x/y/z)) \equiv (Lx/Ly/z) - (Rx/Ry/z) \pmod{\mathcal{F}_{C7}}.$$

The natural choice for c is $gen(G)$ where G is a colored graph as on the left of figure 7, but with the partition $\{a, b, d\}\{c, e\}\{f, g\}$ and with suitable cell orientation. With this choice, the element of $\mathcal{F}_{C7}$ by which the two equivalent elements in (2) differ contains two cubical terms, both with type C6 cores. Define $A'_-(x/y/z)$ similarly so that

$$(3) \quad d(A'_-(x/y/z)) \equiv (Lx/Ry/z) - (Rx/Ly/z) \pmod{\mathcal{F}_{C7}}.$$

Let $B(Rw/Rx/Ry/z)$ be the generator indicated in figure 7. The seven other generators obtained by changing the orientations of the edgesets w , x and z are denoted in the same




FIGURE 7.

way, but with suitable prefixes for w , x and z . Applying lemmas 3.2, 3.3 and 3.5 and their proofs to the six noncanceling faces of $B(Pw/Qx/Sy/z)$, where $P, Q, S \in \{L, R\}$, gives a chain c such that, with $B'(Pw/Qx/Sy/z) = B(Pw/Qx/Sy/z) + c$, we have

$$(4) \quad d(B'(Pw/Qx/Sy/z)) \equiv -(Qx/Sy/z) + (Pw/Sy/z) - (Pw/Qx/z) \pmod{\mathcal{F}_{C7}}.$$

Let $C(Rx/y/z)$ be the generator indicated in figure 7, and let $C(Lx/y/z)$ be the same but with the orientation of the edgeset x reversed.

For each $S \in \{L, R\}$, let c_S be a generator of $C_4(\tilde{X}_n/StN_n)$ such that with $C'(Sx/y/z) = C(Sx/y/z) + c_S$ we have

$$(5) \quad d(C'(Sx/y/z)) \equiv (Sx/Ry/z) - (Sx/Ly/z) \pmod{\mathcal{F}_{C7}}.$$

In both cases, $par(c_S)$ can be taken to be as in C6 of figure 2, but with the basepoint a bivalent vertex on one of the monochromatic loops. Define a homomorphism $f: C_2(\tilde{X}_n/StN_n) \rightarrow \oplus_{1 \leq i, j \leq n} \mathbb{Z}_{i, j}$ by mapping generators according to

$$f(x) = \begin{cases} 0, & \text{if } core(cg(x)) \text{ is not of type } \theta, \\ 1_{ij}, & \text{if } core(cg(x)) = \theta_{ij}. \end{cases}$$

Here each $\mathbb{Z}_{i, j}$ is a copy of $\mathbb{Z}$ and $1_{i, j} \in \mathbb{Z}_{i, j}$ is the generator corresponding to $+1 \in \mathbb{Z}$. Also, in the second line of the definition, we are viewing $core(cg(x))$ as a generator of $C_2(\tilde{X}_n/StN_n)$ with cell orientation that inherited from x . Let Θ be the composite $f \circ d$ where $d: C_3(\tilde{X}_n/StN_n) \rightarrow C_2(\tilde{X}_n/StN_n)$ is the cellular boundary map. Then Θ is trivial on simplicial generators since the cells corresponding to these have no square faces. Also, one can check that Θ is trivial on prismatic generators. The only reduced cubical terms on which Θ is nontrivial are those of type C7, and for these $\Theta(Sx/Ty/z) = 1_{xz} - 1_{yz}$ for all four choices of $S, T \in \{L, R\}$.

Lemma 3.6. *Each $z \in \mathcal{F}_R$ is homologous to a cycle in $\mathcal{F}_{C7}$.*

Proof. Let z_1 be any type C7 cubical term of $z \in \mathcal{F}_R$. Since $\Theta(z) = 0$, there is a cubical term z_2 of z different from z_1 such that some cancelation occurs among the terms of $\Theta(z_1)$ and $\Theta(z_2)$. The following three cases cover the various forms z_2 can have. In all cases, we show that, by adding boundaries to z , it is possible to reduce the total number of type C7 terms of z , while keeping $z \in \mathcal{F}_R$.

Case 1. Assume that $z_1 + z_2 = (Sx/Ty/z) + (Pw/Qx/z)$. By (5) or (2) and (3), we may assume that $S = Q$. If $y \neq w$, then, using (4), we see that $z + d(B'(Pw/Sx/Ty/z))$ is in

$\mathcal{F}_R$ and has one fewer type C7 term than z . If $y = w$, we may assume, again by (5), that $T = P$. Then $z + d(B'(Lw/Sx/Ty/z) + B'(Lw/Ty/Sx/z))$ is in $\mathcal{F}_R$ and has two fewer type C7 terms than z .

Case 2. Assume $z_1 + z_2 = (Sx/Ty/z) - (Px/Qw/z)$. By (2) and (3), we may assume that $S = P$. If $y \neq w$, then $z + d(B'(Sx/Ty/Qw/z))$ is in $\mathcal{F}_R$ and has one fewer type C7 term than z . If $y = w$, and $z_1 + z_2 \neq 0$, then in view of (5), both z_1 and z_2 can be eliminated from z .

Case 3. The last case, in which $z_1 + z_2 = (Sx/Ty/z) - (Pw/Qy/z)$ is similar to the previous ones. First suppose that $x \neq w$. We may assume that $T = Q$ by (5), and then add $-d(B'(Sx/Pw/Ty/z))$ to z . If $x = w$, we may first assume that $S = P$ by (2) and (3), and then finish with the help of (5). $\square$

Both a generator x of $C_3(\tilde{X}_n/StN_n)$ and the corresponding colored graph $cg(x)$ are referred to as θ -prismatic if $core(cg(x))$ is as shown in (a) of figure 6. The u -vertex of a θ -prismatic colored graph is the one corresponding to the vertex labeled u in the figure. The subgroup $\mathcal{F}_{\theta P}$ of the filtration $\mathcal{F}$ is defined by $z \in \mathcal{F}_{\theta P}$ if $z \in \mathcal{F}_{C7}$ and if z contains no θ -prismatic terms.

Lemma 3.7. *Each $z \in \mathcal{F}_{C7}$ is homologous to a cycle in $\mathcal{F}_{\theta P}$.*

Proof. Any θ -prismatic term of z differs by boundaries from a sum of simplicial terms and prismatic terms where each θ -prismatic term in this sum corresponds to a graph with all nonbasepoint vertices of valence 3. To see this, let G be the colored graph corresponding to any θ -prismatic generator. By separating the singleton edges incident with the u -vertex of G , and then doing this with any resulting θ -prismatic colored graphs having u -vertices with valence > 3 , we may assume that the u -vertex of G is trivalent. Then by moving the singleton edges of G incident with the other two non-basepoint vertices toward the basepoint, we may assume that each θ -prismatic term of z corresponds to a graph with all non-basepoint vertices trivalent. Next we show how to eliminate such terms.

Let $c \in C_4(\tilde{X}_n/StN_n)$ be the sum of the seven generators, $g_1, \dots, g_7$ say, corresponding to the seven colored graphs and orientations shown in figure 8. Any consistent choice of orientations of the edgesets may be used. Pairs of arrows pointing at each other indicate cancellation among the terms dg_i of dc . For instance, if G is the colored graph in the upper right of figure 8, and H is the colored graph to its left, then $G_c = H_{a_2}$, and the cell orientations are such that the two corresponding elements of $C_3(\tilde{X}_n/StN_n)$ cancel. With the help of figure 8, it's routine to check that dc is reduced with each cubical term of type C4 or C5, and that dc has exactly one θ -prismatic term coming from the colored graph G_{c_2} . Since each non-basepoint vertex of G_{c_2} is trivalent, all the θ -prismatic terms of z can be eliminated. $\square$

Given $S_1, \dots, S_k \in \{L, R\}$ and distinct positive integers $a_1, \dots, a_{k+1}$, let $F_k(S_1a_1, \dots, S_ka_k, a_{k+1})$, or just $F_k(S_ia_i, a_{k+1})$, be the colored graph with $k+1$ vertices shown on the left of figure 9. Here the edgeset h_{a_i} points away from the basepoint if $S_i = R$ and towards it if $S_i = L$. We also let $F_k(S_ia_i, a_{k+1})$ denote the generator of $C_k(\tilde{X}_n/StN_n)$ corresponding to the colored graph $F_k(S_ia_i, a_{k+1})$ with cell orientation $[a_2^-, a_2^+] \dots [a_{k+1}^-, a_{k+1}^+]$. Here we are assuming that a_i^- meets the basepoint for $2 \leq i \leq k$, and that the orientation of the loop which starts at the basepoint, goes along a_{k+1}^- , and then along a_{k+1}^+ back to the basepoint, agrees with the





FIGURE 8.

FIGURE 9. Give each of g_1^R , g_2 and g_3 the cell orientation $[a_1, a_2][b_1, b_2]$.

orientation of the edgeset $\{a_{k+1}^-, a_{k+1}^+\}$. A hat over a symbol in the following indicates, as usual, omission of that symbol.

Lemma 3.8. *There is a chain $c \in C_4(\tilde{X}_n/StN_n)$ such that*

$$d(F_4(S_i a_i, a_5) + c) \equiv -F_3(S_1 a_1, \widehat{S_2 a_2}, \dots, a_5) + F_3(S_1 a_1, S_2 a_2, \widehat{S_3 a_3}, S_4 a_4, a_5) - F_3(S_1 a_1, \dots, \widehat{S_4 a_4}, a_5) \pmod{\mathcal{F}_{C_3}}.$$

Proof. We denote $F_3(S_1 a_1, \dots, \widehat{S_i a_i}, \dots, a_5)$ by just $F_3(\widehat{S_i a_i})$. Assume that the nonsingleton edgesets of $F_4(S_i a_i, a_5)$ are $\{a_i^-, a_i^+\}$ as in figure 9. Let $d_{a_i^+}(F_4(S_i a_i, a_5))$ be the term of the boundary $d(F_4(S_i a_i, a_5))$ corresponding to the colored graph $F_4(S_i a_i, a_5)$ with the edge a_i^+ blown down. Note that, for $i = 2, 3$ and 4 , the colored graph $cg(d_{a_i^+}(F_4(S_i a_i, a_5)))$ contains a colored subgraph Γ_i which, with its inherited cell orientation, satisfies $gen(\Gamma_i) = (-1)^{i+1} F_3(\widehat{S_i a_i})$.

Next we define a chain $c = \sum_{i=1}^6 c_i \in C_4(\tilde{X}_n/StN_n)$ whose boundary contains terms corresponding to the subgraphs Γ_i and also terms which cancel the $d_{a_i^+}(F_4(S_i a_i, a_5))$ for

$i = 2, 3$ and 4. Let c_1 be obtained from $d_{a_4^+}(F_4(S_i a_i, a_5))$ by separating the edges meeting the 4-valent vertex of $cg(d_{a_4^+}(F_4(S_i a_i, a_5)))$. One cubical term of $d_{a_4^+}(F_4(S_i a_i, a_5)) + dc_1$ is $-F_3(\widehat{S_4 a_4})$. The other is of type C2(a). Define c_2 similarly by separating the edges meeting the 4-valent vertex of $d_{a_3^+}(F_4(S_i a_i, a_5))$. Then define c_3 by separating the edges meeting the 4-valent vertex of the graph corresponding to the cubical term of dc_2 different from $d_{a_3^+}(F_4(S_i a_i, a_5))$ and $F_3(\widehat{S_3 a_3})$. Define c_4 , c_5 and c_6 similarly (by separating edges meeting 4-valent vertices) so as to isolate the subgraph Γ_2 of $d_{a_2^+}(F_4(S_i a_i, a_5))$ corresponding to $-F_3(\widehat{S_2 a_2})$. Its easy to check that with $c = \sum_{i=1}^6 c_i$, the lemma holds. No θ -prismatic terms arise since each edgeset of each colored graph considered contains an edge which meets the basepoint of G . $\square$

Lemma 3.9. *For any choice of $S_1, S_2, S_3 \in \{L, R\}$ and distinct positive integers $a_1, \dots, a_4$, there is a chain $a \in C_4(\tilde{X}_n/StN_n)$ such that*

$$da \equiv F_3(S_1 a_1, S_2 a_2, S_3 a_3, a_4) + F_3(S_1 a_1, S_3 a_3, S_2 a_2, a_4) \pmod{\mathcal{F}_{C_3}}.$$

Proof. Choose a positive integer a_5 distinct from each of $a_1, \dots, a_4$. Use lemma 3.8 to pick c_1 and $c_2 \in C_4(\tilde{X}_n/StN_n)$ such that, mod $\mathcal{F}_{C_3}$,

$$\begin{aligned} d(F_4(S_1 a_1, S_2 a_2, S_3 a_3, S_5 a_5, a_4) + c_1) \equiv \\ -F_3(S_1 a_1, S_3 a_3, S_5 a_5, a_4) + F_3(S_1 a_1, S_2 a_2, S_5 a_5, a_4) - F_3(S_1 a_1, S_2 a_2, S_3 a_3, a_4), \end{aligned}$$

and

$$\begin{aligned} d(F_4(S_1 a_1, S_3 a_3, S_2 a_2, S_5 a_5, a_4) + c_2) \equiv \\ -F_3(S_1 a_1, S_2 a_2, S_5 a_5, a_4) + F_3(S_1 a_1, S_3 a_3, S_5 a_5, a_4) - F_3(S_1 a_1, S_3 a_3, S_2 a_2, a_4). \end{aligned}$$

Adding these two congruences completes the proof. $\square$

Let $E(Ra_1, a_2, a_3)$ be the generator of $C_3(\tilde{X}_n/StN_n)$ indicated in figure 9. In case the edgeset $\{a_1^-, a_1^+\}$ has orientation opposite that shown, then the corresponding generator is denoted by $E(La_1, a_2, a_3)$.

Lemma 3.10. *For any choice of $S_1, S_2 \in \{L, R\}$ and distinct positive integers $a_1, \dots, a_4$, there is a chain $a \in C_4(\tilde{X}_n/StN_n)$ such that, mod $\mathcal{F}_{C_5}$,*

$$da \equiv F_3(S_1 a_1, S_2 a_2, La_3, a_4) - F_3(S_1 a_1, S_2 a_2, Ra_3, a_4) - E(S_1 a_1, a_3, a_4).$$

Proof. Let $a \in C_4(\tilde{X}_n/StN_n)$ be the sum of the seven generators indicated in figure 10. As with figure 8, pairs of arrows pointing at each other indicate canceling terms in da , provided a consistent choice of orientations of edgesets is used. For one such choice, the three specific generators of $C_3(\tilde{X}_n/StN_n)$ on the left of the figure are terms of da . Those terms of da not accounted for in the figure are all prismatic and none are θ -prismatic. $\square$

Let g_1^R , g_2 and g_3 be the generators of $C_2(\tilde{X}_n/StN_n)$ indicated in figure 9. Here $A = \{\alpha_i\}$ and $B = \{\beta_i\}$ are the sets consisting of all the singleton half-edges incident with the vertices indicated in the figure. Define g_1^L just as g_1^R is, except with the orientation of the edgeset y changed. The notation for the g_1^S and the g_i should include some mention of the edgesets y and z and of the sets A and B , however we suppress this. For lemma 3.11 below, we




FIGURE 10.

will count nontrivial commutator relations using the homomorphism $f: C_2(\tilde{X}_n/StN_n) \rightarrow \bigoplus_{\substack{1 \leq x,y,z \leq n \\ S,T \in \{L,R\}}} \mathbb{Z}_{Sx,Ty,z}$ defined on generators by

$$f(g) = \begin{cases} \sum_i 1(\alpha_i, Sy, z), & \text{if } g = g_1^S, \\ \sum_i (1(\alpha_i, Ry, z) - 1(\alpha_i, Ly, z)), & \text{if } g = g_2, \\ -\sum_i 1(\alpha_i, Ry, z) - \sum_j 1(\beta_j, Ly, z), & \text{if } g = g_3, \\ 0, & \text{if } g \neq \pm g_1^S, g_2, g_3. \end{cases}$$

Here each $\mathbb{Z}_{Sx,Ty,z}$ is a copy of $\mathbb{Z}$ with $1(Sx/Ty/z) \in \mathbb{Z}$ corresponding to $+1 \in \mathbb{Z}$.

Let F be the composite $f \circ d$. Then F is trivial on all simplicial generators, and on all prismatic generators except for θ -prismatic ones. Also, F is trivial on all reduced cubical generators except for those of type C5, C6 and C7. So the only terms of any $z \in \mathcal{F}_{\theta P}$ on which the homomorphism F is nontrivial are type C5 and C6 cubical ones. Moreover, for these we have

$$(6) \quad F(F_3(S_1a_1, S_2a_2, S_3a_3, a_4)) = -1(S_1a_1, S_3a_3, a_4) + 1(S_1a_1, S_2a_2, a_4),$$

and

$$(7) \quad F(E(Sa_1, a_2, a_3)) = 1(Sa_1, Ra_2, a_3) - 1(Sa_1, La_2, a_3).$$

For $S \in \{L, R\}$, define $\bar{S}$ by $\{S, \bar{S}\} = \{L, R\}$.

Lemma 3.11. *Each $z \in \mathcal{F}_{\theta P}$ is homologous to a cycle in $\mathcal{F}_{C5}$.*

Proof. Write $z = \sum_{i=1}^n a_i$ where n is minimal and each a_i is a generator of $C_3(\tilde{X}_n/StN_n)$ corresponding to some 3-cell of $\tilde{X}_n/StN_n$. Let $N_5(z)$ be the number of a_i of type C5, and let $N_6(a_i)$ be the number of type C6. Also let $N_{5+6}(z) = N_5(z) + N_6(z)$. We will show that if $N_{5+6}(z) > 0$, then z is homologous to a cycle $z' \in \mathcal{F}_{\theta P}$ with $N_{5+6}(z) < N_{5+6}(z')$.




FIGURE 11.

Since $F(z) = 0$, it follows from (7) that if $N_6(z) > 0$, then $N_5(z) > 0$ as well. So we assume that some a_i , say a_1 , is of type C5 with

$$F(a_1) = 1(Sx, Ty, w) - 1(Sx, Pz, w).$$

Again since $F(z) = 0$, there is another a_i , say a_2 , such that some cancelation occurs between the terms of $F(a_1)$ and $F(a_2)$. Two cases arise.

Case 1. Assume that a_2 is of type C5 with $F(a_2) = 1(Sx, Qu, w) - 1(Sx, Av, w)$. Then either $Qu = Pz$ or $Av = Ty$, say $Av = Ty$. We may assume, by lemma 3.9, that $a_1 = F_3(Sx, Ty, Pz, w)$ and $a_2 = -F_3(Sx, Ty, Qu, w)$. If $u \neq z$, then, as in lemma 3.8, $z' = z + d(F_4(Sx, Ty, Pz, Qu, w) + c)$ is in $\mathcal{F}_{\theta P}$ and satisfies $N_{5+6}(z') = N_{5+6}(z) - 1$.

If $u = z$ and $P \neq Q$, then lemma 3.10 gives a boundary b with $N_{5+6}(z + b) < N_{5+6}(z)$ and $z + b \in \mathcal{F}_{\theta P}$.

Case 2. Assume that a_2 is of type C6 with either

$$F(a_2) = 1(Sx, \bar{T}y, w) - 1(Sx, Ty, w) \text{ or } F(a_2) = 1(Sx, Pz, w) - 1(Sx, \bar{P}z, w).$$

Then either $a_2 = \pm E(Sx, y, w)$ or $a_2 = \pm E(Sx, z, w)$. In the first (resp. second) case, we may assume, by lemma 3.9, that $a_1 = -F_3(Sx, Pz, Ty, w)$ (resp. $F_3(Sx, Ty, Pz, w)$), and then lemma 3.10 gives a boundary b with $z + b \in \mathcal{F}_{\theta P}$ and $N_{5+6}(z + b) < N_{5+6}(z)$. $\square$

Lemma 3.12. *Each $z \in \mathcal{F}_{C5}$ is homologous to a cycle in $\mathcal{F}_{C4}$.*

Proof. Let c_1 and c_2 be the generators indicated in figure 11. It's easy to check that $d(c_1 + c_2)$ consists of two cubical terms of type C2(a), two prismatic terms (neither of which is θ -prismatic), and one cubical term of type C4. $\square$

Lemma 3.13. *Each $z \in \mathcal{F}_{C4}$ is homologous to a cycle in $\mathcal{F}_{C3}$.*

Proof. Let $g \in C_4(\tilde{X}_n/StN_n)$ correspond to the colored graph shown on the right of figure 11 with either cell orientation. Then it's routine to check that there is a chain $c \in C_4(\tilde{X}_n/StN_n)$ consisting of 2,1,1-terms such that $d(g + c)$ is in $\mathcal{F}_{C4}$ and contains exactly one term of type C3. Indeed, we can take $c = \sum_{i=1}^4 c_i$ where the colored graphs $cg(c_i)$, for $i = 1, 2$ and 3, are defined by separating edges incident with the 4-valent vertices of the three colored graphs corresponding to the terms of dg containing such vertices. Then just one cubical term of $d(g + \sum_{i=1}^3 c_i)$ corresponds to a colored graph with a 4-valent vertex, and c_4 is defined by separating edges incident with it. $\square$




FIGURE 12.

For the next lemma we will use a homomorphism $r_3: C_2(\tilde{X}_n/StN_n) \rightarrow \bigoplus_{\substack{1 \leq a,b,c,d \leq n \\ S,T \in \{L,R\}}} \mathbb{Z}_{\{Sa/b, Tc/d\}}$ which counts trivial commutator relations. Here each $\mathbb{Z}_{\{Sa/b, Tc/d\}}$ is a copy of $\mathbb{Z}$ with distinguished generator $1\{Sa/b, Tc/d\} = 1\{Tc/d, Sa/b\}$. Let $g' \in C_2(\tilde{X}_n/StN_n)$ be the generator indicated on the left of figure 12 where, if $h_a = \{a_-, a_+\}$ and $h_b = \{b_-, b_+\}$, we are assuming that $a < b$, and where $A = \{\alpha_i\}$ and $B = \{\beta_j\}$ are the indicated sets of half-edges. The homomorphism r_3 is defined on generators by

$$r_3(x) = \begin{cases} \sum_{i,j} 1\{\alpha_i/a, \beta_j/b\}, & \text{if } x = g' \\ 0, & \text{if } x \neq \pm g'. \end{cases}$$

Let R_3 be the composite $\mathcal{F}_{C3} \hookrightarrow C_3(\tilde{X}_n/StN_n) \xrightarrow{d} C_2(\tilde{X}_n/StN_n) \xrightarrow{r_3} \bigoplus \mathbb{Z}$. Then R_3 is trivial on simplicial and prismatic terms as can be checked, and is only nontrivial on generators of type C2.

More preliminaries to the proof of the next lemma are needed. View $\tilde{X}/StN$ as the direct limit of the spaces $\tilde{X}_S/StN_S$ where S varies over all finite subsets of $\mathbb{N} = \{1, 2, \dots\}$. Given a generator g of $C_*(\tilde{X}_n/StN_n)$, let $EC(g)$, the essential colors of g , be the subset of $\mathbb{N}$ of minimal cardinality such that there is an element $E(g) \in C_*(\tilde{X}_{EC(g)}/StN_{EC(g)})$ agreeing with g in $C_*(\tilde{X}_n/StN_n)$. Given generators $g_1, g_2 \in C_*(\tilde{X}_n/StN_n)$ with $EC(g_1) \cap EC(g_2) = \emptyset$, let $g_1 \vee g_2$ be the generator of $C_*(\tilde{X}_n/StN_n)$ corresponding to the one point union of $cg(E(g_1))$ and $cg(E(g_2))$ with the natural coloring and with cell orientation that of g_1 followed by that of g_2 .

Let $g = F_3(Sx, Aw, Ty, z) \vee F_1(Qu, v)$. Then there is a chain $c \in C_4(\tilde{X}_n/StN_n)$, obtained by separating edges as in the proof of lemma 3.8, such that

$$(8) \quad d(g + c) \equiv -(F_2(Sx, Ty, z) \vee F_1(Qu, v)) + F_2(Sx, Aw, z) \vee F_1(Qu, v) \pmod{\mathcal{F}_{C2}}.$$

Now let g, c_1 and c_2 be the generators of $C_4(\tilde{X}_n/StN_n)$ indicated in figure 12. Then, with appropriate cell orientations, we have

$$(9) \quad d(g + c_1 + c_2) \equiv F_2(Sx, Ty, z) \vee F_1(Qu, v) - F_2(Sx, \bar{Ty}, z) \vee F_1(Qu, v) \pmod{\mathcal{F}_{C1}}.$$

Finally, let $g = F_2(Sx, Ty, z) \vee F_2(Qu, Aw, v)$. Then there are generators c_1 and c_2 in $C_4(\tilde{X}_n/StN_n)$ (obtained by separating edges of colored graphs corresponding to terms of dg) such that

$$(10) \quad d(g + c_1 + c_2) \equiv -F_1(Sx, z) \vee F_2(Qu, Aw, v) - F_2(Sx, Ty, z) \vee F_1(Qu, v) \pmod{\mathcal{F}_{C1}}.$$

Lemma 3.14. *Each $z \in \mathcal{F}_{C3}$ is homologous to a cycle in $\mathcal{F}_{C2}$.*

Proof. Let $z \in \mathcal{F}_{C3}$. We show that there is a 4-chain x such that $z + dx$ is in $\mathcal{F}_{C3}$ and contains fewer type C2 terms than z . Write $z = \sum_i z_i$.

Case 1. Assume some z_i , say z_1 , is of type C2(a) with

$$z_1 = \pm(F_2(Sx, Ty, z) \vee F_1(Qu, v)).$$

Then $R_3(z_1) = \mp 1\{Sx/z, Qu/v\}$. So, since $R_3(z) = 0$, there is some other z_i , say z_2 , with $R_3(z_2) = -R_3(z_1)$. Since $x \neq u$, either

$$(11) \quad z_2 = \mp(F_2(Sx, Aw, z) \vee F_1(Qu, v))$$

or

$$(12) \quad z_2 = \pm(F_1(Sx, z) \vee F_2(Qu, Aw, v))$$

for some $A \in \{L, R\}$ and positive integer w . If (11) holds and $w \neq y$, use the congruence (8) to decrease the number of type C2 terms of z by two, while keeping $z \in \mathcal{F}_{C3}$. If $w = y$ and $A \neq T$, use (9) to do the same. If (12) holds, then by (8), we may assume that $w \neq y$. Then use (10).

Case 2. If z has a type C2(b) term, an argument like that of the previous case can be used. The colored graphs corresponding to the three generators denoted by g in (8), (9) and (10) should be modified so that the two singleton edges of each of these are replaced with a single singleton edge. $\square$

The two cases in the previous proof could easily be combined, but then the wedge notation (which eliminates the need for several figures) could not have been easily used.

Lemma 3.15. *Each $z \in \mathcal{F}_{C2}$ is homologous to a cycle in $\mathcal{F}_{C1}$.*

Proof. If x is a generator of $C_3(\tilde{X}_n/StN_n)$ of type C1, then $x = g \vee F_1(Sa, b)$ for some $g \in C_2(\tilde{X}_n/StN_n)$. Let w be a color distinct from the five or six essential colors making up the set $E(x)$. Then there is a generator c , obtained, as usual, by separating edges, such that $d(g \vee F_2(Sa, Sw, b) + c) = -x$. $\square$

4. $\mathbb{R}^n$ IN X_k

Let $k = 2n - 2$. In this brief section, which will be used at the end of the next, we define a subgroup $\mathbb{Z}_*^k$ of $\text{Aut}(F_n)$ isomorphic to $\mathbb{Z}^k$, the free abelian group of rank k , and also define a subcomplex $\mathbb{R}_*^k$ of X_n . The group $\mathbb{Z}_*^k$ acts on $\mathbb{R}_*^k$ in the usual way, and thus so does $\mathbb{Z}^k$, via an isomorphism $\mathbb{Z}^k \approx \mathbb{Z}_*^k$ we specify. We also give $\mathbb{R}^k$ a standard triangulation and the usual $\mathbb{Z}^k$ -action, and then prove the following.

Lemma 4.1. *There is a $\mathbb{Z}^k$ -equivariant simplicial homeomorphism from $\mathbb{R}^k$ to $\mathbb{R}_*^k$. Moreover, the classifying space $\mathbb{R}_*^k/\mathbb{Z}_*^k$ for $\mathbb{Z}^k$ is a subcomplex of $X_n/\text{Aut}(F_n) = \tilde{X}_n/StN_n$.*

Assume that F_n is free on the set $\{x_1, \dots, x_n\}$. The generators of $\mathbb{Z}_*^k$ are, by definition, the k automorphisms α_i and β_i , for $1 \leq i \leq n - 1$, defined by $\alpha_i(x_i) = x_n x_i$, $\beta_i(x_i) = x_i x_n$, and $\alpha_i(x_j) = x_j = \beta_i(x_j)$ if $i \neq j$. We identify $\mathbb{Z}^k$ with $\mathbb{Z}_*^k$ by letting $(m_1, \dots, m_k) \in \mathbb{Z}^k$ correspond to the automorphism $\alpha_1^{m_1} \beta_1^{-m_2} \dots \alpha_{n-1}^{m_{k-1}} \beta_{n-1}^{-m_k}$.

A cell of X_n is, by definition, in $\mathbb{R}_*^k$ if it corresponds to some blowdown of a colored graph of the sort shown in figure 13(a) (where π is any permutation of the set $\{1, \dots, k\}$) with marking such that any (and hence every) 0-cell in the boundary of the cell corresponds to an




FIGURE 13.

automorphism of $\mathbb{Z}_*^k$. Thus $\mathbb{R}_*^k$ consists of those cells whose 0-cells all correspond to elements of $\mathbb{Z}_*^k$.

The unit cube $[0, 1]^k \subset \mathbb{R}^k$ is triangulated by the $k!$ k -simplices $\sigma_{\pi(1)} \times \dots \times \sigma_{\pi(k)}$ where π ranges over all permutations of the set $\{1, \dots, k\}$ and $\sigma_i = (0, \dots, 0, 1, \dots, 1)$ is the (degenerate) k -simplex with i zeros and $k + 1 - i$ ones. (Here we are using the notation of, for instance, p. 244 of Mac lane [10].) Translation of this triangulation by the elements of $\mathbb{Z}^k$ defines a $\mathbb{Z}^k$ -invariant triangulation of all of $\mathbb{R}^k$.

proof of lemma 4.1. Let $[a_0, \dots, a_k]_{\pi, id}$ be the k -simplex of X_n corresponding to the colored graph shown in figure 13(a) with marking specified by $[a_0] = id \in \text{Aut}(F_n)$. Mapping, for each permutation π , the k -simplex $\sigma_{\pi(1)} \times \dots \times \sigma_{\pi(k)}$ of the cube $[0, 1]^k$ to $[a_0, \dots, a_k]_{\pi, id}$ so that the ordering of the vertices is preserved, and then extending equivariantly defines the desired homeomorphism.

Now assume that $g(a) = g(b)$ where $g: \mathbb{R}_*^k / \mathbb{Z}_*^k \rightarrow X_n / \text{Aut}(F_n)$ is the map induced by the inclusion $\mathbb{R}_*^k \rightarrow X_n$. Then there are points $a', b' \in \mathbb{R}_*^k \subset X_n$ representing a and b and some $f \in \text{Aut}(F_n)$ with $f a' = b'$. Since f acts cellularly, there are vertices v and w of $\mathbb{R}_*^k$ with $f v = w$. Since v and w correspond to elements of $\mathbb{Z}_*^k$, so does f . Thus g is injective. $\square$

Remark 4.2. While the number n plays a special role in the definitions of $\mathbb{Z}_*^k$ and $\mathbb{R}_*^k$, these could just as well be defined using any of $1, \dots, n - 1$ in place of n , and the lemma would still be true.

4.1. Let $\mathbb{R}_{i*}^k$ denote the subcomplex of $\tilde{X}_n / \text{St}N_n$ analogous to $\mathbb{R}_*^k = \mathbb{R}_{n*}^k$, but defined with i in place of n . We conclude this section with an explicit description of a basis for $H_2(\mathbb{R}_{i*}^k / \mathbb{Z}^k)$ which will be used in Lemmas 5.8 and 5.10. Order the half-edges of any colored graph by $Si < Tj$ if either $i < j$ or, if $i = j$, $S = L$ and $T = R$. Let $S_i(\alpha, \beta)$ (resp. $S_i(\beta, \alpha)$) be the generator of $C_2(\mathbb{R}_{i*}^k / \mathbb{Z}^k)$ indicated in figure 13(b) (resp. figure 13(c)) where we are assuming that $\alpha < \beta$. Denote by $1_i\{\alpha, \beta\} = 1_i\{\beta, \alpha\}$ the element of $H_2(\mathbb{R}_{i*}^k / \mathbb{Z}^k)$ which the cycle $S_i(\alpha, \beta) + S_i(\beta, \alpha)$ represents. It follows from the computation of $H_2(\mathbb{Z}^k)$ and lemma 4.1 that the $1_i\{\alpha, \beta\}$ for all distinct half-edges α and β different from Li and Ri form a basis for $H_2(\mathbb{R}_{i*}^k / \mathbb{Z}^k)$.

5. ELIMINATING PRISMATIC AND SIMPLICIAL TERMS

We say that a colored graph G has type Pn , for $n \neq 2$, if the core of G is isomorphic to the core of the partitioned graph labeled Pn in figure 14. Type P2(a), (b) and (c) colored




FIGURE 14. Near classification of prismatic colored graphs.

graphs are defined similarly, and a colored graph is of type P2 if it is of type P2(a), (b) or (c).

In order to eliminate all prismatic terms from z , we extend the filtration $\mathcal{F}$ of the group $Z_3(\tilde{X}_n/StN_n)$ of cycles to

$$\mathcal{F}_{C1} \supset \mathcal{F}_{P5} \supset \mathcal{F}_{P4} \supset \dots \supset \mathcal{F}_{P1} \supset \mathcal{F}_S$$

where $z \in \mathcal{F}_{Pn}$ if none of the terms of z correspond to colored graphs of type Pm , for $m \geq n$. (An additional technical requirement is placed on the elements of $\mathcal{F}_{P4}$ in Def. 5.4 below.) Also, $z \in \mathcal{F}_S$ if none of the terms of z are simplicial. Lemmas 5.3 to 5.10 below show that any cycle in a subgroup of the above filtration is homologous in the space $\tilde{X}_n/StN_n$ to a cycle in the next smaller subgroup of the filtration. Since $\mathcal{F}_{C1} \subset \mathcal{F}_{\theta P}$, it follows from the next lemma that the smallest subgroup $\mathcal{F}_S$ of the filtration is the trivial group. Thus, every element of $\mathcal{F}_{C1}$ is a boundary.

Lemma 5.1. *If G is a prismatic colored graph, then either G is of one of the types shown in figure 14, or G is θ -prismatic.*

θ -prismatic graphs were defined just before Lemma 3.7 above.

Proof. Let G be any prismatic colored graph. Since G has four vertices, the three edges of G of the same color form a connected subgraph of G . If they form a monochromatic loop, then, disregarding basepoints, the core of G is as shown in either P2(a) or P2(b) of figure 14. Otherwise the core is as shown in P5 of figure 14. Each of the possible locations for the basepoint on these three cores gives either one of the graphs in figure 14 or a θ -prismatic graph. $\square$

Definition 5.2. A colored graph G with three vertices is of type θ (resp. x , y , w) if the core of G is equivalent, for some choice of colors, to the core of the graph labeled type θ (resp. x , y , w) in figure 15. The incidence number of a colored graph of any of the three types θ , x , or y is the cardinality of the set of half-edges corresponding to the set labeled T in the appropriate graph in the figure. The incidence number of a type w colored graph is $|T| + |F|$, where T and F are the sets analogous to those in the figure. The incidence number of a generator g of $C_2(\tilde{X}/StN)$ with $cg(g)$ of type x , y , w or θ is the incidence number of $cg(g)$.

Let G be a colored graph of either type P5 or θ . The initial vertex of the first edge of the (oriented) edgeset through the basepoint is called the initial vertex of G . We say that G is reduced if its initial vertex is trivalent.

Lemma 5.3. *Each $z \in \mathcal{F}_{C1}$ is homologous in $\tilde{X}_n/StN_n$ to some element of $\mathcal{F}_{P5}$.*




FIGURE 15.




FIGURE 16.

Proof. By adding to z generators of $C_4(\tilde{X}_n/StN_n)$ corresponding to (suitably oriented) colored graphs of the sort shown in figure 16(a) and (b), we may assume that, for each type P5 term z_i of z , $cg(z_i)$ is reduced.

Given a type θ colored graph G which is not reduced, there is exactly one reduced colored graph of type P5 having a blowdown equal to G . Thus, if x is a term of dz with $cg(x)$ of type θ , then $cg(x)$ is reduced. It follows that all the type P5 terms of z correspond to colored graphs of the sort shown in figure 16(c) where B , but not A , may be empty. Such colored graphs are said to have type $P5^+$. We denote by $\theta_{pq}(A, B)$ the colored graph in figure 16(c) and by $\theta_{pq}(A, B, C)$ the one in figure 16(d). Here q is the color of the edgesets shown as dotted and p is the color of the edges in the other edgeset with more than one edge.

Assuming $z \notin \mathcal{F}_{P5}$, let $m(z)$ be the maximal incidence number of all the type θ faces of the terms of z . Choose two terms z_1 and z_2 of z such that dz_1 contains a term which realizes $m(z)$ and cancels with a term of dz_2 . Then both z_1 and z_2 correspond to type $P5^+$ colored graphs, and if $cg(z_i) = \theta_{pq}(A_i, B_i)$ for $i = 1$ and 2 , then the set $A_1 \cup B_1 = A_2 \cup B_2$ has cardinality $m(z)$.

If one of the $B_i = \emptyset$, say B_1 , then the boundary of the generator of $C_4(\tilde{X}_n/StN_n)$ corresponding to the suitably oriented colored graph $\theta_{pq}(A_2, B_2, \emptyset)$ is $z_1 + z_2$ plus two more terms. One of these is of type P3. The θ -faces of the other, which is of type $P5^+$, all have incidence number $\leq |B_2| < m(z)$. Thus, if a $B_i = \emptyset$, the total number of terms of z with a face realizing $m(z)$ can be decreased.

Now we assume that neither $B_i = \emptyset$ and show that this is still true. We first assume that $m(z) > 2$. Then if, say $|A_1| > 1$, any proper subset a of A_1 can effectively be deleted from A_1 and added to B_1 by adding to z the boundary of the generator of $C_4(\tilde{X}/StN)$ corresponding to the (suitably oriented) colored graph G shown in figure 17(a). This may also add to z a type P3 term corresponding to G_x and a type $P5^+$ one corresponding to G_w . But, again, the two θ -faces of this last term have incidence number $\leq |a| + |B_1| < m(z)$. Similarly any proper subset of B_1 can be moved from B_1 to A_1 without increasing $m(z)$. By using either




FIGURE 17.

one, two or three of these moves it's possible to eliminate z_1 and z_2 from z . This reduces the total number of terms of z which have a type θ face realizing $m(z)$. Thus, if $m(z) > 2$, the number of such terms can be reduced to zero, thereby reducing $m(z)$.

So now we assume that $m(z) = 2$ and, by the above, that neither $B_i = \emptyset$. Then $A_1 = B_2$ and $B_1 = A_2$, and, with suitable orientations,

$$z_1 + z_2 + d(\text{gen}(\theta_{pq}(A_1, B_1, \emptyset)) - \text{gen}(\theta_{pq}(B_1, A_1, \emptyset)))$$

has just two terms. One is of type $P3$. The θ -faces of the other, which is of type $P5^+$, have incidence number ≤ 1 . So $m(z)$ can also be reduced if it equals 2. This completes the proof since $m(z)$ cannot equal 1. $\square$

Definition 5.4. A type $P2$ colored graph G is reduced if all its type x and y faces have incidence number one. All type $P2$ terms of elements of $\mathcal{F}_{P4}$ are required to be reduced.

Lemma 5.5. Each $z \in \mathcal{F}_{P5}$ is homologous to a cycle in $\mathcal{F}_{P4}$.

Proof. If a type $P2$ colored graph G is drawn as in P2(a), (b) or (c) of figure 14, then G is reduced if the set of half-edges which are not incident with any dotted edge has cardinality one. We may assume that all the $P2$ terms of z are reduced by repeatedly separating the half-edges of any of these sets which have cardinality > 1 .

Assume $z \notin \mathcal{F}_{P4}$. The argument that z is homologous to a cycle in $\mathcal{F}_{P4}$ is similar to parts of the proof of lemma 5.3. Let $m(z)$ now be the maximal incidence number of all the type x faces of the terms of z . Let z_1 and z_2 be two terms of z having a common canceling face which realizes $m(z)$. Since the $P2$ terms of z are reduced, both of the colored graphs $cg(z_1)$ and $cg(z_2)$ are of type $P4$. For $i = 1$ and 2 , we denote by A_i the set of half-edges of $cg(z_i)$ corresponding to the set labeled A of the graph in the lower right of figure 18, and by B_i the one corresponding to B . Then $A_1 \cup B_1 = A_2 \cup B_2$, and this set has cardinality $m(z)$. If $m(z) > 2$ then the argument in the proof of 5.3 showing $m(z)$ can be reduced is now applicable with G the colored graph in figure 17(b) rather than 17(a). Note that all the type x and y faces of G_x and G_w have incidence number $< m(z)$.

So we now assume that $m(z) = 2$. Then, using the above notation, we have $A_1 = B_2$ and $B_1 = A_2$, and we assume that $A_1 = \{\alpha\}$ and $B_1 = \{\beta\}$. Figure 18 indicates a 4-chain c with four terms such that $z_1 + z_2 \pm dc \in \mathcal{F}_{P1}$ for one of the choices of the signs. (Each graph in the figure is identical to the one above or below it except for the orientation of the edgeset $\{x, y, z\}$. The two colored graphs on the right with the indicated orientations correspond to $\pm(z_1 + z_2)$.) This completes the proof. $\square$




FIGURE 18.




FIGURE 19.

Lemma 5.6. *Each $z \in \mathcal{F}_{P_4}$ is homologous to a cycle in $\mathcal{F}_{P_3}$.*

Proof. Assuming $z \notin \mathcal{F}_{P_3}$, let $m(z)$ now denote the maximal incidence number of all the type y faces of the terms of z . The argument of the previous lemma with some changes shows that we may assume that $m(z) = 2$. These changes include using type y colored graphs in place of type x ones, using the graph in figure 17(c) (with the edgeset $\{x, y, z, w\}$ suitably oriented) instead of the one in figure 17(b), and substituting type P_3 for P_4 .

We will use a homomorphism $f: C_2(\tilde{X}_n/StN_n) \rightarrow \oplus_i \mathbb{Z}_i$ where there is one $\mathbb{Z}_i \approx \mathbb{Z}$ summand in the range corresponding to each colored graph shown in figure 19(a). Here Sa and Tb are half-edges so that $S, T \in \{L, R\}$. Also a, c and d are distinct colors, as are b, c and d . We let $1\{Sa/c, Tb/d\} = 1\{Tb/d, Sa/c\}$ denote a generator of the $\mathbb{Z}$ summand corresponding to the colored graph in figure 19(a). Let $G_1^\pm$, G_2 and G_3 be the colored graphs indicated in figure 19 where we assume $A = \{\alpha_i\}$ and $B = \{\beta_i\}$. Let $g_1^\pm = \text{gen}(G_1^\pm)$ and $g_i = \text{gen}(G_i)$ be the corresponding generators of $C_2(\tilde{X}_n/StN_n)$ with the orientations of $G_1^\pm$ and the G_i given by $[c_1, c_2][d_1, d_2]$ if $c < d$, and opposite this if $c > d$. Let $G^\pm$ be the two colored graphs




FIGURE 20.

indicated in figure 20 where α and β are single half-edges. Let $g_{cd}^{\pm}(\alpha, \beta, E) = \text{gen}(G^{\pm})$ be the generators of $C_3(\tilde{X}_n/StN_n)$ corresponding to G^+ and G^- , both with the orientation $[c_1, c_2, c_3][d_1, d_2]$.

The homomorphism f is defined by

$$f(g_1^{\pm}) = f(g_2) = f(g_3) = \begin{cases} \sum_{i,j} 1\{\alpha_i/c, \beta_j/d\} & \text{if } B \neq \emptyset, \\ 0 & \text{if } B = \emptyset, \end{cases}$$

and $f(g) = 0$ for all other generators g of $C_2(\tilde{X}_n/StN_n)$. It's routine to check that $f(d(g_{cd}^{\pm}(\alpha, \beta, E))) = 1\{\alpha/c, \beta/d\}$ where $d: C_3(\tilde{X}_n/StN_n) \rightarrow C_2(\tilde{X}_n/StN_n)$ is the boundary homomorphism. Also, if a generator g of $C_3(\tilde{X}_n/StN_n)$ is simplicial or of type $P1$ or $P2$, then $f(d(g)) = 0$.

As usual let z_1 and z_2 be terms of z with a common canceling face realizing $m(z) = 2$. The colored graph corresponding to this face is either G_1^+ or G_1^- , say it's G_1^+ (with a similar argument if it's G_1^-). Then, since the $P2$ terms of z are reduced,

$$z_1 + z_2 = g_{cd}^+(\alpha, \beta, E) - g_{cd}^+(\beta, \alpha, E)$$

for some $c, d \in \mathbf{n}$, half-edges α and β , and set of half-edges E . Let $g_4^+(\alpha, \beta, E)$ be the generator of $C_4(\tilde{X}_n/StN_n)$ corresponding to the colored graph shown in figure 20(a) with orientation $[c_1, c_2, c_3][d_1, d_2, d_3]$. Then

$$z_1 + z_2 - (g_{cd}^+(\alpha, \beta, \emptyset) - g_{cd}^+(\beta, \alpha, \emptyset)) + d(g_4^+(\alpha, \beta, E) - g_4^+(\beta, \alpha, E)) \in \mathcal{F}_{P2}$$

with all the $P2$ terms reduced. So we assume that $E = \emptyset$.

Since $f(d(z_1 + z_2)) = 1\{\alpha/c, \beta/d\} - 1\{\beta/c, \alpha/d\} \neq 0$ and z is a cycle, z must contain a term, z_3 say, distinct from z_1 and z_2 , such that $f(d(z_3))$ cancels with a term of $f(d(z_1 + z_2))$. Let z_4 be a term of z with a face canceling the one of z_3 which realizes $m(z) = 2$. Then, for some set of half-edges E' , either

$$(13) \quad z_3 + z_4 = -g_{cd}^+(\alpha, \beta, E') + g_{cd}^+(\beta, \alpha, E')$$

or

$$(14) \quad z_3 + z_4 = -g_{cd}^-(\alpha, \beta, E') + g_{cd}^-(\beta, \alpha, E').$$

In both cases we may assume, as above, that $E' = \emptyset$. Then if (13) holds, $\sum_{i=1}^4 z_i = 0$. To treat the other case, let $s_4(\alpha, \beta)$ be the generator of $C_4(\tilde{X}_n/StN_n)$ corresponding to the colored graph shown in figure 20(b) with orientation $[c_1, c_2, c_3][d_1, d_2, d_3]$. Then, if (14) holds,




FIGURE 21.

we have

$$d(s_4(\alpha, \beta) - s_4(\beta, \alpha) \pm c) - \sum_{i=1}^4 z_i \in \mathcal{F}_{P2}$$

for a 4-chain c like that indicated in figure 18. Moreover, all the $P2$ terms are reduced. So in both cases the number of terms of z realizing $m(z) = 2$ can be reduced. $\square$

Definition 5.7. The lower incidence number of a type x (resp. y) colored graph with set E as shown in figure 15 is $|E| + 2$ (resp. $|E| + 1$). The lower incidence number of a generator g of $C_2(\tilde{X}/StN)$ with $cg(g)$ of type x or y is the lower incidence number of $cg(g)$.

Lemma 5.8. Each $z \in \mathcal{F}_{P3}$ is homologous to a cycle in $\mathcal{F}_{P2}$.

Proof. Terms of z of type $P2(a)$ can be exchanged for simplicial and reduced type $P2(b)$ and (c) ones by using colored graphs as shown in figure 16(e). Thus, we assume that none of the terms of z are of type $P2(a)$.

If z contains any terms of type $P2(b)$, let $m(z)$ be the maximal lower incidence number of all the type x faces of the terms of z . We may assume that $m(z) = 2$ by using the usual argument, this time with colored graphs as, for instance, shown in figures 17(d) and 16(e). Let z_1 and z_2 be two type $P2(b)$ terms of z with faces realizing $m(z) = 2$. We may assume that the colored graphs $cg(z_1)$ and $cg(z_2)$ are identical except for the orientations of the edgesets with two edges. Then, for one choice of the plus or minus sign,

$$dy \pm (z_1 + z_2)$$

consists of just reduced type $P2(c)$ terms where y is the sum of the seven generators of $C_4(\tilde{X}_n/StN_n)$ indicated in figure 21 with suitable choices of colors. So we now assume that z contains no terms of either type $P2(a)$ or (b).

The following notation will be used in the remainder of the proof. Let $F_i^+(Ra, \alpha, \beta)$ (resp. $-F_i^-(Ra, \alpha, \beta)$) be the generator of $C_4(\tilde{X}_n/StN_n)$ corresponding to the colored graph and orientation in the upper left (resp. lower left) of figure 22. Here we are assuming that $h_a = \{x, y\}$ and $h_i = \{s, t, u\}$. Define $F_i^+(La, \alpha, \beta)$ (resp. $F_i^-(La, \alpha, \beta)$) as $F_i^+(Ra, \alpha, \beta)$ (resp. $F_i^-(Ra, \alpha, \beta)$) is defined, but switch the orientation of the edgeset h_a .




FIGURE 22.

If z contains any terms of type $P2(c)$, then, as usual, we may assume that $m(z) = 2$ where now $m(z)$ is the maximal lower incidence number of all the type y faces of the terms of z . Write

$$z = \rho + \sum_{i=1}^n \zeta_i$$

where ρ contains all the type $P1$ terms of z , and ζ_i contains all the type $P2(c)$ and simplicial terms corresponding to colored graphs whose edgesets with cardinality > 2 have color i . Choose a $q \in \mathbf{n}$ such that ζ_q contains a type $P2(c)$ term. Write

$$\zeta_q = s + \sum_j (z_j + z'_j)$$

where s contains all the simplicial terms of ζ_q and where, for each j , z_j and z'_j have a common canceling face realizing $m(z)$. Since $m(z) = 2$, we have, for each j ,

$$(15) \quad z_j + z'_j = \pm(F_q^+(S_j a_j, \alpha_j, \beta_j) - F_q^-(S_j a_j, \alpha_j, \beta_j))$$

for some $S_j \in \{L, R\}$, $a_j \in \mathbf{n}$, and half-edges α_j and β_j , and where the choice of the plus or minus sign depends on j . The left part of figure 22 indicates that each $d(z_j + z'_j)$ is in the image of the homomorphism $C_2(\mathbb{R}_{q*}^k/\mathbb{Z}^k) \xrightarrow{i_*} C_2(\tilde{X}_n/StN_n)$. Here $\mathbb{R}_{q*}^k/\mathbb{Z}^k$ is the subcomplex of $\tilde{X}_n/StN_n$ defined in §4.1 and i is the inclusion. So $d\zeta_q$ is also in the image of i_* . By a similar argument, each $d\zeta_i$ is in the image of the map $C_2(\mathbb{R}_{i*}^k/\mathbb{Z}^k) \rightarrow C_2(\tilde{X}_n/StN_n)$ induced by inclusion. Thus, since all the terms of $d\rho$ correspond to square 2-cells, each ζ_i is a cycle, as is ρ .

The 3-chain s is itself in the image of $C_3(\mathbb{R}_{q*}^k/\mathbb{Z}^k) \xrightarrow{i_*} C_3(\tilde{X}_n/StN_n)$. Thus ds , viewed as a 2-chain in $\mathbb{R}_{q*}^k/\mathbb{Z}^k$, represents the trivial element of $H_2(\mathbb{R}_{q*}^k/\mathbb{Z}^k)$. Since ζ_q is a cycle, $d\left(\sum_j (z_j + z'_j)\right)$, viewed in the same way, also does. But, with the notation of §4.1, we have




FIGURE 23.




FIGURE 24.

$d(z_j + z'_j) = \pm 1_q \{\alpha_j, \beta_j\} \in H_2(\mathbb{R}_{q*}^k / \mathbb{Z}^k)$, as figure 22 indicates. Thus, there are two values of j , say $j = 1$ and $j = 2$, such that $d(z_1 + z'_1 + z_2 + z'_2) = 0 \in H_2(\mathbb{R}_{q*}^k / \mathbb{Z}^k)$. So, with the notation introduced in (15), either

$$(16) \quad \alpha_1 = \alpha_2 \quad \text{and} \quad \beta_1 = \beta_2,$$

or

$$(17) \quad \alpha_1 = \beta_2 \quad \text{and} \quad \beta_1 = \alpha_2.$$

To treat these two cases, we introduce two elements c_1 and c_2 of $C_4(\tilde{X}_n / StN_n)$. Assume that the finely dotted edges (resp. other dotted edges, solid edges) of the graphs in figure 23 have color a (resp. b , q). Let c_1 be the sum of the seven generators indicated in figure 23, plus three more corresponding to the three graphs on the right of the figure, but with the colors a and b switched, and also with the cell orientations of all three graphs switched. Then, for any consistent choice of orientations of the edgesets a and b , we have

$$(18) \quad dc_1 - (F_q^+(Sa, \alpha, \beta) - F_q^-(Sa, \alpha, \beta) - F_q^+(Tb, \alpha, \beta) + F_q^-(Tb, \alpha, \beta)) \in \mathcal{F}_{P2}$$

where $S, T \in \{L, R\}$.

With the same coloring scheme, let c_2 be the sum of the two generators of $C_4(\tilde{X}_n / StN_n)$ indicated in figure 24, plus two more like these but with the orientations of the edgesets q reversed, and also with the cell orientations reversed, plus three more like those on the right of figure 23, plus another five. These five more are similar to the five 2,4 terms of c_2 already described, but with both the colors a and b and the half-edges α and β interchanged. Then,




FIGURE 25.

with cell orientations chosen properly,

$$(19) \quad dc_2 - (F_q^+(Sa, \alpha, \beta) - F_q^-(Sa, \alpha, \beta) - F_q^+(Tb, \beta, \alpha) + F_q^-(Tb, \beta, \alpha)) \in \mathcal{F}_{P_2}.$$

Now, in view of (19), and (15) with $j = 1$ and 2 , we may assume that (16) holds, and not (17). Also, because of (18), we may assume that $a_1 \neq a_2$. Then adding to z one of $\pm dc_1$ with $a = a_1$ and $b = a_2$ decreases by four the number of terms of z with faces realizing $m(z) = 2$. This only adds type P1 and simplicial terms to z , and so completes the proof. $\square$

Lemma 5.9. *Each $z \in \mathcal{F}_{P_2}$ is homologous to a cycle in $\mathcal{F}_{P_1}$.*

Proof. Let $m(z)$ be the maximal incidence number of all the type w faces of the terms of z . By using colored graphs like the two on the left of figure 25, we may assume that $m(z) = 3$. More specifically, assume z_1 and z_2 are two terms of z with canceling type w faces realizing $m(z) > 3$. If $cg(z_1)$ and $cg(z_2)$ have monochromatic loops of length two of different colors, then there is a generator g with $cg(g)$ as shown in the lower left of figure 25 such that $m(z_1 + z_2 + dg) < m(z_1 + z_2)$. If the monochromatic loops have the same color then adding one or two generators corresponding to the colored graph in the upper left achieves the same result.

Now the type P1 terms of z can be paired off, with the elements of each pair corresponding to colored graphs and orientations like those on the right of figure 25. The central portion of the figure shows that these are the boundary of a 4-chain. $\square$

Lemma 5.10. *Each $z \in \mathcal{F}_{P_1}$ is homologous to $0 \in C_3(\tilde{X}_n/StN_n)$.*

Proof. Write $z = \sum_{i=1}^n \zeta_i$ where each ζ_i is a cycle in the subcomplex $\mathbb{R}_{i*}^k/\mathbb{Z}^k$ of $\tilde{X}_n/StN_n$. Each basis element of $H_2(\mathbb{R}_{i*}^k/\mathbb{Z}^k)$ described in §4.1 is trivial in $H_2(\tilde{X}_n/StN_n)$ as figure 22 indicates. Thus, each ζ_i , and so also z , is trivial in $H_2(\tilde{X}_n/StN_n)$. $\square$

Given the remarks preceeding lemmas 3.2 and 5.1, this completes the proof of Theorem 3.1.

6. COMPUTATION OF $H_2(\tilde{X}_n/StN_n)$

This section is devoted to the proof of the following.

Theorem 6.1. *The second homology group $H_2(\tilde{X}_n/StN_n)$ is free abelian of rank n provided $n \geq 5$.*

We assume throughout this section that $n \geq 5$, although this is only needed in Lemma 6.9. Let $\mathbb{Z}^n$ be the free abelian group of rank n with basis $\{e_1, \dots, e_n\}$. We will define a homomorphism $\Theta': C_2(\tilde{X}_n/StN_n) \rightarrow \mathbb{Z}^n$ which roughly counts how many terms of any given chain correspond to a colored graph which contains one of the colored graphs θ_{ij} as a subgraph. (The colored graph θ_{xy} is shown on the left of Figure 6 in §3.) The lemmas in this section will show that Θ' induces an isomorphism $\Theta: H_2(\tilde{X}_n/StN_n) \rightarrow \mathbb{Z}^n$, thereby proving the theorem.

The homomorphism Θ' is defined on oriented generators x of $C_2(\tilde{X}_n/StN_n)$ by

$$\Theta'(x) = \begin{cases} 0, & \text{if } \text{core}(cg(x)) \text{ is not a } \theta_{ij}, \\ e_j, & \text{if } \text{core}(cg(x)) = \theta_{ij} \text{ with the same cell orientations.} \end{cases}$$

In the second line of the definition, we are assuming the cell orientation of θ_{ij} is as shown in the upper left of Figure 6.

Lemma 6.2. *The homomorphism Θ' vanishes on the image of the cellular boundary map $\partial: C_3(\tilde{X}_n/StN_n) \rightarrow C_2(\tilde{X}_n/StN_n)$.*

Proof. If G is a colored graph with $\text{gen}(G) \in C_3(\tilde{X}_n/StN_n)$ and no blowdown of G has core one of the θ_{ij} , then clearly $\Theta' \circ \partial(\text{gen}(G)) = 0$. So we will be concerned with those G having an edge x such that the core of the blowdown G_x is a θ_{ij} . Given such G and G_x , let v be the vertex of G_x which the edge x is mapped to by the blowdown $G \rightarrow G_x$. Observe that the colored graph G is determined, up to the color of the edge x , by the colored graph G_x , the vertex v , and a partition into two nonempty sets P_1 and P_2 of the half-edges of G_x meeting v . (In case v is the basepoint of G_x , the basepoint of G must also be specified as one of the two vertices of x . This, however, will not be necessary for our purposes.)

Now we fix G_x to be any colored graph with core some θ_{ij} and use the observation in the previous paragraph to determine all possibilities for G .

As one of essentially two possibilities for the vertex v , we first assume that it is either of the two non-basepoint vertices of G_x . Let a be the edge of the subgraph θ_{ij} which joins the basepoint $*$ to v , and let c and d be the edges forming the monochromatic loop of θ_{ij} .

As a first possibility for the partition, assume that $\{a, c, d\} \subseteq P_1$, where we are not distinguishing between edges and half-edges incident with v . Since G is colored, the color of the edge x of G must be different from the two of θ_{ij} , and so x is a nonsingleton edge of G . Let x' be the edge of G_x having the same color as x . Then, since G is colored, both x and x' are incident with the vertex of G not meeting a , c and d . For each of the resulting three possibilities for the core of G , the map Θ' takes the corresponding generator to 0 since the subgraph θ_{ij} occurs in two faces of the boundary with opposite orientations. The same is true, except in one case, for the other possibilities for G . We will not point this out in the rest of the proof for each G , but will comment on the exceptional case.

As another possibility for the partition, assume $\{a\} \subseteq P_1$, $\{c, d\} \subseteq P_2$. If G is prismatic, the color of x must be the color of a . If G is cubical, with edges x and x' of the same




FIGURE 26. Two views of a generator of $H_2(\tilde{X}_n/StN_n)$.

color, then, in order that G be colored, one vertex of x' must be incident with the non-basepoint vertex of a . Of the two possibilities for the other vertex of x' , one gives a colored graph with x and x' forming a loop, the other the colored graph shown in the middle of Figure 6 in §3.

The latter possibility is the exceptional one in that a θ_{ij} occurs in four faces of the cell corresponding to G . Two of these faces are identical, but have opposite orientations, so cancel in the boundary. Of the other two, one contains θ_{ik} , the other θ_{jk} (for suitable distinct i, j and k). These have opposite orientations, and so Θ' applied to their sum is 0.

As the last possibility for the partition, assume $\{c\} \subseteq P_1$, $\{a, d\} \subseteq P_2$, although we could just as well assume that $\{d\} \subseteq P_1$, $\{a, c\} \subseteq P_2$. If G is prismatic, the color of x must be that of c and the core of G is as shown in Figure 6(a). If G is cubical with x and x' of the same color, then G is only colored if x and x' form a loop in G .

Now assume that v is the basepoint of G_x . Let b be the fourth edge of θ_{ij} different from the edges a, c and d specified above. Assume $\{a, b\} \subseteq P_1$. Then G must be cubical, and the edges x and x' of the same color of G must both be incident with the vertex of G disjoint from a and b .

Finally assume $\{a\} \subseteq P_1$, $\{b\} \subseteq P_2$. If G is prismatic, then x has the color of a and b , giving a case already covered. If G is cubical, the only possibility for the core of G is the partitioned graph C6 as shown in Figure 1 of §3. $\square$

Lemma 6.3. *The homomorphism $\Theta: H_2(\tilde{X}_n/StN_n) \rightarrow \mathbb{Z}^n$ induced by Θ' is onto.*

Proof. In §10 we will exhibit elements of $H_2(\tilde{X}_n)$ whose images in $H_2(\tilde{X}_n/StN_n)$ are mapped to the generators of $\mathbb{Z}^n$ by Θ . Alternatively, the left portion of Figure 26 indicates a cycle in $\tilde{X}_n/StN_n$ which Θ takes to the generator e_k of $\mathbb{Z}^n$. $\square$

This last cycle, which we denote by t , is carried by the image in $\tilde{X}_n/StN_n$ of a genus two surface. The map of this surface to $\tilde{X}_n/StN_n$ is indicated on the right of the figure.

Remark 6.4. We note that t may be viewed as an element of the homology of certain triangular subgroups of $\text{Aut}(F_n)$. Specifically, let σ be any partial order of the set $\mathbf{n}$ with $i > k$, $j > k$, $i \neq j$, and let T be the corresponding triangular subgroup T_n^σ as defined in

§2.6. Consider the maps

$$C_2(ET/T) \xrightarrow{i_*} C_2(\tilde{X}_n^{alg}/StN_n) \xleftarrow{\mu_*} C_2(\tilde{X}_n/StN_n)$$

where i_* is induced by the inclusion $ET \hookrightarrow \tilde{X}_n^{alg}$, and where μ_* is induced by the map $\tilde{\mu}_n: \tilde{X} \rightarrow \tilde{X}_n^{alg}$ defined in §2.10. Choose $t^t \in C_2(ET/T)$ with $i_*(t^t) = \mu_*(t)$. We will see in Theorem 12.3 that t^t represents a nontrivial element of $H_2(T)$ which is closely related to exotic elements of $K_3(\mathbb{Z})$.

A square (resp. triangular or simplicial) colored graph is a colored graph which corresponds to a 1,1-cell (resp. 0,2-cell). Such cells were discussed near the beginning of §3. Generators of $C_2(\tilde{X}_n/StN_n)$ and $C_2(\tilde{X}_n)$ are called square or triangular or simplicial if the corresponding colored graphs are.

The following lemma is used to show that Θ is injective and will also be used in the proof of Lemma 9.1. We say that a chain c in $C_2(\tilde{X}_n/StN_n)$ or $C_2(\tilde{X}_n)$ is reduced if, for each square term of c , all the non-basepoint vertices of the corresponding colored graph have valence three.

Lemma 6.5. *Let G be any square colored graph with $gen(G) \in C_2(\tilde{X}_n/StN_n)$. Then there is a reduced chain c in $C_2(\tilde{X}_n/StN_n)$ such that $\partial c = \partial(gen(G))$ and $c - gen(G) = 0$ in $H_2(\tilde{X}_n/StN_n)$. If G is marked as well as colored with $gen(G) \in C_2(\tilde{X}_n)$, then the same conclusion holds if the space $\tilde{X}_n/StN_n$ is replaced with $\tilde{X}_n$.*

Proof. Disregarding basepoints, there are only two possibilities for the core of G . Either it consists of two monochromatic loops sharing one vertex or of a single monochromatic loop with the two other edges meeting all three vertices of G . Considering basepoints, there are essentially four cases to consider, each of which is straightforward. The key point is that prismatic cells can be used to decrease the valence of selected vertices of colored graphs corresponding to square cells, at the expense of introducing triangular cells. $\square$

A similar lemma, but applicable to triangular rather than square graphs is given next. It will be cited in the last lemma of this section and is also used in the proof of Lemma 9.1.

We will use the following notation. Let G be a simplicial colored graph. Assume that the monochromatic edge loop of G of length > 1 is $e_0, e_1, \dots, e_k$ where $init(e_0) = term(e_k) = *$, the basepoint of G . Let A_i be the set of half edges of G , other than e_{i-1}^+ and e_i^- , which are incident with the vertex $term(e_{i-1}) = init(e_i)$. Then we write $[A_1|A_2|\dots|A_k]_a$ for the oriented simplicial cell $[e_0, e_1, \dots, e_k]$ of $\tilde{X}_n/StN_n$, where a is the color of the loop $e_0, \dots, e_k$. Often the subscript a is dropped.

If G is also marked, then the corresponding oriented cell $[e_0, e_1, \dots, e_k]$ of $\tilde{X}_n$ is also denoted by $[A_1|A_2|\dots|A_k]_a$. This notation makes no mention of the marking, which is usually clear from context.

With this bar notation, the boundary maps for both the space $\tilde{X}_n$ and $\tilde{X}_n/StN_n$ are

$$d_0[A_1|A_2|\dots|A_k] = [A_2|\dots|A_k], \quad d_k[A_1|A_2|\dots|A_k] = (-1)^k[A_1|\dots|A_{k-1}],$$

and, if $0 < i < k$,

$$d_i[A_1|A_2|\dots|A_k] = (-1)^i[A_1|\dots|A_{i-1}|A_i \cup A_{i+1}|A_{i+2}|\dots|A_k].$$

By the cardinality of the cell $[A_1|A_2|\dots|A_k]$, we mean the maximum cardinality of the sets A_i .

Lemma 6.6. *If c is a simplicial 2-chain in $H_2(\tilde{X}_n/StN_n)$ or $H_2(\tilde{X}_n)$ with each 1-cell in the boundary of c having cardinality one, then c is homologous rel boundary (in the sense of Lemma 6.5) to a 2-chain whose terms all correspond to simplicial cells of cardinality two.*

Geometrically, this says that c is homologous to a chain consisting of pairs of simplicial 2-cells meeting along 1-cells of cardinality 2. Each such pair forms a square corresponding to one of the relations R1.

Proof of 6.6. Among all the edges of the 2-cells of c , let e be one of maximum cardinality. Orient e and assume that $[A|B]$ and $[A'|B']$ are 2-cells corresponding to terms of c with $d_1[A|B] = e = d_1[A'|B']$. Thus $[A \cup B] = [A' \cup B'] = e$. We will show that, if the cardinality of e is > 2 , then the square consisting of $-[A|B]$ and $[A'|B']$ is homologous to a simplicial 2-cycle whose cells all have cardinality less than that of e . This will complete the proof.

As a first case, assume that $A' = A - a$, where $a \subsetneq A$, and $B' = B \cup a$. Then the 3-cell $[A'|a|B]$ provides the desired homology since

$$d[A'|a|B] = [a|B] - [A|B] + [A'|B'] - [A'|a]$$

and the first and last terms have cardinality less than that of e .

For another case, assume that $A' = A \cup b - a$ and $B' = B \cup a - b$, where $a \subset A$ and $b \subset B$ both consist of one element. Since the set $A \cup B$ has cardinality > 2 , one of the containments is proper, say $a \subsetneq A$. Then the boundary of the 3-cycle $[A - a|a|B] - [A - a|b|B']$ is $-[A|B] + [A'|B']$ plus four terms corresponding to cells with cardinality less than that of e .

For the last case, assume that $A' = A \cup b - a$ and $B' = B \cup a - b$ as above, where now $a \subset A$ and $b \subset B$ are nonempty subsets with $a \cup b$ containing at least three elements. Say $a = a_1 \sqcup a_2$, where both of the a_i are nonempty. Then the 3-chain

$$[A - a_1|a_1|B] - [A - a_1|b|a_1 \cup B - b] + [A'|a_2|B - b \cup a_1]$$

provides a homology between $-[A|B] + [A'|B']$ and six 2-cells, each with cardinality less than that of e . $\square$

To show that Θ is injective we work with a filtration

$$\ker \Theta' \cap Z_2(\tilde{X}_n/StN_n) \supset \mathcal{F}_R \supset \mathcal{F}_\theta \supset \mathcal{F}_F \supset \mathcal{F}_{Sq} \supset 0$$

by subgroups of $\ker \Theta' \cap Z_2(\tilde{X}_n/StN_n)$, where $\mathcal{F}_R$ consists of the reduced cycles in $\ker \Theta'$. Other terms in the filtration will be defined as we go.

It follows from Lemma 6.5 that all the chains in $\ker \Theta' \cap Z_2(\tilde{X}_n/StN_n)$ are homologous to reduced ones. The remaining lemmas in this section will show that the elements of each subgroup of the filtration are homologous to ones of the next smaller subgroup of the filtration. This will show that Θ is injective.

The subgroup $\mathcal{F}_\theta$ is defined as all the elements of $\mathcal{F}_R$ with no term corresponding to a colored graph with core a θ_{ij} .

Lemma 6.7. *Each element of $\mathcal{F}_R$ is homologous to one of $\mathcal{F}_\theta$.*

Proof. Let $z \in \mathcal{F}_R$. Then, since $\Theta'(z) = 0$, if x is a term of z with $\Theta'(x) \neq 0$, there is another term x' of z with $\Theta'(x + x') = 0$. Note that if $cg(x) = \pm\theta_{ik}$, then (assuming $x \neq -x'$) $cg(x') = \mp\theta_{jk}$ for some $j \neq i$.

Now let g_7 be the generator of $C_2(\tilde{X}_n/StN_n)$ corresponding to the partitioned graph C7 in Figure 1, oriented and colored so that, for a suitable sum p of prismatic terms, the boundary

$\partial(g_7 + p)$ contains terms which cancel $x + x'$ and the cycle $z + \partial(g_7 + p)$ is reduced and has fewer terms on which Θ' is nonzero than z . A cycle homologous to z with no such terms is obtained by, if necessary, adding more boundaries to z like $\partial(g_7 + p)$, but with perhaps different colorings of the corresponding graphs. $\square$

We say that a colored graph is of type F if it is square, reduced, and has a single monocromatic loop. We use the notation $F_2(Si/Tj/k)$, or just $F(Si/Tj/k)$ for type F graphs as in §3 just before Lemma 3.8.

The group $\mathcal{F}_F$ in the filtration is defined as all chains in $\mathcal{F}_\theta$ with no terms corresponding to a colored graph of type F.

Lemma 6.8. *Each element of $\mathcal{F}_\theta$ is homologous to one of $\mathcal{F}_F$.*

Proof. Let D be the free abelian group $\bigoplus_{1 \leq i, j \leq n, i \neq j} (\mathbb{Z}_{ij}^L \oplus \mathbb{Z}_{ij}^R)$ where each $\mathbb{Z}_{ij}^L$ and $\mathbb{Z}_{ij}^R$ is a copy of $\mathbb{Z}$. We let $e_{ij}^L \in \mathbb{Z}_{ij}^L$ and $e_{ij}^R \in \mathbb{Z}_{ij}^R$ be fixed generators. Let $f: C_1(\tilde{X}_n/StN_n) \rightarrow D$ be the homomorphism defined on oriented, reduced generators x by

$$f(x) = \begin{cases} e_{ij}^L, & \text{if } x \text{ corresponds to } L_{ij}, \\ e_{ij}^R, & \text{if } x \text{ corresponds to } R_{ij}^{-1}. \end{cases}$$

On the remaining (unreduced) generators, we define f so that $f(\partial s) = 0$ for all oriented triangular generators s of $C_2(\tilde{X}_n/StN_n)$.

We also define a homomorphism $F: C_2(\tilde{X}_n/StN_n) \rightarrow D$ by $F(x) = f(\partial s)$ for each $x \in C_2(\tilde{X}_n/StN_n)$. It is easy to check that F vanishes on all square reduced generators in $\mathcal{F}_\theta$ except for those of type F, and, for these, $F(F_2(Si/Tj/k)) = \pm e_{ik}^S$.

Now let $z \in \mathcal{F}_\theta$. Assume z contains a term c of type F. Say $c = \pm \text{gen}(F(Si/Tk/j))$. Then, since $F(z) = f(\partial z) = f(0) = 0$, there is another term $c' = \mp \text{gen}(F(Si/Al/j))$ of z such that $F(c') = -F(c)$.

If $k = l$, then there is a single prismatic term p and a generator g_6 of $C_3(\tilde{X}_n/StN_n)$ corresponding to the partitioned graph C6 of Figure 1 suitably colored such that $z + \partial(g_6 + p)$ is in $\mathcal{F}_\theta$ and contains two fewer type F terms than does z . Adding $\partial(g_6 + p)$ to z essentially trades c and c' for two triangular terms and one square one, not of type F.

If $k \neq l$, then there is a sum p of prismatic terms such that $z + \partial(F_3(Si/Tk/Al/j) + p)$ is in $\mathcal{F}_\theta$ and has two fewer type F terms than z . This time c and c' are exchanged for a number of triangular and square terms, where again the square terms are not of type F. $\square$

Let $\mathcal{F}_{S_q}$ be the elements of $\mathcal{F}_F$ (or of $\mathcal{F}_R$) with no square terms.

Lemma 6.9. *Each element of $\mathcal{F}_F$ is homologous to one of $\mathcal{F}_{S_q}$.*

Proof. There are two types of square colored graphs which terms of elements of $\mathcal{F}_F$ may correspond to. Both have two monocromatic loops, and one type has rank three, the other rank four. Generators corresponding to both types have boundary zero.

Given a generator x corresponding to either of these types of colored graphs, there is a chain c consisting of one cubical and one prismatic term with boundary x . The cubical term corresponds to the partitioned graph C2(a) of Figure 1, if $cg(x)$ has rank 4, and to C2(b), if it has rank 3. The prismatic term essentially isolates x . $\square$

Remark 6.10. The boundary of a lift to $\tilde{X}_n$ of one of the chains c in the previous proof in the rank 4 case can be represented by a picture like that on the left of Figure 39, but with

the label of the bottom circle changed from R_{ij} to R_{kl} . All the generators corresponding to the crossings in the picture cancel in $C_2(\tilde{X}_n/StN_n)$ except for the one just above the middle of the picture. This crossing can be taken to correspond to x .

The proof of the Theorem 6.1 is completed by the following.

Lemma 6.11. *Each element of $\mathcal{F}_{S_q}$ is homologous to $0 \in C_2(\tilde{X}_n/StN_n)$.*

Proof. Let $z \in \mathcal{F}_{S_q}$. Note that the terms of z are all triangular. By adding canceling pairs of more triangular terms to z , we can write z as a sum of chains whose boundaries all have cardinality 1. So by Lemma 6.6, we may assume that all the terms of z have cardinality 2. These can be paired off into cycles of two types—type R2 and type R3, where both colored graphs corresponding to the terms of a type Rn cycle have rank n.

Let y be a cycle of type R2. Then there is a 3-chain $c = p_1 + p_2 + S$ with $\partial c = y$. Both of the p_i are prismatic terms and S is a sum of three simplicial terms. One p_i corresponds to the (suitably colored) partitioned graph shown in Figure 39. The other corresponds to the same colored graph, but with the orientation of the edgeset j reversed. Cell orientations are chosen so that contracting the edge j_0 in both graphs gives canceling terms in ∂c . The three simplicial terms effectively decrease the cardinality of two of the triangular terms in $\partial(p_1 + p_2)$ from 3 to 2, thereby isolating y in ∂c .

If y has type R3, then there is a 3-chain c' with $\partial c' = y$. The chain c' is like c , but the singleton edge of color i in each of the five colored graphs corresponding to the terms of c is replaced with two singleton edges. Each of these meets the basepoint of the relevant graph and one of the vertices which i meets. $\square$


As in Remark 6.10, the picture in figure 39 represents a lift of a cycle of type R2 to $\tilde{X}_n$. The crossing just above the center of the picture is the only one that does not cancel in $C_2(\tilde{X}_n/StN_n)$.

7. THE TREE COMPLEX $T(G)$

Fix a graph G (equipped with neither a partition nor a basepoint). After some preliminary definitions, we define the 3-skeleton of a CW-complex $T(G)$. The 2-skeleton of $T(G)$ was implicitly defined in [4], where it was shown that $T(G)$ is 1-connected. We show that $T(G)$ is 2-connected and use related facts in the proof of Lemma 11.5, which in turn is used in the proof of Theorem 11.7.

A graph G' is said to be a quotient of G if it is obtained from G by collapsing each component of an acyclic subgraph of G to a point. If G' is a quotient of G , with corresponding quotient map $q: G \rightarrow G'$, we take $E(G') = \{e \in E(G) : q(e) \text{ is not a point}\}$, with q inducing the identity on $E(G) \cap E(G')$. Two quotients G' and G'' of G are equal if and only if $E(G') = E(G'')$. Thus collapsing distinct acyclic subgraphs (with no components consisting of just a vertex) gives, by definition, distinct quotients, even if these are homeomorphic.

The partitioned quotients of G form a poset $\mathcal{PQ}(G)$, where $G' \leq G''$ if G' is a blowdown of G'' (in which case the partition of G' is obtained by deleting the edges from the edgesets of G'' which are collapsed in the blowdown.) Maximal elements of $\mathcal{PQ}(G)$ correspond to the various partitions of G . If $G' \in \mathcal{PQ}(G)$, let $\mathcal{PQ}_{\leq G'}$ be the full subposet of $\mathcal{PQ}(G)$ consisting of all $H \in \mathcal{PQ}(G)$ with $H \leq G'$. The analog of lemma 2.12 with $\mathcal{PQ}_{\leq G'}$ in place of bG follows immediately from the definition of $\mathcal{PQ}(G)$. So the analog of Prop. 2.14 also holds. That is, the geometric realization $|\mathcal{PQ}(G)|$ of $\mathcal{PQ}(G)$ has a CW-structure with one k -cell

FIGURE 27. Cells of $T(G)$ added to $|\mathcal{PQ}(G)|$.

for each partitioned quotient of G with $k + 1$ vertices. Specifically, if G' is such a partitioned quotient, the geometric realization $|\mathcal{PQ}_{\leq G'}|$ of $\mathcal{PQ}_{\leq G'}$ forms a k -cell of $|\mathcal{PQ}(G)|$.

Note that the 0-cells of $|\mathcal{PQ}(G)|$ correspond to the maximal trees of G . Let T_i be the maximal tree corresponding to the 0-cell v_i . A 1-cell of $|\mathcal{PQ}(G)|$ joins two 0-cells v_i and v_j if the number of edges in $T_i \cap T_j$ is one less than the number of edges in any maximal tree of G . Thus the 1-cells of $|\mathcal{PQ}(G)|$ correspond to elementary moves as defined on p. 225 of [4].

7.1. The definition of $T(G)$. The 2-skeleton of $T(G)$ is defined by attaching a class of 2-cells, called w -triangles, to $|\mathcal{PQ}(G)|$. (In the language of pseudo-isotopy theory, these w -triangles correspond to dovetail singularities. The 2-cells of $|\mathcal{PQ}(G)|$ correspond to the other codimension two singularities.) For each quotient G' of G with two vertices, there is one w -triangle corresponding to each subgraph of G' of the sort shown at the top of figure 27(i). In other words, there is one w -triangle corresponding to each triple $\{T_1, T_2, T_3\}$ of distinct maximal trees of G with $T_1 \cap T_2 \cap T_3$ consisting of one fewer edge than T_1 . The lower part of figure 27(i) indicates the attaching map of the w -triangle corresponding to the subgraph above it. The edge labeled $a \wedge b$, for example, maps to the 1-cell of $|\mathcal{PQ}(G)|$ corresponding to G' with the partition $\{a, b\}$. The map is such that the vertex on the left maps to the 0-cell corresponding to the maximal tree of G which contains a .

To complete the definition of $T(G)$, three types of 3-cells, called w -prisms, w -tetrahedra and pillows, are attached to the 2-skeleton of $T(G)$. For each subgraph of the type shown in figure 27(ii) of each quotient of G with two vertices, or, equivalently, for each quadruple $\{T_1, T_2, T_3, T_4\}$ of distinct maximal trees of G with $T_1 \cap \dots \cap T_4$ consisting of one fewer edge than T_1 , there is one w -tetrahedron attached as indicated in figure 27(ii). Specifically, the four faces of the tetrahedron in (ii) are attached to the w -triangles corresponding to the four subgraphs consisting of three edges of the graph in (ii).

Now let G' be any quotient of G with three vertices. There is one w -prism as shown in figure 27(iii) corresponding to each of the two subgraphs of G' of the types shown at the top of (iii). The two triangular faces of the w -prism in (iii) attach to w -triangles. The three rectangular faces correspond to partitions of the graph in (iii), and so attach to cells of $|\mathcal{PQ}(G)|$. There are two w -prisms corresponding to each subgraph of G' of the type shown in figure 27(iv). One of these two w -prisms is as in (iv), the other is as in (iii). There is one pillow as shown in (v) corresponding to each subgraph of G' of the type shown in (v). Two

of the faces of this pillow attach to w -triangles. The other three faces attach to the 2-cells of $|\mathcal{PQ}(G)|$ corresponding to the graph in (v) with the three partitions $\{a, c, d\}$, $\{b, c, d\}$ and $\{a, b\}, \{c, d\}$. This completes the definition of $T(G)$.

As examples, let $G_1, \dots, G_6$ be the graphs from left to right in Figure 27. Then for $i = 1, 2, 4$ and 6 , the complex $T(G_i)$ is the one just below G_i in the figure. $T(G_3)$ consists of the w -prism in (iii), with one w -tetrahedron attached at each end, and with three pillows then attached to this.

It is not difficult to see that $T(G)$ is connected. This is the content of lemma 1.3 of [4]. Lemma 1.5 of the same paper implies that $T(G)$ is simply connected. For relations (1) and (3) of that lemma correspond to 2-cells of $T(G)$ associated with partitioned graphs, and relation (2) to w -triangles.

7.2. Simplification of cell-to-cell maps. A map of CW-complexes is said to be cell-to-cell if each closed cell of the domain is mapped homeomorphically to a closed cell of the range. The CW-complex with underlying space the 2-sphere S^2 and CW-structure $\mathcal{C}$ is denoted by $(S^2, \mathcal{C})$.

Let $f: (S^2, \mathcal{C}) \rightarrow T(G)$ be a cell-to-cell map. The following four types of operations will be used in the proof of lemma 7.1 below to progressively simplify f .

- (a) A deformation of f to $f': (S^2, \mathcal{C}') \rightarrow T(G)$ which is supported on the interior of a closed disk D equal to the union of 2-cells of $\mathcal{C}$ and such that $f(D) \cup f'(D)$ is the boundary of a 3-cell or chain τ of $T(G)$. Such deformations arise below when the image of D is deformed, rel its boundary, through τ . By subdividing D , we may assume that f' is cell-to-cell.

Assume that the 2-cells σ and τ of $\mathcal{C}$ share at least an edge and that f maps σ and τ to the same 2-cell of $T(G)$. As a subspace of S^2 , $\sigma \cup \tau$ is either a disk, an annulus, or all of S^2 (since the boundary of each 2-cell of $T(G)$ consists of at most four 1-cells).

- (b) If $\sigma \cup \tau$ is a disk D , a deformation of f to $f': (S^2, \mathcal{C}') \rightarrow T(G)$ which is supported on the interior of D and which has the effect that $f'(D) = f(\partial D)$. By increasing the support of the deformation slightly to a neighborhood of D , we may assume that f' is cell-to-cell, where $\mathcal{C}'$ is obtained from $\mathcal{C}$ by deleting the interior of D , and then identifying points in ∂D with the same image under f .
- (c) In case $\sigma \cup \tau$ is an annulus A , delete the interior of A from S^2 and identify the points of the remaining edges of σ and τ which have the same image under f . The resulting quotient space Q consists of two 2-spheres, and f induces a cell-to-cell map $f/\sim$ from Q to $T(G)$. The operation we will use amounts to replacing f with the two cell-to-cell maps defined by restricting $f/\sim$ to each of these 2-spheres.
- (d) If $\sigma \cup \tau$ is all of S^2 , any deformation of f to a constant map.

Lemma 7.1. *Let $f: (S^2, \mathcal{C}) \rightarrow T(G)$ be a cell-to-cell map. Then some sequence of the above four operations transforms f to a constant map or, if operation (c) is used, to several constant maps.*

It follows that $T(G)$ is 2-connected, cf. lemma 11.2, but we will not use this fact.

Before proving 7.1, some terms and lemmas are given. For each closed edge e_i of $T(G)$, pick a point m_i , called the midpoint of e_i , in the interior of e_i . The components of $e_i - m_i$ are called half-edges. If v is a vertex of $T(G)$, let $T(v)$ be the maximal tree of G associated with v . Let e be an edge of $T(G)$ with vertices a and b . If e_a is the halfedge of e containing




FIGURE 28.

a , let $E_G(e_a)$ be the edge $T(a) - T(b)$ of the graph G . This defines a function E_G , called the edge function, from the halfedges of the complex $T(G)$ to the edges of the graph G . A halfedge h with $E_G(h) = \beta$ is called a β -halfedge of $T(G)$.

Now let h be any halfedge of $T(G)$. Assume that h is contained in the 2-cell σ of $T(G)$, and that $E_G(h) = \beta$. Then, it follows from the definition of $T(G)$ that σ contains exactly one other β -halfedge, h' say. The two β -halfedges h and h' meet in a vertex of σ in case σ is a w -triangle. For the other 2-cells σ of $T(G)$, they are disjoint, and there is a unique edge of σ which meets both h and h' . This unique edge is said to be β -parallel for σ . An edge of $T(G)$ is said to be β -parallel if it is β -parallel for at least one of the 2-cells of $T(G)$ containing it.

Assume now that $f: (S^2, \mathcal{C}) \rightarrow T(G)$ is a cell-to-cell map. Use f to pullback the definitions of midpoints, halfedges, the edge function $E(G)$, β -halfedges and β -parallel edges to the CW-structure $\mathcal{C}$ on S^2 .

For each of the following three lemmas, which will be used in the proof of lemma 7.1, we assume that σ and τ are 2-cells of $\mathcal{C}$.

Lemma 7.2. *If $\sigma \cap \tau$ contains an edge which is β -parallel for σ but not τ , then there is a deformation (consisting of type (a) operations above) of f to a cell-to-cell map $f': (S^2, \mathcal{C}') \rightarrow T(G)$ such that*

- (1) *the CW-structures $\mathcal{C}$ and $\mathcal{C}'$ agree except on the interior of the disk $\sigma \cup \tau$, as do the functions f and f' ,*
- (2) *the two edge functions associated with the two CW-structures $\mathcal{C}$ and $\mathcal{C}'$ have the same image, and*
- (3) *the β -parallel edges of $\mathcal{C}'$ in the closed disk $\sigma \cup \tau$ are the two or three edges of τ different from $\sigma \cap \tau$.*

Proof. Figure 28 shows all possibilities for σ and τ , assuming these are as in the statement of the lemma. The six cases below treat these six possibilities, with case (j) treating the configuration shown in part (j) of the figure. In each case, G' denotes the graph obtained from G by blowing down the edges common to all the maximal trees of G associated with the vertices of $T(G)$ in the closed disk $f(\sigma \cup \tau)$.

Case (i). First assume that $x = b$. Then we claim that G' is partitioned by $\{\{a, e\}, \{\beta, b, y\}\}$. (Singleton edges are not shown in the notation for partitions.) Indeed, let ω be any cycle in G' . If ω_y , the cycle in G'_y corresponding to ω , contains either a singleton edge of G'_y or $\{a, e\}$, then ω contains either an edge of G' different from those in $\{a, e, \beta, b, y\}$ or ω contains $\{e, a\}$. Otherwise $\omega_y \supset \{\beta, b\}$. Then $\omega_\beta \supset \{b, y\}$ since G'_β is partitioned by $\{\{b, y\}, \{a, e\}\}$. So $\{\beta, b, y\} \subset \omega$, and G' is partitioned as claimed. A deformation of type (a) above through the corresponding 3-cell of $|\mathcal{PQ}(G)|$ completes case (i) provided $x = b$.

Now assume that $x \neq b$. The following two observations will be used.

- (1) No length two cycle of G' , except for possibly $\{b, x\}$, contains one element from two of the three sets $\{a, e\}$, $\{x, y\}$ and $\{\beta, b\}$.
- (2) No length three cycle of G' , except for possibly $\{a, b, x\}$ or $\{e, b, x\}$, contains one element from each of the three sets $\{a, e\}$, $\{x, y\}$ and $\{\beta, b\}$.

To prove (1), assume ω is a cycle of G' of length two consisting of one edge from each of the sets $\{a, e\}$ and $\{x, y\}$. Then ω_β would not be colored in G_β , a contradiction. Similarly, ω does not consist of one element from each of $\{a, e\}$ and $\{\beta, b\}$ since G_y is partitioned by these two sets. If ω consists of one element from each of $\{x, y\}$ and $\{\beta, b\}$ and $\beta \in \omega$, then ω_β is not colored in G'_β , a contradiction. Similarly, $y \notin \omega$. This leaves $\omega = \{b, x\}$ as the only possibility and completes the proof of (1).

For the proof of (2), assume ω consists of one element from each of $\{a, e\}$, $\{x, y\}$ and $\{\beta, b\}$. If $\beta \in \omega$, then ω_β is not colored in G'_β , and if $y \in \omega$, then ω_y is not colored in G'_y . So $\beta \notin \omega$ and $y \notin \omega$ which proves (2).

If G' is partitioned by $\{\{a, e\}, \{x, y\}, \{\beta, b\}\}$, then $f(\sigma)$ and $f(\tau)$ are two faces of the corresponding 3-cell of $|\mathcal{PQ}(G)|$. A deformation, rel the boundary of the disk $f(\sigma) \cup f(\tau)$, across this 3-cell finishes this case.

So assume that G' is not partitioned by $\{\{a, e\}, \{x, y\}, \{\beta, b\}\}$, and let ω be any cycle of G' which is contained in $\{a, e, x, y, \beta, b\}$ and which contains none of the three elements of $\{\{a, e\}, \{x, y\}, \{\beta, b\}\}$. Then the length of ω is either two or three, and so, by (1) and (2), ω equals either $\{b, x\}$, $\{a, b, x\}$ or $\{e, b, x\}$.

First assume that $\omega = \{b, x\}$. If b, x, β and y all meet the same vertex, then (1) implies that the other vertices of b, β and y are distinct. But then of the 4 choose 2 ways that e can be attached to the four vertices of G' , three are ruled out by (1) and the other three by (2). It follows from this contradiction, and the fact that $\{\beta, y\}$ is not a cycle, that x, y and β form either a path or loop of length 3 in G' . If they form a loop, then it follows from (1) that the vertex of G' not in this loop meets both e and a . And if they form a path and not a loop, then it follows from (1) and (2) that both e and a meet both endpoints of this path. Either way, G' is partitioned by $\{\{e, a\}, \{x, y, \beta\}\}$ and case (i) is completed (assuming $\omega = \{b, x\}$) using two deformations of type (a) above through the two 3-cells of $T(G)$ shown in figure 29(a).

Now assume $\omega = \{e, b, x\}$ (with a similar argument if $\omega = \{a, b, x\}$). By (1) and (2), both of the edges y and β meet the vertex of G' which is not contained in the cycle ω . Also y and β do not form a cycle of length two. We consider three cases depending on which of the three vertices of ω the edge β meets.

First assume β meets $x \cap e$. Then y meets $b \cap x$, because if y met $b \cap e$ then $\{y, \beta, e\}$ would be a cycle of length three, contradicting (2). Now, by (1), either a meets both vertices of e , in which case a deformation as indicated in figure 29(a) completes the proof, or a meets




FIGURE 29.

$y \cap \beta$ and $b \cap e$. But the latter case cannot occur since it would result in the cycle $\{a, b, y\}$ in contradiction to (2).

Now assume β meets $b \cap e$. Then, by (2), y meets $x \cap b$. So, by (1), either a meets both vertices of e , or a meets $\beta \cap y$ and $e \cap x$. In the first case, use the deformation indicated by figure 29(a), and in the second, use the one indicated by figure 29(b).

Finally assume that β meets $b \cap x$. Then, using (1) and (2), either y meets $b \cap e$ and a meets both vertices of e , or y meets $e \cap x$ in which case a meets both vertices of e or a meets $b \cap e$ and $\beta \cap y$. For all three possibilities, a deformation like that indicated by figure 29(a) can be used to complete the proof.

Case (ii). Assume first that $x \neq b$. In case $\{b, \beta\}$ is a loop in G' , then, since G' is partitioned by $\{\{a, e\}, \{\beta, b\}\}$, the edges e and a both meet the vertex of G' not in the loop $\{b, \beta\}$. So does the edge x , since τ is a w -triangle corresponding to the graph G'_β . Thus there is a w -prism having σ and τ as faces. This 3-cell of $T(G)$ can be used to complete the proof (assuming $\{b, \beta\}$ is a loop).

So assume $\{b, \beta\}$ is not a loop in G' . Then $\{e, a\}$ is. If x meets both vertices of e , then, as just above, a single w -prism can be used to complete the proof. If x meets the vertex $\beta \cap b$, and so also $e \cap b$, then a deformation as indicated in figure 29(c) completes the proof.

Now assume that $x = b$. Then $\{b, \beta\}$ is not a loop in G' , so $\{e, a\}$ is, and the vertex $b \cap \beta$ is not contained in this loop. Therefore there is a pillow in $T(G)$ having σ and τ as faces which can be used to complete the proof.

Case (iii). We will use the following.

- (1) No cycle of length three in G' , except possibly for $\{a, e, b\}$, contains two edges from $\{a, e, x\}$ and one from $\{\beta, b\}$.
- (2) The only possible cycle of length two in G' made up of edges from the set $\{a, e, x, \beta, b\}$ is $\{\beta, b\}$.

To prove (1), assume ω is a cycle of length three with two edges in $\{a, e, x\}$ and one in $\{\beta, b\}$. If $\beta \in \omega$, then ω_β is not colored in G'_β . And if $x \in \omega$, then ω_x is not colored in G'_x . So $\beta \notin \omega$ and $x \notin \omega$.

For (2), let ω be a cycle of length two and assume that $\omega \subset \{a, e, x, \beta, b\}$ and $\omega \neq \{\beta, b\}$. If $\beta \in \omega$, then ω_β is not colored in G'_β . If $b \in \omega$, then ω_x is not colored in G'_x . And if $\{b, \beta\} \cap \omega = \emptyset$, then $\omega \subsetneq \{a, e, x\}$, and so is not colored in G'_β . These are all contradictions, so (2) is proved.

If G' is partitioned by $\{\{a, e, x\}, \{\beta, b\}\}$, then a deformation across the corresponding O-prism of $T(G)$ which contains σ and τ as faces completes the proof. So assume G' is not




FIGURE 30.

partitioned by $\{\{a, e, x\}, \{\beta, b\}\} = \{h_i\}$, where, as usual, singleton h_i are not shown in the set on the left of the equals sign. Let ω be a loop in G' which contains none of the h_i . Then ω has length two or three, and so it follows from (1) and (2) that $\omega = \{a, e, b\}$. Now (1) and (2) imply that both β and x meet the vertex of G' not in ω . If β meets $a \cap e$, then each of the three ways of attaching x contradicts either (1) or (2). So β meets one vertex of b and, by (1) and (2) again, x meets the other. Thus G' is partitioned by $\{a, e, x, \beta\}$, and figure 30(d) shows how to complete the proof.

Case (iv). If G' is partitioned by $\{\{x, y\}, \{a, e, \beta\}\}$, then use the corresponding O-prism of $T(G)$ with σ and τ as faces. If not, then, just as in case (iii), $\{a, e, x\}$ and $\{x, y, \beta\}$ are loops in G' . So G' is partitioned by $\{\{x, y, \beta\}, \{a, e\}\}$ and figure 30(e) indicates how to complete the proof.

Case (v). Since G' is partitioned by $\{a, e, \beta\}$, the edges a, e and β form a loop in G' . Because of the 2-cell τ , the edge x meets $a \cap e$ and one vertex of β . If x meets $\beta \cap a$, then G' is partitioned by $\{\{a, x\}, \{\beta, e\}\}$ and the pillow with σ, τ and the corresponding 2-cell as faces can be used to complete the proof. If x meets $\beta \cap e$, use the pillow with faces σ, τ and the 2-cell corresponding to G' partitioned by $\{\{e, x\}, \{\beta, a\}\}$.

Case (vi). Let ω be a loop in G' consisting of a subset of $\{a, e, x, \beta\}$. Then $\{a, e, \beta\} \subset \omega_x$ since G'_x is partitioned by $\{a, e, \beta\}$. Similarly, $\{a, e, x\} \subset \omega_\beta$. Therefore $\omega = \{a, e, x, \beta\}$ and so G' is partitioned by $\{a, e, x, \beta\}$. Use the corresponding 3-cell of $T(G)$ to complete the proof of this case and the lemma. $\square$

Lemma 7.3. *Assume that $f(\sigma)$ and $f(\tau)$ are distinct 2-cells of $T(G)$. If $\sigma \cap \tau$ consists of a single edge which is β -parallel for both σ and τ , then the same conclusions as in the previous lemma hold, except that (3) of that lemma is replaced by:*

- (3') *The β -halfedges of $\mathcal{C}'$ in the closed disk $\sigma \cup \tau$ are the same as those of $\mathcal{C}$ and are contained in two w -triangles of $\mathcal{C}'$.*

Proof. We consider all possible configurations for σ and τ in the following two cases. Define the graph G' as at the start of the proof of lemma 7.2.

Case (i). Assume σ and τ are as in figure 28(i), but with $y = \beta$. If $\{a, e\}$ is a loop in G' , then the edges β, b , and x all meet the vertex of G' not in the loop $\{a, e\}$. So there is a w -prism with $f(\sigma)$ and $f(\tau)$ as faces. If $\{a, e\}$ is not a loop, then both $\{\beta, x\}$ and $\{\beta, b\}$ are, and so again $f(\sigma)$ and $f(\tau)$ are faces of some w -prism in $T(G)$. Deformations across these w -prisms complete the proof in both cases.

Case (ii). Assume σ and τ are as in figure 28(iv), but with $y = \beta$. Then, since G' is partitioned by $\{a, e, \beta\}$, the edges a , e and β form a loop in G' . So $\{a, e\}$ is not a loop in G' , which implies that $\{x, \beta\}$ is, since G' is also partitioned by $\{\{a, e\}, \{x, \beta\}\}$. Use the pillow with $f(\sigma)$ and $f(\tau)$ as faces to complete the proof of this case and the lemma. $\square$

Let h be any halfedge of $(S^2, \mathcal{C})$. Say $E_G(h) = \beta$. Let $\{\sigma_i\}$ be the set of 2-cells of $\mathcal{C}$ which contain a β -halfedge. For each σ_i , let m_i and m'_i be the midpoints of the two edges of σ_i which contain β -halfedges. Also let a_i be an arc which joins m_i and m'_i and which is otherwise contained in the interior of σ_i . The union of the a_i form a collection of disjoint circles in S^2 , called β -circles.

Lemma 7.4. *Assume c is a β -circle in S^2 contained in $\bigcup_{i=1}^n \sigma_i$ where each σ_i is a closed w -triangle of $\mathcal{C}$ meeting c . Then f can be deformed to either a cell-to-cell or to a constant map f' with the effect that*

- (1) c is eliminated and no new β -circles are introduced and
- (2) if E_G (resp. E'_G) is the edge function associated with f (resp. f'), the image of E'_G is contained in the image of E_G .

Proof. Assume $c \subset \bigcup_{i=1}^n \sigma_i$ as in the statement of the lemma. We say that two of the σ_i are adjacent if they share at least one edge. If f takes two adjacent σ_i to the same cell of $T(G)$, then n can be decreased by two by using a deformation of either type (b) or (d) above. If adjacent σ_i map to different cells, then n can be decreased by at least one with a type (a) deformation across the w -tetrahedron of $T(G)$ having the adjacent σ_i as faces. $\square$

Proof of lemma 7.1. Let β be any element of the image of the edge function associated with the cell-to-cell map $f: (S^2, \mathcal{C}) \rightarrow T(G)$, and let $\{c_i\}$ be the set of β -circles in S^2 . Of the two disks in S^2 which each c_j bounds, let D_j be the one which contains all the β -parallel edges for the 2-simplices meeting c_j . Let D' be one of the D_j . Among all the D_j contained in D' , let D be an innermost one. If none of the c_i are contained in the interior of D , we say that D is strictly innermost.

First assume that D is strictly innermost. Let c be the β -circle which D bounds. Consider the set $\mathcal{B}$ of β -parallel edges which are contained in 2-cells meeting c . We assume, for now, that $\mathcal{B} \neq \emptyset$. If some $e \in \mathcal{B}$ is contained in the intersection, $\sigma \cap \tau$ say, of two 2-cells of $\mathcal{C}$ and e is β -parallel for σ but not τ , then an application of lemma 7.2 decreases by one the total number of 2-cells contained in D . Thus, since D is strictly innermost, we may assume, by repeated applications of lemma 7.2, that, for each $e \in \mathcal{B}$, both 2-cells containing e meet the β -circle c .

So now, if $e \in \mathcal{B}$, either lemma 7.3 or one of the operations (b), (c) or (d) above can be applied to the 2-cells, σ and τ say, containing e . Applying the appropriate one decreases the cardinality of $\mathcal{B}$ by one by eliminating e from it. If lemma 7.3 is applied, the β -circle c is surgered into two β -circles and the portion of these in the disk $\sigma \cup \tau$ is contained in two w -triangles. If (b) or (c) is applied, then c is again transformed into two β -circles. If (c) has been applied, these β -circles are in two different 2-spheres. (The edges of the CW -structures of these 2-spheres are identified with those of $\mathcal{C}$ in the natural way.)

By repeating the above procedure, once for each of the remaining edges in $\mathcal{B}$, we may assume that $\mathcal{B} = \emptyset$. Then either c has been eliminated or c has been transformed into some number of β -circles. In the latter case, each of the β -circles is contained in a union of w -cells and so can be eliminated by lemma 7.4. This shows that if D is strictly innermost, then

the number of β -circles can be decreased by one, and the image of the edge function is, if changed, made smaller. (In case operation (c) is used, then f' has been transformed into some number of cell-to-cell maps $f_i: S^2 \rightarrow T(G)$. Then “number of β -circles” means “the sum of the number of β -circles in the domains of the f_i ” and “image of the edge function” means “the union of the images of the edge functions of the f_i ”.)

Assume now that D is innermost, but not strictly so, and let R denote the component of $S^2 - \bigcup c_i$ in D having c as limit points. Define the nonempty set $\mathcal{B}$ as above and, as above, use lemma 7.2 to decrease the number of 2-cells in R until some edge of $\mathcal{B}$ is contained in a 2-cell which meets a β -circle different from c . Then apply either operation (b) or lemma 7.3 to the two 2-cells containing this edge. This process joins the two β -circles into one and so again decreases the number of β -circles by one without increasing the image of the edge function.

Thus all β -circles can be eliminated in such a way that the image of the edge function becomes a proper subset of what it was originally. This completes the proof since β was an arbitrary element of the image of the edge function. $\square$

8. THE SIMPLICIAL COMPLEX $\mathcal{T}$

Let $\mathcal{S}_T$ denote the sphere complex SA_n defined in §2.4 of [5] together with, for each vertex (G, ρ) of SA_n , some choice of a maximal tree in G . In more detail, each vertex of the sphere complex SA_n , which we denote by $\mathcal{S}_n$, or just $\mathcal{S}$, is an equivalence class of pointed marked graphs of rank n . Usually the adjective pointed is dropped in this context. A maximal tree of any one representative of an equivalence class serves to specify a maximal tree in every representative. Such a compatible choice of maximal trees is what is meant by a choice of a maximal tree in G .

We point out that there is a simplicial map $\mathcal{F}: \tilde{X}_n \rightarrow \mathcal{S}_n$ which forgets partitions and colorings. It will be used in §11.

If $v = (G, \rho)$ is a vertex of $\mathcal{S}_T$ which is not a rose, let $p(v)$ denote the edge of $\mathcal{S}$ which joins v to the marked rose G/T , where T is the chosen maximal tree in G . With the edges of $\mathcal{S}$ oriented so that each points towards the vertex corresponding to the graph with fewer vertices, we write *term* $p(v)$ for the terminal vertex of $p(v)$. If v is a rose, then $p(v)$ is, by definition, just the vertex v , and we say, for convenience, that $p(v)$ terminates at v and set *term* $p(v) = v$. Let $\mathcal{C}$ be the subcomplex of $\mathcal{S}$ consisting of those simplices $\{v_0, \dots, v_n\}$ with *term* $p(v_0) = \dots = \text{term } p(v_n)$.

Lemma 8.1. *Each component of $\mathcal{C}$ is contractible and contains exactly one marked rose.*

Proof. Let C_i be a component of $\mathcal{C}$, and let $\{v_j\}$ be the set of vertices of C_i . Since C_i is connected all the $p(v_j)$ terminate at the same marked rose, r say. Therefore C_i is a cone on r and so is contractible. $\square$

Let $\mathcal{T}$ be the simplicial complex obtained from $\mathcal{S}$ by contracting the components of $\mathcal{C}$ to distinct points. Since $\mathcal{C}$ contains all the vertices of $\mathcal{S}$, $\mathcal{T}$ has one vertex for each component C_i of $\mathcal{C}$, and $\{C_0, \dots, C_n\}$ is a simplex of $\mathcal{T}$ if and only if there is a simplex $\{v_0, \dots, v_n\}$ of $\mathcal{S}$ with $v_i \in C_i$ for $i = 0, \dots, n$. Equivalently, the vertices of $\mathcal{T}$ correspond to the roses of $\mathcal{S}$, and $\{R_0, \dots, R_n\}$ is a simplex of $\mathcal{T}$ if and only if there is a simplex $\{v_0, \dots, v_n\}$ of $\mathcal{S}$ with *term* $p(v_i) = R_i$ for $i = 0, \dots, n$. The simplicial quotient map $q: \mathcal{S} \rightarrow \mathcal{T}$, a homotopy equivalence by lemma 8.1, can be defined by taking each vertex v of $\mathcal{S}$ to the vertex *term* $p(v)$ of $\mathcal{T}$. A simplex $\tilde{\sigma}$ of $\mathcal{S}$ is called a lift of the simplex σ of $\mathcal{T}$ if $q(\tilde{\sigma}) = \sigma$ and if the dimensions

of σ and $\tilde{\sigma}$ are the same. Prop. 8.2 below shows that among all the graphs corresponding to the lifts of a given simplex, the one with the fewest number of vertices is unique. Moreover, any such graph blows down to this unique lift.

Let $\sigma = \{R_0, \dots, R_n\}$ be an n -simplex of $\mathcal{T}$, and let $\tilde{\sigma} = G_n \rightarrow G_{n-1} \rightarrow \dots \rightarrow G_0$ be a lift of σ to $\mathcal{S}$. (So the R_i are marked roses, the G_i are marked graphs with markings compatible with the blowdowns $G_i \rightarrow G_{i-1}$, and we may assume that $q(G_i) = R_i$.) For $i = 0, \dots, n-1$, let T_i be the maximal tree of G_n which is blowdown by the composite $G_n \rightarrow \dots \rightarrow G_i \xrightarrow{q} R_i$. Also let T_n be the maximal tree of G_n associated with the vertex G_n of $\mathcal{S}_T$. Let $G_{\tilde{\sigma}}$ be the marked, pointed graph obtained from G_n by collapsing the edges of $\bigcap_{i=0}^n T_i$.

Proposition 8.2. *The vertex of the sphere complex $\mathcal{S}$ corresponding to the marked graph $G_{\tilde{\sigma}}$ does not depend on the choice of lift $\tilde{\sigma}$ of σ .*

The following definitions and conventions will be used in the proof. Let G be a graph with edges oriented and labeled. If H is any subgraph of G , let $L(H)$ denote the set of labels of the edges of H , and let $L(H)^\pm = L(H) \cup \{e^{-1} : e \in L(H)\}$.

If $x \in L(H)$, then $init(x)$ and $term(x)$ denote the initial and terminal vertices, respectively, of the edge x labels. The same edge as x , but with the opposite orientation, is denoted by x^{-1} . Thus $init(x) = term(x^{-1})$ and $term(x) = init(x^{-1})$. An oriented edgepath in G is specified by a word in the elements of $L(G)^\pm$ in the obvious way.

Assume now that G is marked by $g: R_0 \rightarrow G$ where the edges of R_0 are labeled by elements of the set $\{r_i\}$. We often write $g_E(r_i)$ for the (reduced) word corresponding to the edgeloop $g(r_i)$. Here we have identified a label of an edge with the edge itself. We often do this. If e is an edge of G , let $e \cdot g(r_i)$ be the number of occurrences of e in the (reduced) word $g(r_i)$ plus the number of occurrences of e^{-1} in the same. Let $e \cdot im(g) = \sum_i e \cdot g(r_i)$. Also, if e_i and e_j are edges of G , let $e_i e_j \cdot im g = \sum_k e_i e_j \cdot g(r_k)$ where $e_i e_j \cdot g(r_k)$ is the number of occurrences of the subword $e_i e_j$ in $g(r_k)$ plus the number of occurrences of $e_j^{-1} e_i^{-1}$ in the same.

Proof of 8.2. We first assume that $\sigma = \{R_1, R_2\}$ is a 1-simplex of $\mathcal{T}$ with lifts $\tilde{\sigma} = G_1 \rightarrow G_0$ and $\tilde{\sigma}' = G'_1 \rightarrow G'_0$ in $\mathcal{S}$. Let G denote the marked graph $G_{\tilde{\sigma}}$ and let S and T be the maximal trees of G which are the image under $G_1 \rightarrow G$ of the maximal trees T_0 and T_1 of G_1 . Define G' , S' and T' similarly. Then there is a homotopy commutative diagram

$$\begin{array}{ccccc}
 & & G & & \\
 & \swarrow S & \uparrow f & \searrow T & \\
 R_2 & \xleftarrow{h} & R_0 & \xrightarrow{g} & R_1 \\
 & \swarrow S' & \downarrow f' & \searrow T' & \\
 & & G' & &
 \end{array}$$

where f and f' are the markings of G and G' , and where the arrows labeled S , T , S' and T' are blowdowns of the maximal tree of the label.

By viewing the diagram as four edges in $\mathcal{S}$, and then acting on them with a suitable element of $\text{Aut}(F_n)$, we may assume that h is homotopic to a homeomorphism. We often use this homeomorphism to identify the edges of R_2 with those of R_0 .

We say that an edge e of R_1 is paired with the edge e' of R_2 if $e \cdot g(r_i) = e' \cdot h(r_i)$ for every edge r_i of R_0 .

Lemma 8.3. *An edge e of R_1 is paired with the edge e' of R_2 if and only if $S(b) = e'$ and $T(b) = e$ for some edge b of G .*

Proof. Choose labels for the edges of G and let $B = L(G) - L(S \cup T)$. Label the edges of R_1 and R_2 so that $S(e) = e$ if e is an edge of G not in the tree T , and $T(e) = e$ if e is not in the tree S . Then

$$L(R_2) = L(T) \cup B \quad \text{and} \quad L(R_1) = L(S) \cup B.$$

If $x \in L(R_2) \approx L(R_0)$, then $x \cdot \text{im}(h) = 1$ since h is homotopic to a homeomorphism. If $s \in L(S) \subset L(R_1)$, then $s \cdot \text{im}(g) \geq 2$ by the following lemma. So an edge e of R_1 can be paired with an edge e' of R_2 only if $e \in B \subset L(R_1)$. Since h is homotopic to a homeomorphism, it is easy to see that each $e \in B \subset L(R_1)$ is paired with the unique edge of $B \subset L(R_2)$ with the same label as e .

The implication in the other direction is easy. $\square$

Lemma 8.4. *If $s_1 \dots s_n$ is an edgepath in the tree S terminating at the basepoint of G , then*

$$2 \leq s_1 \cdot \text{im } g < s_2 \cdot \text{im } g < \dots < s_n \cdot \text{im } g.$$

Proof. Label the edges of G and the R_i as in the proof of lemma 8.3. If $t \in L(T)^\pm \subset L(R_2)^\pm \approx L(R_0)^\pm$, then, since h is homotopic to a homeomorphism, $g_E(t)$ is a word in the letters $L(S)^\pm$, and is obtained from $f_E(t)$ by deleting the unique occurrence of t in $f_E(t)$. Also if $b \in B$, then $g_E(b) = f_E(b)$. So the lemma follows since each vertex of G (other than the basepoint) has valence ≥ 3 . $\square$

Lemma 8.5. *If $ss' \cdot \text{im } g > 0$ where $s, s' \in L(S)^\pm$, then*

$$\begin{aligned} ss' \cdot \text{im } g = 1 &\Leftrightarrow \text{there is a } t \in L(T)^\pm \text{ such that } sts' \text{ is an edge path in } G, \text{ and} \\ ss' \cdot \text{im } g \geq 2 &\Leftrightarrow ss' \text{ is an edgepath in } G \text{ where } \text{init}(s') = \text{term}(s) \neq *. \end{aligned}$$

Proof. If, in G , $\text{term}(s) \neq \text{init}(s')$, then both sides of the first of the two equivalences in the statement of the lemma hold. Otherwise, the same is true of the second equivalence by 8.4 and a little more. $\square$

Returning now to the proof of 8.2, label the edges of R_1 and R_2 so that two edges have the same label if and only if they are paired. Then label the edges of G so that an edge e of G has the same label as an edge r of R_i if the blowdown $G \rightarrow R_i$ takes e to r . Such a labeling is possible by lemma 8.3. Next, orient the edges of G and the R_i so that the blowdowns S and T preserve orientations. Label and orient the edges of G' similarly. Let $B = L(R_1) \cap L(R_2)$. Then we have

$$L(G) = L(R_0) \cup L(R_1) = L(B) \sqcup (L(R_1) - B) \sqcup (L(R_2) - B) = L(G').$$

Thus a bijection from the edges of G to those of G' is defined by taking an edge e of G to the edge of G' with the same label as e . We will show that this respects incidence relations and so defines a homeomorphism $G \rightarrow G'$.

As a first step we show that f_E is determined by g_E . Let $t \in L(T)^\pm$. Then $g_E(t) = s_1 \dots s_n$ for some s_i in $L(S)^\pm$. Assume, for now, that $n > 1$. If $s_i s_{i+1} \cdot \text{im } g = 1$ for some consecutive pair of letters s_i and s_{i+1} in $g_E(t)$, then, by 8.5 and the fact that T has no cycles, $f_E(t) = s_1 \dots s_i t s_{i+1} \dots s_n$. Otherwise, $f_E(t)$ equals either $s_1 \dots s_n t$ or $t s_1 \dots s_n$. Which, is determined by g , according to 8.4.

If $n = 1$ so that $g_E(t) = s_1$, then $s_1 \cdot \text{im } g \geq 2$ by 8.4. Thus there is an $e \in L(R_0) = L(R_2)$ such that $s_1 \cdot g(e) > 0$ and $e^{\pm 1} \neq t^{\pm 1}$. Assume $g_E(e) = s_1^{\pm 1}$. If $e \in L(T)^\pm$, then the edges

e and t would form a cycle in T . And if $e \in L(B)$, then $g_E(e)$ would contain the letter e , another contradiction. So the word $g_E(e)$ contains at least two letters, with either the first or last being s_1 or s_1^{-1} . If the first is s_1 or the last is s_1^{-1} , then $f_E(t) = s_1 t$. Otherwise, $f_E(t) = t s_1$. Since $f_E(b) = g_E(b)$ if $b \in L(B)$, we have shown how to compute f_E from g_E . Similar arguments apply just as well to G' and f'_E , so we conclude that $f_E = f'_E$.

Now, for convenience, assume that the edges of the tree S are oriented so that they point toward the basepoint of G . Let $x \in L(G)^\pm = L(G')^\pm$, and let $y \in L(S)$. Then

$$\begin{aligned} * \neq \text{term}(x) = \text{init}(y) \text{ in } G &\Leftrightarrow xy \cdot \text{im } f_E > 0 \\ &\Leftrightarrow xy \cdot \text{im } f'_E > 0 \Leftrightarrow * \neq \text{term}(x) = \text{init}(y) \text{ in } G'. \end{aligned}$$

For edges meeting the basepoint of G and G' , we have

$$\begin{aligned} \text{init}(x) = * \text{ in } G &\Leftrightarrow f_E(z) \text{ begins with } x \text{ for some } z \in L(R_0)^\pm \\ &\Leftrightarrow f'_E(z) \text{ begins with } x \text{ for some } z \in L(R_0)^\pm \Leftrightarrow \text{init}(x) = * \text{ in } G'. \end{aligned}$$

It follows that a homeomorphism $h: G \rightarrow G'$ is defined by identifying edges of G and G' with the same labels in an orientation preserving manner. Since $f_E = f'_E$, we have $h \circ f = f'$. Therefore G and G' are equivalent as marked graphs.

Now let $\sigma = \{R_1, \dots, R_{n+1}\}$ be an n -simplex of $\mathcal{T}$ with lift $\tilde{\sigma} = G_1 \rightarrow \dots \rightarrow G_{n+1}$ in $\mathcal{S}$. Let $G = G_{\tilde{\sigma}}$ be the marked graph with maximal trees T_i as defined above. Also let T_i , for $i = 1, \dots, n+1$, denote the image of T_i under the blowdown $G_0 \rightarrow G_{\tilde{\sigma}}$. Thus the T_i are maximal trees in G with $\bigcap_i T_i$ containing no edges. Each edge $G_i \rightarrow G_j$ of $\tilde{\sigma}$ gives rise to a commutative diagram

$$\begin{array}{ccccc} & & G & & \\ & \swarrow T_i & \downarrow & \searrow T_j & \\ R_i & \longleftarrow G/T_i \cap T_j & \longrightarrow & R_j & \\ & \swarrow g_i & \uparrow f_{ij} & \searrow g_j & \\ & & R_0 & & \end{array}$$

where all the arrows are blowdowns except for g_i , g_j , and f_{ij} which are markings. These markings are induced by the marking $f: R_0 \rightarrow G$ of G which is not shown in the diagram.

Label the edges of G and the R_i as above (in the case where σ was a 1-simplex) so that an edge e of G has the same label as an edge e' of R_i if and only if T_i sends e to e' . Note that whether an edge of R_i has the same label as an edge of R_j can be determined from just g_i and g_j by precomposing g_i with a suitable homotopy equivalence of R_0 and then using 8.3. Also orient the edges of G and the R_i so that the maps T_i preserve the orientations of the edges. We will show that the function f_E from $L(R_0)^\pm$ to the set of words in the elements of $L(G)^\pm$ is determined by the g_i . Since $L(G) = \bigcup_i L(R_i)$, the proof can then be completed as above assuming, as we may, that $g_0: R_0 \rightarrow R_0$ is homotopic to a homeomorphism, and then using the tree T_1 of G in place of the tree S used in the above argument.

Since the edge $G_1 \rightarrow G_i$ of $\tilde{\sigma}$ is a lift of the 1-simplex $\{R_1, R_i\}$ of $\mathcal{T}$, the functions $f_i^E \equiv f_{0i}^E$ (where E has been written as a superscript rather than a subscript) are determined by the g_i . Assume $f_i^E(r) = S_i r S'_i$ where $r \in L(R_0)$, and where the S_i and S'_i are words in the letters $L(T_1 - T_i)^\pm \subset L(T_1)^\pm$. Since $\bigcup_i L(T_1 - T_i) = L(T_1)$, the union of the letters in the words S_i form an edgepath in the tree T_1 of G from $*$ to $\text{init}(r)$. Since $x \cdot \text{im } f = x \cdot \text{im } g_i$ if $x \in L(T_1 - T_i) \subset L(G)$, the order of the letters in this edgepath can be determined using

the g_i and lemma 8.4. A similar argument using the S'_i shows that the g_i determine $f_E(r)$ and so also f_E . $\square$

8.1. The CW-complex $\mathcal{T}^*$. Because of the proposition just proved, we henceforth denote $G_{\bar{\sigma}}$ by just G_{σ} . Note that the rule $\sigma \mapsto G_{\sigma}$ is functorial in the sense that it induces a simplicial map from the barycentric subdivision of $\mathcal{T}$ to $\mathcal{S}$.

If G is a marked, pointed graph, let $T_M(G)$ denote the CW-complex $T(G)$ defined in section 7.1, but with, in addition, the graphs associated with the cells of $T_M(G)$ marked as blowdowns of G . The basepoint of G does not figure into the definition of $T(G)$, but is needed for the definition of $T_M(G)$ since markings necessarily preserve basepoints. The map $T_M(G) \rightarrow T(G)$ which forgets markings is a cellular isomorphism, and T_M becomes a functor if blowdowns $G \rightarrow H$ of marked, pointed graphs are sent to the natural inclusions $T_M(H) \hookrightarrow T_M(G)$. Composition of functors gives the functor $\sigma \mapsto T_M(G_{\sigma})$ from the simplices of $\mathcal{T}$ to CW-complexes.

We next use this last functor to define a cellular subdivision $\mathcal{T}^*$ of the 2-skeleton of $\mathcal{T}$. (Only a subdivision of a finite subcomplex of $\mathcal{T}$ will be needed in applications.) Let e be any edge of $\mathcal{T}$, and let v_1 and v_2 be the vertices of e . Each v_i corresponds to a marked rose which is a blowdown of G_e , and so also corresponds to a vertex v'_i of $T_M(G_e)$. Choose a subdivision e^* of e so that there is a cellular isomorphism which takes v_i to v'_i and which takes e^* onto a shortest edgepath in $T_M(G_e)$ joining v'_1 with v'_2 . For example, if G_e has just two vertices, then $e^* = e$. Subdivide each edge of $\mathcal{T}$ in this way to obtain the 1-skeleton of $\mathcal{T}^*$. Each 1-cell of $\mathcal{T}^*$ corresponds to a marked partitioned graph with two vertices.

Let τ be any 2-simplex of $\mathcal{T}$, and let e_1 , e_2 , and e_3 be the three faces of τ . Define a map $f: \partial\tau \rightarrow T_M(G_{\tau})$ by piecing together the three composites $e_i \rightarrow T_M(G_{e_i}) \rightarrow T_M(G_{\tau})$. The first map in each of these was chosen above, and the second is provided by the functor $\sigma \mapsto T_M(G_{\sigma})$. If $f(\partial\tau)$ is a contractible subspace of $T_M(G_{\tau})$, then τ is called a degenerate simplex and is not subdivided. Otherwise, let τ^* be a cellular subdivision of τ such that f extends to a cell-to-cell map $\tau^* \rightarrow T_M(G_{\tau})$. Subdivide each 2-simplex of $\mathcal{T}$ in this way and let $\mathcal{T}^*$ be the resulting CW-complex. Also fix choices of the above cell-to-cell maps. Then the 2-skeleton of $\mathcal{T}^*$ comes equipped with a map G to the space $\bigsqcup_{\sigma \in \mathcal{T}} T_M(G_{\sigma}) / \sim$ where $x \sim y$ if there is a simplex τ of $\mathcal{T}$ with a face σ such that the inclusion $T_M(G_{\sigma}) \hookrightarrow T_M(G_{\tau})$ takes one of x or y to the other. Degenerate simplices of $\mathcal{T}$ are mapped by G to the image of their boundaries.

8.2. Standard w -crossings. If an element w of the group StN is either w_{ij}^L or w_{ij}^R , as defined in §2.8, then, according to §1.4 of [7], the conjugate $wg^{-1}w^{-1}$ is a defining generator of the group StN , provided g is. The corresponding relation in the group StN can be realized (as consequences of defining relations for StN) in $\tilde{X}$ as the union of 2-cells. We will soon make specific choices for these 2-cells and call their union a standard w -crossing. Each standard w -crossing is topologically a disk with boundary a loop corresponding to the relation the standard w -crossing represents.

Figure 31(i) shows the standard w -crossing corresponding to the relation $w_{ij}^L R_{ik} (w_{ij}^L)^{-1} = L_{jk}$, where any consistent markings of the graphs in the figure can be used. Of course, different markings correspond to different conjugates of the relation. We mention that figure 31(i) is essentially the pictorial version of the first computation in the proof of 1.3 in [7]. Figure 31(ii) is an abbreviation for the standard w -crossing in figure 31(i), and figure 31(iii) is an even briefer abbreviation for the same. Changing the orientation of the i edgesets in 31(i)




FIGURE 31.




FIGURE 32.

gives the standard w -crossing in 31(iv), shown in abbreviated form. Figure 31(v) shows (the abbreviated form of) another standard w -crossing obtained by changing the orientations of the j edgesets in 31(i) and then adding two triples of 2-cells representing the relation $w_{ij}^L = w_{ij}^R$. These triples are indicated by asterisks in 31(v). (Figure 31(iv) represents the second computation in the proof of 1.3 in [7], and figure 31(v) corresponds to a relation covered by 1.4, but not 1.3, in [7].

Figure 32(i) shows a second type of standard w -crossing. It is superimposed on the faint outlines of the six corresponding 2-cells of $\tilde{X}$. The colored graphs corresponding to these 2-cells should be clear. Any consistent markings may be used. Figure 32(ii) is an abbreviation which we will use for the standard w -crossing of figure 32(i).

Figure 32(iii) indicates a third type of standard w -crossing and figure 32(iv) shows an abbreviation for it. This crossing will only be used in Figure 55 below and discussion relevant to it. The use of the abbreviation is justified for our purposes since the choice of k and the orientations of the k -edgesets will not matter by Lemma 10.3. There are four standard w -crossings in 32(iii): two are as in 32(ii), but with the orientations of the edgesets changed, and two are as in 31(iii).

The last type of standard w -crossing are those involving four distinct subscripts and are abbreviated as in figure 32(iv). Each of these crossings consists of three 2-cells in $\tilde{X}$, where each 2-cell represents a trivial commutator relation.

The 2-cells in $\tilde{X}$ obtained from any one of the above standard w -crossings by changing the orientations of any combination of the i -, j -, or k -edgesets in all the cells of the crossing is also a standard w -crossing.

Let C be any standard w -crossing, thought of as in abbreviated form, which contains no asterisks. Then three other standard w -crossings are obtained from C by adding asterisks on either or both sides of the broken line in C , as in figure 31(v).

Let $\langle\langle A, x_k \rangle\rangle$ be any Whitehead element of the group StN_n , and let w be any generator of W_n . We next define a standard w -crossing C corresponding to the relation $w\langle\langle A, x_k \rangle\rangle w^{-1} = \langle\langle B, x_l \rangle\rangle$. C is the union of triangles in $\tilde{X}$ realizing the relations $\langle\langle A, x_k \rangle\rangle = \prod_{a \in A} \langle\langle a, x_k \rangle\rangle$ and $\langle\langle B, x_l \rangle\rangle = \prod_{b \in B} \langle\langle b, x_l \rangle\rangle$ with the standard w -crossings involving w and the $\langle\langle a, x_k \rangle\rangle$. The order of the factors in the products is arbitrary except that if $\langle\langle a_p, x_k \rangle\rangle$ is the p^{th} factor in $\prod_{a \in A} \langle\langle a, x_k \rangle\rangle$, then $w_{ij}^L \langle\langle a_p, x_k \rangle\rangle (w_{ij}^L)^{-1}$ must equal the p^{th} factor in $\prod_{b \in B} \langle\langle b, x_l \rangle\rangle$. Varying the order gives different standard w -crossings representing the same relations as C does, as does varying the standard w -crossings in C .

8.3. The subcomplex $\tilde{W}$ of $\tilde{X}$. For $n > 4$, let W_n be the group with generators W_{ij}^L and W_{ij}^R , where i and j are distinct integers between 1 and n , and the eleven relations

$$\begin{array}{ll}
(1^L) & [W_{ij}^L, W_{kl}^L] \\
(2^L) & W_{ij}^L W_{ik}^L (W_{ij}^L)^{-1} W_{jk}^L \\
(3^L) & W_{ij}^L W_{ki}^R (W_{ij}^L)^{-1} W_{kj}^L \\
(4^L) & W_{ij}^R W_{jk}^L (W_{ij}^L)^{-1} (W_{ik}^L)^{-1} \\
(5^L) & W_{ij}^L W_{kj}^L (W_{ij}^L)^{-1} (W_{ki}^L)^{-1} \\
(6) & W_{ij}^L = W_{ij}^R \\
(1^R) & [W_{ij}^R, W_{kl}^R] \\
(2^R) & W_{ij}^R W_{ik}^R (W_{ij}^R)^{-1} W_{jk}^R \\
(3^R) & W_{ij}^R W_{ki}^L (W_{ij}^R)^{-1} W_{kj}^R \\
(4^R) & W_{ij}^L W_{jk}^R (W_{ij}^R)^{-1} (W_{ik}^R)^{-1} \\
(5^R) & W_{ij}^R W_{kj}^R (W_{ij}^R)^{-1} (W_{ki}^R)^{-1}
\end{array}$$

where i, j, k , and l are distinct. Note that each of the relations (n^R) is obtained from (n^L) by interchanging each occurrence of L and R , and so is a consequence of (n^L) and relation (6).

Lemma 8.6. *There is an injective homomorphism $i: W_n \rightarrow StN_n$ induced by $i(W_{ij}^L) = w_{ij}^L$ and $i(W_{ij}^R) = w_{ij}^R$.*

Proof. Figure 33 shows how the image under i of the relations (2^L) and (3^L) are consequences of the defining relations of the group StN_n . Both 2-cycles in the figure consist of the union of three standard w -crossings, and, in the case of part (2^L) of the figure, a triple of 2-cells realizing the relation $w_{ij}^R = w_{ij}^L$. Figure 1 indicates, up to markings, these three 2-cells.

Inverting x_i in part (3^L) of the figure (by which we mean changing each occurrence of x_i to x_i^{-1} and each occurrence of x_i^{-1} to x_i) shows how the image of the relation (5^L) is a consequence of defining relations in StN_n . Similarly, inverting both x_j and x_k , and then all of x_i, x_j and x_k shows that relations (5^R) and (3^R) , respectively, map to relations in StN_n . Also, inverting x_i , then both of x_j and x_k , and finally all of x_i, x_j , and x_k in part (2^L) of the figure shows that relations (4^L) , (4^R) , and (2^R) , respectively, map to relations in StN_n . Each of (1^L) and (1^R) clearly map to relations, each of which are consequences of nine trivial




FIGURE 33. Realizing the relations (2^L) and (3^L) in $\tilde{X}_n$.

commutator relations in the definition of StN_n . Also, relation (6) maps to a defining relation for StN_n . This shows that i is well-defined.

That i is injective follows, for instance, from Theorem 3.3 and its proof in [7]. $\square$

Let $BW_n^{(2)}$ be the CW -complex with a single vertex, with oriented edges corresponding to the generators of W_n , and with 2-cells (attached in the usual way) corresponding to the above defining relations for W_n . The homomorphism i , both parts of Fig. 33 and the figures obtained from them by the inversions mentioned above, and Fig. 1 serve to define a cellular map $f: BW_n^{(2)} \rightarrow \tilde{X}_n/StN_n$. Let $EW_n^{(2)}$ be the universal cover of $BW_n^{(2)}$, and let $\tilde{f}: EW_n^{(2)} \rightarrow \tilde{X}_n$ be any lift of f which takes the vertices of $EW_n^{(2)}$ to those vertices of $\tilde{X}_n$ corresponding to the elements of the subgroup $i(W_n)$ of StN_n .

Let $\tilde{W}_n$ be the image of $\tilde{f}$, and let $\tilde{W}$ be the limit of the $\tilde{W}_n$ as $n \rightarrow \infty$. The image under $\tilde{f}$ of each 2-cell of $EW^{(2)}$ other than those corresponding to the relation (6) is called a standard ww -crossing. Such crossings may, by definition, also contain asterisks.

Abbreviations for standard ww -crossings are also shown in Figure 33. The asterisk in the abbreviation for the one corresponding to the relation (2^L) is often omitted, as are the labels, (2^L) and (3^L) in the figure, of the relations they correspond to.

9. LEMMA T

We next describe a number of types of 2-cycles in $\tilde{X}_n$. Any cycle of one of these types is called a conjugate cycle. Such cycles arise in §11 as part of the ambiguity in passing from 2-cycles of $\mathcal{T}$ to those of $\tilde{X}$.

Two examples of conjugate cycles are indicated in Figure 34 using the same conventions as in Figures 31 and 32. In general, there is, by definition, one type of conjugate cycle corresponding to each of the defining relations (R1)-(R3) in §8.3 for the group StN . Let C be a 2-chain in $\tilde{X}$ realizing such a relation. Each conjugate cycle of the corresponding type consists of standard w -crossings arranged along a circle surrounding C , as in the figure, together with a translate C^w of C on the opposite side of the circle. The chain C^w is only implied by the loose ends in the figure which are understood to join in a point like that in




FIGURE 34.



FIGURE 35. All reduced colored graphs with three vertices.

the center of (i) and (ii). The cells of C and C^w are called conjugate cells. We note that the marked colored graphs corresponding to the conjugate cells of C are the same as those of C^w , except for the color of two edgesets of each graph.

The circle in a conjugate cycle is called an ij -circle if crossing it in some direction corresponds to either w_{ij}^L or w_{ij}^R . An $\{i,j\}$ -circle is either an ij -circle or a ji -circle. Some points on an ij -circle may correspond to the relation $w_{ij}^L = w_{ij}^R$, as, for example, in Figure 34(ii). Such points are indicated by asterisks.

The relations (R1)-(R3) are realized by the reduced colored graphs with three vertices. These are shown in Figure 35. Thus conjugate cycles could have been defined in terms of these graphs. We generalize the definition of conjugate cycles slightly so that there is one type corresponding to each (not necessarily reduced) colored graph with three vertices. So the $\{i,j\}$ -circle of a conjugate cycle may surround a chain corresponding to any number of the relations (R1)-(R3), rather than just one.

Figure 34(i) indicates a conjugate cycle corresponding to a relation of type (R2) and 34(ii) to one of type (R1). The right of Figure 40 below shows another two conjugate cycles. Both correspond to a relation of type (R3). These two types of conjugate cycles are the only ones which are nontrivial in $H_2(\tilde{X})_{StN}$.

Lemma 9.1. *If $n \geq 5$, all conjugate cycles in $\tilde{X}_n$ are trivial as elements of $H_2(\tilde{X}_n)_{StN_n}$, except for those which map to nontrivial elements of $H_2(\tilde{X}_n/StN_n)$.*

Proof. We begin by showing that all conjugate cycles whose conjugate cells are reduced are trivial in $H_2(\tilde{X}_n)_{StN_n}$, except for the noted exceptions.

We first assume that the conjugate cells correspond to the graph numbered 1 in the figure, where k is the color of the two edges forming a cycle and i is the color of the other two edges. If the solid circle in the corresponding conjugate cycle is labeled either w_{kn} or w_{nk} , then the cycle is nontrivial in $H_2(\tilde{X}_n)$ since the composite $H_2(\tilde{X}_n) \rightarrow H_2(\tilde{X}_n/StN_n) \xrightarrow{\Theta} \mathbb{Z}^n$ takes this




FIGURE 36.

cycle to $\pm(e_k \pm e_n)$. The map Θ was defined in §6. So we assume that the subscripts are i and n , and further assume that the conjugate cycle is as shown in Figure 3, since changing the orientations of the edgesets in this figure give all other possibilities.

Let G and H be the colored graphs indicated in figure 36 marked in any way such that the marked roses $G_{b,d,f}$ and $H_{a,d,e}$ coincide. With cell orientations as indicated in the figure, let $g = \text{gen}(G)$ and $h = \text{gen}(H)$ be the corresponding generators of $H_3(\tilde{X})$. Then there is a 3-chain c in $\tilde{X}$ consisting of just additive prismatic terms such that $d(g + h + c)$ is the cycle indicated in figure 36. This cycle consists of six standard w -crossings, plus the terms corresponding to four relations of type (iv) in the group StN . The portion of the figure above the dotted line corresponds to terms of dg , the portion below to those of dh .

By adding two θ -prismatic terms to $g + h + c$, each of the two triples of cells nearest the dotted line realizing the relation $w_{ij}^L = w_{ij}^R$ in figure 36 can effectively be moved around a standard w -crossing so as to transform the picture in the center of figure 36 into the picture shown in the upper right of this figure. Thus the 2-cycle indicated in this last figure is a boundary. Changing the orientation of all of the i -edgesets in the above argument shows that the 2-cycle like that in the upper right of figure 36, but with the relations 3^R and -3^L replaced by 5^R and -5^L , is also trivial in $H_2(\tilde{X})$.

Now we assume that the conjugate cells of a conjugate cycle correspond to the graph numbered 2 in Figure 35. Let i, j and k be the colors of the edges of the graph corresponding to one of the conjugate cells. There are then six cases depending on whether the w -crossings involve w_{in} , w_{jn} or w_{kn} , or the same, but with the order of the subscripts switched. After, if necessary, moving asterisks across standard w -crossings as in the previous case, two of these six cases are treated in Figure 37. Each of the other cases are similar to one of these two. The six 3-cells on the left of the figure correspond to the colored graphs near them. The markings of these graphs can be any such that the indicated cancelations in the boundaries occur. The boundary of these six 3-cells is then a conjugate cycle of the type represented by




FIGURE 37.

Figure 34(i). In a similar way, the three 3-cells on the right show that conjugate cycles like those in Figure 34(i), except with w_{ni} in place of w_{jn} , are trivial as elements of $H_2(\tilde{X}_n)$.

Assume now that the conjugate cell corresponds to a colored graph with the underlying partitioned graph $G_1 \vee G_2$ numbered 3 in Figure 35. If at least one of i or j of the $\{i, j\}$ -circle of a corresponding conjugate cycle differs from the colors of $G_1 \vee G_2$, then the conjugate cycle is the boundary of the 3-chain corresponding to the marked colored graphs making up a w -crossing, each wedged with G_1 or G_2 . If both i and j are colors of $G_1 \vee G_2$, let $G'_1 \vee G_2$ be the colored graph C2(a) of figure 1 in eliminating cubical terms section with the colors of the singleton edges and monochromatic loops of G'_1 the same as those of G_1 . Consider the 3-chain C corresponding to $G'_1 \vee G_2$ and the prismatic cell which makes all the cells of ∂C reduced. Then the sum of the conjugate cycles corresponding to the 2-cells in ∂C is itself a boundary. But all of these conjugate cycles, except for three, have been considered in previous cases, so are boundaries. One of the three exceptional cycles, x say, corresponds to $G_1 \vee G_2$. The other two, call them y and y' , correspond to the same colored graph, but with opposite orientations. Thus $y + y' = 0$ in $H_2(\tilde{X})_{StN}$. So $x = 0$ in $H_2(\tilde{X})_{StN}$ since $x + y + y' = 0$ in $H_2(\tilde{X})$.

Assume now that the conjugate cycle corresponds to the graph numbered 4 in figure 35, where i is the color of the singleton edge and j and k are the colors of the other two edgesets. If the conjugate cycle contains either a $\{j, n\}$ - or a $\{k, n\}$ -circle, then previous cases essentially show that it is a boundary. If it contains an $\{i, n\}$ -circle, then replace each of the two conjugate cells with those five cells in the boundary of the 3-cell corresponding to the colored graph in Figure 36 containing two concentric circles which differ from the cell corresponding to earmuffs. Previous cases essentially show that the five conjugate cycles corresponding to these five cells are boundaries, which completes this case unless the conjugate cycle contains a $\{j, k\}$ -circle. This last case is similar to the last one considered above involving $G_1 \vee G_2$, but with the graph C2(b) in Figure 2 in place of $G'_1 \vee G_2$ of C2(a).

Now let the conjugate cell correspond to the graph numbered 5, with the color of the monochromatic loop k , and i and j the other two colors. Cases like previous ones show




FIGURE 38.

that conjugate cycles containing $\{i, n\}$ - and $\{j, n\}$ -circles are trivial. So we assume that the conjugate cycle Z has an nk -circle, specifically one corresponding to $w_{nk}^L = R_{nk}^{-1} L_{kn}^{-1} L_{nk}$. We show that Z is the boundary of a 3-chain in $\tilde{X}_n$ corresponding to 18 oriented 3-cells. Cells of these 18 which contain an edge corresponding to, respectively, R_{nk} , L_{kn} and L_{nk} , are called, respectively, lower, middle and upper cells. These three types of cells partition the 18 into three sets of six elements each.

The lower cells of Z are in bijective correspondence with the permutations of the set $\{i, j, n\}$, and, accordingly, triangulate a cube C in $\tilde{X}_n$ with four (oriented) edges corresponding to each of R_{ik} , R_{jk} and R_{nk} . Specifically, the lower cell which corresponds to the permutation p is $[p(n)^+ | p(j)^+ | p(i)^+]_k$. The edge path $[p(i)^+], [p(j)^+], [p(n)^+]$ consisting of three of the edges of this cell is also an edgepath in the boundary of C , one of the six (of length three) from the vertex $init([p(i)^+])$ of C to the diagonally opposite one $term([p(n)^+])$.

Let $P(a, b, c, d)$ and $V(a, b, c, d)$ be the colored graphs shown to the right of Figure 38. The rest of the figure indicates the colored graphs corresponding to four of the middle cells. The two other middle cells, which are simplicial, are also indicated, using bar notation. Markings and orientations are such that the indicated cancelations occur and such that two terms $[j^+ | i^+]_n$ and $[i^+ | j^+]_n$ to the lower right cancel with one face of the cube C . Cells enclosed in dotted circles correspond to terms of the conjugate cycle Z . This accounts for all the terms of the boundary of the middle cells except for the six indicated by the upward pointing arrows at the top center of the figure.

The upper cells correspond to the same colored graphs as do the middle cells, except that the colors k and n are switched, as are i and j . Orientations and markings are such that the six unaccounted for terms of the boundary of the middle cells cancel with the analogous six terms in the boundary of the upper cells. The remaining terms of the boundary of the upper cells which do not cancel among themselves are part of the conjugate cycle Z .




FIGURE 39.

The case of conjugate cycles involving the graph numbered 5 and an $\{i, j\}$ -circle is similar to the following case.

Assume now that the conjugate cell is rank 2, where the monochromatic cycle has color j and the other edge has color i . This cell and an adjacent one correspond to the crossing x just above the middle of the picture on the left of Figure 39. Note that previous cases have dealt with conjugate cycles containing conjugate cells corresponding to all the other crossings in the picture. So, to complete this case, it's enough to show that this picture represents a boundary. Indeed it's the boundary of a 3-chain c with two prismatic terms: one corresponds to the colored graph shown in the middle of Figure 39 and the other to the same colored graph, but with the orientation of the edgeset j reversed. Colorings and orientations are chosen so that contracting the edge j^0 in both graphs gives canceling terms in the boundary of the 3-cycle. The chain c also has three simplicial terms which isolate the crossing x from the one just to its right in the figure.

This completes the proof for conjugate cycles whose conjugate cells are reduced. By lemmas 6.5 and 6.6, we may replace any nonreduced conjugate cell with a homologous reduced chain. Previous cases then apply. $\square$

The following weaker version of Lemma 9.1 will often be cited.

Lemma 9.2 (Three Circles Lemma). *Let $n \geq 5$. Let P be a picture in $\tilde{W}_n$ (or a translate of it in $\tilde{X}_n$) which consists of three circles as shown on the right of Figure 39 with each crossing corresponding to a defining relation of the group W_n . If the only asterisks of P are on its single solid circle, then P represents the trivial element of $H_2(\tilde{X}_n)_{StN_n}$. If there are asterisks on any of the other edges of P , then P may only represent the trivial element of the quotient $H_2(\tilde{X}_n)_{StN_n}/\langle \emptyset \rangle$.*

10. PICTURES AND THEIR SIMPLIFICATION

Theorem 11.7 below implies that each element of $H_2(\tilde{X})_{StN}$ is represented by a 2-cycle in $\tilde{W}$. Such 2-cycles are studied in this section. Figure 40 shows four examples of these, drawn using the notation introduced at the end of §8.3. Numbers in the figure near each crossing indicate which defining relation of §8.3 the crossing corresponds to. Following Igusa, representations of simple spherical cycles as in the figure are called pictures.

Different pictures may represent the same cycle in $\tilde{W}$. For instance, pictures which differ only as shown within the dotted circles in either (a) or (b) of figure 41 are equal in $\tilde{W}$.




FIGURE 40. Two preimages of an exotic element of $K_3(\mathbb{Z})$, and another pair of nontrivial elements of $H_2(\tilde{X})_{StN}$.



FIGURE 41.

Since $\tilde{W}$ is a subcomplex of $\tilde{X}$, any picture gives a cycle in $\tilde{X}$. Let E be the subgroup of $H_2(\tilde{X})_{StN}$ generated by the classes which the pictures C_{ij}^2 and C_{ij}^4 for $1 \leq i, j$ and $I_1^{-3,-4}$ represent. Notation is that of Figure 40. A subgroup E_n of $H_2(\tilde{X}_n)_{StN_n}$ is defined similarly, but with $i, j \leq n$. The letter E is used in the notation for these subgroups since, as we will see, the picture $I_1^{-3,-4}$ maps to what has been called an exotic element of $K_3(\mathbb{Z})$. (Exotic elements are those not in the image of the map $\pi_3(B\Sigma^+) \rightarrow \pi_3(BGL(\mathbb{Z})^+)$.) Let $\langle \mathcal{O} \rangle$ be the subgroup of E or E_n generated by the classes of the pictures C_{ij}^2 and C_{ij}^4 on the right of Figure 40. Most of this section is devoted to proving the following.

Theorem 10.1. *Let P be any picture in $\tilde{W}_n$ where $n \geq 5$. Then the class of P in $H_2(\tilde{X}_n)_{StN_n}$ is in the subgroup E_n .*

This, along with theorem 11.7 below, shows that $H_2(\tilde{X}_n)_{StN_n} = E_n$.

Numerous preparatory lemmas and some definitions proceed the proof proper of the theorem. The first lemma we give is a convenience which shows that deleting the L and R superscripts and the asterisks from any picture leaves a well-defined element of $H_2(\tilde{X})_{StN}/\langle \mathcal{O} \rangle$.




FIGURE 42.

Lemma 10.2. *Changing the superscripts L and R and adding or deleting asterisks in any picture does not change the class of the picture in $H_2(\tilde{X}_n)_{StN_n}/\langle \mathcal{O} \rangle$ if $n \geq 5$.*

This follows from the triviality of the conjugate cycles corresponding to the relation (R3) considered at the start of the proof of Lemma 9.1.

By forgetting some information, any picture P in $\tilde{W}_n$ may be thought of as a graph, also denoted by P , on S^2 with vertices, all 4-valent, corresponding to the crossings (and with some of the edges of the graph possibly containing asterisks). Thus we will speak of edges of a picture. (Edges of pictures will be given a somewhat more general meaning after lemma 10.5 below.) Because of lemma 10.2, transversely oriented edges of pictures are often labeled by just two letters, with, for instance, ij denoting either w_{ij}^L or w_{ij}^R . Of course, a picture drawn with this abbreviated notation only determines a well-defined element of $H_2(\tilde{X}_n)_{StN_n}/\langle \mathcal{O} \rangle$ and not a homology class in $\tilde{W}_n$, nor in $\tilde{X}_n$.

Figure 42 shows a few pictures drawn with this abbreviated notation. The top and bottom portions of the picture on the left are examples of what we call swirls. Swirls are of two types: knot-like and circular. A knot-like (resp. circular) swirl is, by definition, any 2-chain in $\tilde{X}$ with corresponding unlabeled partial picture as shown in the top (resp. bottom) portion of the picture on the left of Figure 42. There are eight types of knot-like swirls, the two shown in figure 42 on the right, with three more obtained from each of these by changing the direction of each of i , j , and k . By changing the direction of, for instance, i , we mean changing the orientation of all the i -edgesets of the colored graphs corresponding to the terms of the cycle a given picture represents. For instance, the swirl at the bottom of the picture on the left of Figure 42 is obtained from the one in the upper right by changing the direction of j . Another four knot-like swirls are shown in figure 51 below.

Swirls will often arise in the course of the proof of theorem 10.1. In fact, the proof will show that, by introducing swirls in any given picture P , the number of crossings in P which are not part of swirls, can always be reduced. Thus P is homologous in $\tilde{X}$ (modulo elements

of $\langle \mathcal{O} \rangle$) to a picture consisting solely of swirls. Lemma 10.19 deals with pictures of this last sort.

Given a picture P , let P_k be the subgraph of P consisting of all the edges having label w_{ab}^L or w_{ab}^R where either $a = k$ or $b = k$. Edges of P_k are often referred to as k -edges. Each vertex of P_k is either bivalent or trivalent as a quick examination of the defining relations in §8.3 shows. Delete the bivalent vertices from P_k , and join the two edges meeting each of these into one edge. Denote by P_k^* the result of this process. Note that, as subsets of S^2 , we have $P_k^* = P_k$. In case P_k has any components with only bivalent vertices, the corresponding components of P_k^* are circles with no vertices. These circles are referred to as circular edges. The (possibly empty) complement of the circular edges in P_k^* has the structure of a graph with all vertices trivalent. Any reference to a vertex, edge, or cycle of P_k^* refers to one of this graph.

Here is another application of lemma 9.1.

Lemma 10.3. *Let P be a picture in $\widetilde{W}_n$. Changing the transverse orientations of all the k -edges in any one component of P_k does not change the class of P in $H_2(\widetilde{X}_{n+2})_{StN_{n+2}}/\langle \mathcal{O} \rangle$, nor does changing the subscript k of all the k -edges in one of the components to $n+2$.*

Proof. Write the cycle P as $C_k + C'_k$ where the terms of C_k are the generators of $C_2(\widetilde{X}_n)$ corresponding to the crossings in one component of P_k and the terms of C'_k are those of the other components. Let C_{n+1} be the same chain as C_k , except with each appearance of the subscript k in the corresponding cycle changed to $n+1$. Then

$$P = (C_k - C_{n+1}) + (C_{n+1} + C'_k)$$

where both $C_k - C_{n+1}$ and $C_{n+1} + C'_k$ are cycles. In $H_2(\widetilde{X}_{n+2})_{StN_{n+2}}/\langle \mathcal{O} \rangle$, we have

$$C_k - C_{n+1} = w_{n+2,k}^L \cdot (C_k - C_{n+1}) = C_{n+2} - C_{n+1},$$

where C_{n+2} is the chain C_k with k changed to $n+2$ in the corresponding picture. Thus $P = C_{n+2} + C'_k$, which proves the last assertion of the lemma. The first is proved similarly by replacing $w_{n+2,k}^L$ in the above equation with $(w_{n+2,k}^L)^2$. $\square$

Let P be a picture such that P_k^* contains a circular edge e . Let $P - e$ denote the picture obtained from P by deleting all the edges of P which are contained in e . The next lemma is often used and, for instance, shows, along with lemma 9.1, that swirls can be moved from one side of an edge of a picture to another.

Lemma 10.4 (Broken Circles Lemma). *If $n \geq 5$, then P and $P - e$ represent the same class in $H_2(\widetilde{X}_n)_{StN_n}/E_n$.*

The following proof actually shows that P and $P - e$ represent the same class in $H_2(\widetilde{X})_{StN}/\langle \mathcal{O} \rangle$. The weaker statement of the lemma is explained by remark 10.20 below.

Proof. Let e be a circular edge as in the statement of the lemma. Let α_1 and α_2 be the two loops in S^2 obtained by deforming e slightly off itself in each of the two possible directions. Let A_1 and A_2 be thin annular neighborhoods of α_1 and α_2 disjoint from e , and let A be the annulus consisting of $A_1 \cup A_2$ together with the region between α_1 and α_2 which contains e . Since all the vertices of P_k contained in e are bivalent, we may assume that the two words in W_{n-1} associated with the loops α_1 and α_2 are the same. Denote this one word, the trivial element of W_n , by w . Pick a specific trivialization of w using the defining relations of the group W_{n-1} , that is, write w as a product of conjugates of these relations. (For this, we




FIGURE 43.




FIGURE 44.

need that the map $W_{n-1} \rightarrow W_n$ is injective, which follows, for instance, from the proof of 3.3 in [7].) Then use this trivialization (four times) to deform P only in the neighborhoods A_1 and A_2 to a picture P' which is disjoint from α_1 and α_2 . Do this so that $A \cap P'$ admits an involution which preserves labels and transverse arrows, which fixes e and which interchanges the two portions of $A \cap P'$ on each side of e .

Let C be the component of P' which contains the component e . The involution pairs those vertices of C which are not contained in e . These pairs can be eliminated, one pair at a time, using the three circles lemma 9.2 with either a deformation like that in Fig. 43 or, if the crossings at the paired vertices v_1 and v_2 in the figure are different, by using a deformation somewhat simpler than the one in the figure. Both types of deformations preserve the $\mathbb{Z}_2$ -symmetry of $a \cap P'$ and do not change P_k^* . Thus we may assume that C consists of the circle e together with solid circles. These solid circles can be eliminated (using lemma 9.1 if necessary).

So now we assume that C consists of just the circle e . Delete e from P' . This does not change the class of P' in the group $H_2(\tilde{X})_W$ of coinvariants. Next eliminate the vertices in $A_1 \cup A_2$ of the remaining picture with a deformation which essentially reverses the way in which they were introduced. The resulting picture equals $P - e$ in $H_2(\tilde{X}_n)_{StN_n}/\langle \mathcal{O} \rangle$. $\square$

Lemma 10.5 (Swirl Moving). *Let $n \geq 5$. If two pictures agree except on neighborhoods as shown in (a) and (b) or as in (c) and (d) of figure 44, then the two pictures are equal in $H_2(\tilde{X}_n)_{StN_n}/\langle \mathcal{O} \rangle$.*

Proof. It follows from the broken circles lemma 10.4 that two pictures which differ only as shown in (c) and (d) are equal. If they differ as shown in (a) and (b) they are equal by lemma 9.1, with the help of Lemma 10.3 if necessary. $\square$

Since swirls can be moved “across” edges, they will be, to some extent, be ignored till lemma 10.19 below, where pictures consisting solely of swirls are considered. So the collection of edges making up a swirl will often itself be referred to as an edge. Pictures are often drawn accordingly without explicitly showing swirls (as in figure 45) which may be contained in their edges. Or a swirl may be indicated as on the right of figure 46(c). Edges which contain no swirls may be referred to as swirl-free.




FIGURE 45.




FIGURE 46.

We next discuss conventions which are used in Figure 45 and later figures. Edges in these labeled by i and j are i - and j -edges, respectively, of P_k^* , while dotted edges are edges of P , but not P_k^* . In pictures without arrows, any consistent choice of these can be used.

Lemma 10.6. *If two pictures differ only as shown in (a) and (b) or in (a) and (c) of Figure 45, then, after introducing a swirl on one of the edges in the figure, the two pictures are equal in $H_2(\tilde{X}_n)_{StN_n}/\langle \mathcal{O} \rangle$ if $n \geq 5$.*

Note that an application of this lemma change P_k , but not the unlabeled graph P_k^* .

Proof. There are essentially eight cases depending on which of the four defining relations of the group W the vertex in Fig. 45(a) corresponds to and whether the two pictures differ as shown in (a) and (b) or in (a) and (c). These cases correspond to the eight types of swirls mentioned above. For one case, assume that a picture P^a has a neighborhood like that shown in figure 46(a), while another picture P^b is obtained from P^a by changing this neighborhood to the one shown in Figure 46(b). By lemma 10.5, we may assume that the edge of P_k^* (completely) shown in 45(b) is swirl-free. Then $P_a - P_b$ is the picture of 46(c) which is clearly trivial for a suitably chosen swirl.

For another case, assume the labels ik in figure 46 are changed to ki . Then the swirl appears on a vertical k -edge of figure 46(b), but otherwise this case is similar to the previous one. The remaining cases are similar to the two just considered. $\square$

Circular swirls will arise with applications of the following.

Lemma 10.7. *Assume a picture P has a neighborhood like that in Fig. 47(a). Then this neighborhood can be deformed to one like that in either (b) or (c) giving a picture P' equal to P in $H_2(\tilde{X})_{StN}/\langle \mathcal{O} \rangle$, provided a suitable swirl is introduced on one of the ij -edges of P' .*

Proof. There are numerous cases depending on the directions of the transverse arrows on the edges and the labels of the edges. For one case, assume two pictures differ by the neighborhoods shown in (a) and (b) of figure 48. By the swirl moving lemma 10.5, we may




FIGURE 47.




FIGURE 48.

assume that the three edges forming the triangle in figure 48(a) are swirl-free. Then the difference of these two pictures in $H_2(\tilde{X})$ is represented by the picture in figure 48(c). This is trivial for a suitable choice of a circular swirl. Other cases are similar. $\square$

An edgepath in a graph G (which could contain circular edges) is a sequence $e_1, \dots, e_n$ of oriented (noncircular) edges of G such that $\text{term}(e_i) = \text{init}(e_{i+1})$ for $i = 1, \dots, n-1$. An edgepath $e_1, \dots, e_n$ is a cycle if $\text{term}(e_n) = \text{init}(e_1)$, and a simple cycle if, in addition, the vertices $\text{term}(e_i), i = 1, \dots, n$ are distinct.

The next lemma is used implicitly in the definition following it.

Lemma 10.8. *Every cycle of P_k^* consists of at least two edges.*

Proof. Suppose a single edge e forms a cycle in P_k^* . Then the loop in S^2 associated with e can be deformed slightly to a loop ω such that the intersection $P_k \cap \omega$ is transverse and consists of a single point contained in that edge of P_k which is incident with $\text{init}(e) = \text{term}(e)$ and which is different from e . But the element of the group W associated with this loop cannot be trivial since, for instance, the element of $\text{Aut}(F_n)$ it maps to does not take x_k to x_k . This contradiction completes the proof. $\square$

Let P be a picture. Fix an orientation of the 2-sphere containing P_k . This determines, at each vertex v of P_k^* , a cyclic ordering of the three edges of P_k^* incident with v . An edgepath e_i, e_j is said to be right-turning (at the vertex $\text{term}(e_i) = \text{init}(e_j)$) if e_i, e_j, e is this cyclic ordering at the vertex $\text{term}(e_i)$, where e is the third edge of P_k incident with $\text{term}(e_i)$. A cycle $e_1, \dots, e_n$ in P_k^* is right-turning if all the subedge-paths e_i, e_{i+1} for $i = 1, \dots, n-1$ as well as e_n, e_1 are right-turning. A left-turning cycle is one which is right-turning with respect to the orientation opposite to the fixed one. A cycle is monotonic if it is either left-turning or right-turning. A cycle in P_k is monotonic if the corresponding cycle in P_k^* is monotonic.

Lemma 10.9. *Every monotonic cycle in P_k is simple.*

Proof. It suffices to show that monotonic cycles in P_k^* are simple. Suppose $\omega = e_1, \dots, e_n$ is a monotonic cycle in P_k^* , right-turning say, which is not simple. Choose a simple subcycle ω' of ω which is right-turning at all its vertices except for one, v say. Then ω' can be deformed slightly to a loop in S^2 which meets P_k transversely in a single point, namely a point of that edge of P_k incident with v which is not part of ω' . The proof can now be completed as was the previous one. $\square$

Each monotonic cycle ω bounds two closed disks in S^2 . One of these contains the component C of P_k which contains ω . The other has interior disjoint from C . This latter disk is called a face of C , while the former is called a coface. The interiors of the faces of C partition $S^2 - C$. The edges of P_k^* in the cycle forming the boundary of a face are often referred to as the edges of the face. A face with n edges is called an n -gon. We call 4-gons, 3-gons, and 2-gons, respectively, squares, triangles, and bigons, respectively.

Fix a point $x \in S^2 - P$. Associate with each component C of P_k the coface of C which does not contain x . These cofaces have disjoint boundaries, and so are partially ordered by inclusion. Components of P_k which correspond to minimal or innermost such cofaces are called innermost components of P_k .

Let C_k be an innermost component of P_k , and let C_k^* denote the component of P_k^* corresponding to P_k . The face of C_k containing x is called the backface of C_k . The other faces are sometimes called frontfaces. The interiors of frontfaces are disjoint from P_k , while the interior of each backface meets P_k , unless P_k is connected.

A face of P_k is, by definition, a clear face if the interior of the face does not meet P . We say that a frontface F can be cleared if there is a deformation of P which makes F a clear face without changing P_k^* .

Lemma 10.10 (Clearing Lemma). *Every front-face of an innermost component C_k of a picture P can be cleared.*

Proof. Let F be such a front-face. Use lemma 10.6 to deform P so that $\text{int}(F) \cap P = \emptyset$ near each vertex of C_k^* in ∂F . Then vertices of $\text{int}(F)$ which are joined by a single edge of P to a vertex of ∂F can be eliminated, one at a time, using three circles lemma 9.2 or the broken circles lemma 10.4. Figure 49 shows one way to do this for vertices in $\text{int}(F)$, which are joined by an unbroken edge to ∂F . The deformation for the other type of vertex in $\text{int}(F)$ is simpler.

So now we assume that any remaining vertices in $\text{int}(F)$ are contained in components of P different from the one of ∂F . An edge of one of these components can be deformed to meet ∂F in two unbroken crossings. Then the vertices of what was this component can be eliminated from $\text{int}(F)$ as above.

By repeating the deformations described above as necessary, we may assume that all edges of P in $\text{int}(F)$ meet two vertices in ∂F . These edges can be deformed outside of F using the three circles lemma and lemma 10.7. $\square$

The clearing lemma just above often makes some of the following four lemmas applicable.

Lemma 10.11. *If F is a clear face of C_k with just one l -edge, for some fixed l , then there is a deformation of P which decreases the number of vertices of P_k^* .*

Proof. Assume F is a clear face with just one l -edge. Use lemma 10.6, if necessary, to make the single l -edge of F solid at both its vertices. If F is a bigon, it is then easy to deform P




FIGURE 49.




FIGURE 50.

in a neighborhood of F so as to eliminate these two vertices. This decreases the number of vertices of P_k^* by two.

If F is an n -gon, where $n > 2$, then a deformation like that shown in figure 50 (or a similar deformation if the j -edge in the figure is broken at both of its vertices) produces a clear face with one l -edge and one fewer edge than F . This deformation changes C_k^* , but does not change the number of vertices of C_k^* . Repeating such deformations gives a clear bigon. The number of vertices of P_k^* can then be reduced as above. $\square$

Two l -edges of a clear face F are said to be consecutive if there is an edgepath in ∂F which joins the two edges and contains no other l -edges.

Lemma 10.12. *Assume F is a clear face of C_k . If two consecutive l -edges of F have either both inward- or both outward-pointing transverse arrows, then there is a deformation of P which decreases the number of vertices of P_k^* .*

Proof. Let e and f be consecutive l -edges of F which have either both inward- or both outward-pointing transverse arrows. Surgery in F as in Figure 41(a) using e and f produces two clear faces, at least one of which has just one l -edge. So the previous lemma can be applied. $\square$

Lemma 10.13. *If F is a clear face with an odd number of l -edges, then there is a deformation of P which decreases the number of vertices of P_k^* .*

Proof. If the number of l -edges is odd and greater than one, then the arrows on all pairs of consecutive l -edges do not alternate between inward- and outward-pointing. So the previous lemma applies. $\square$

Lemma 10.14. *If F is a clear face of C_k with i -, j -, and l -edges, where i, j , and l are distinct, then there is a deformation of P which decreases the number of vertices of P_k^* .*

Proof. If ∂F contains three types of edges, then it has three consecutive edges of different types. We may assume that these are arranged as in figure 50(a). Then either F has an odd number of j -edges or the deformation in figure 50 produces a clear face with an odd number of j -edges. Either way, the previous lemma can be applied. $\square$

While the clearing lemma 10.10 will be used to provide clear faces to which lemmas 10.11–10.14 can be applied, the clearing lemma itself can only be applied to frontfaces. The next three lemmas show that faces to which lemmas 10.11–10.14 can potentially be applied occur in numbers greater than one. Since C_k^* has only one back-face, at least one of these must be a front-face.

Lemma 10.15. *No component C_k^* of P_k^* can have exactly one face with an odd number of edges.*

Proof. Let F be a face of C_k^* with an odd number of edges. Since the interior of each edge of C_k^* meets two faces we have

$$2(\text{number of edges of } C_k^*) = \text{number of edges of } F + \sum_{G \neq F} (\text{number of edges of } G)$$

where the sum is over all faces G of C_k^* different from F . It follows that some face other than F has an odd number of vertices. $\square$

We mention that with slight modifications, the above proof shows that if all the faces of C_k^* are $2n$ -gons, then every cycle in C_k^* consists of an even number of edges.

We say that a face F of C_k^* is surrounded by squares if each edge of F is also the edge of a square (different from F , in case F is itself a square). We say that C_k^* is n -regular if C_k^* has two disjoint n -gons F_1 and F_2 and another n squares for faces, if each edge of C_k^* not in $F_1 \cup F_2$ meets both F_1 and F_2 , and if whenever two of these edges meet consecutive vertices of F_1 , they also meet consecutive vertices of F_2 .

If C_k^* is n -regular then, using the above notation, both F_1 and F_2 are surrounded by squares. Conversely, it can be shown that if C_k^* has an n -gon which is surrounded by squares, then C_k^* is n -regular. But we only need and prove the following partial converse.

Lemma 10.16. *Assume all faces of C_k^* are $2m$ -gons for (possibly varying) $m > 1$. If some face of C_k^* with n edges is surrounded by squares, then C_k^* is n -regular.*

Proof. Assume F is a face of C_k^* which is surrounded by squares. Suppose an edge of $C_k^* - \partial F$ joins two vertices, say v and w , of F . Let $e_1, \dots, e_p$ be a simple edge path in ∂F from v to w . Since all faces of C_k^* are $2m$ -gons, all cycles in C_k^* have even length, and so p is odd. Note that $p \neq 1$ since none of the faces of C_k^* are bigons by assumption. Since e_1, e, e_p is a monotonic path meeting, but not contained in F , there must be an edge not in ∂F joining $\text{init}(e_1)$ with $\text{term}(e_p)$. Here we are using the orientations which make e_1, e, e_p an edgepath. Repeating this argument as necessary shows that e_m , where $m = \frac{1}{2}(p+1)$, is the edge of a bigon. Therefore no edge of $C_k^* - \partial F$ joins two vertices of F .

Now let v be a vertex of F and let e be the unique edge of $C_k^* - \partial F$ incident with v . Let $f(v)$ be the other vertex of e . This defines a function f from the vertices of F to the vertices of $C_k^* - \partial F$. Suppose f is not injective. Choose consecutive vertices $v_1, \dots, v_n$ of F such that $f(v_1) = f(v_n)$ and such that f is injective when restricted to each of the sets $\{v_1, \dots, v_{n-1}\}$ and $\{v_2, \dots, v_n\}$. Note that $n > 2$ since C_k^* has no cycles of length three.

If $n = 3$, consider the square S_1 with vertices $f(v_1), v_1, v_2$, and $f(v_2)$ and the square S_2 with vertices $f(v_3), v_3, v_2$, and $f(v_2)$. Note that the single edge joining v_2 to $f(v_2)$ is an edge of both squares. Let e_1 be the edge of S_1 joining $f(v_2)$ and $f(v_1)$, and let e_2 be the edge of S_2 joining $f(v_2)$ and $f(v_3) = f(v_1)$. If $e_1 \neq e_2$, then the vertex $f(v_1)$ has valence > 3 , a contradiction. And if $e_1 = e_2$, then S_1 and S_2 share two consecutive edges, another contradiction.

So assume $n > 3$. But then each of the four distinct vertices $v_1, f(v_2), f(v_{n-1})$, and v_n is joined by an edge to the trivalent vertex $f(v_1)$, which is impossible. This proves that the function f is injective. The proof of the lemma is now easily completed. $\square$

By a biggest face of C_k^* , we mean a face which has as many or more edges than all the other faces of C_k^* .

Lemma 10.17. *Assume all faces of C_k^* are $2m$ -gons, where $m > 1$. If F is a biggest (nonsquare) face of F which is not surrounded by squares, then at least two edges of F are not edges of squares.*

Proof. Suppose that F is a biggest face of C_k^* which is not surrounded by squares, and that all the edges of F , except one, are edges of squares. Let c be the one exceptional edge of F , and let G be the face of C_k^* other than F having c as an edge. Note that $F \cap G = c$. The first paragraph of the previous proof can now be applied (provided the edgepath $e_1, \dots, e_p$ in that proof is chosen so that it does not include c) and shows that an edge of $C_k^* - \partial F$ cannot join two vertices of F . The second paragraph also applies (provided the consecutive $v_1, \dots, v_n$ are chosen so that both vertices of c are not among them) and shows that the function f defined as above is injective.

Next we show that the image of f contains only two vertices of G . Suppose there is an edge e not in $F \cup G$ which joins a vertex v_F of F to a vertex v_G of G . Let $e_1, \dots, e_n$ be a simple edgepath in $\partial F \cup \partial G - \text{int}(c)$ from v_F to v_G . Here $\text{int}(c)$ is the interior of the edge c . Note that n is odd as remarked after the proof of Lemma 10.15. Since e_1 is an edge of F different from c , the monotonic path e_1, e, e_n is contained in a square. Let e' be the fourth edge of this square. Note that e' is not in $F \cup G$ since faces don't share consecutive edges. If e' is incident with a vertex of c , then this vertex would have valence > 3 , a contradiction. Otherwise, repeat the above argument with e' in place of e and the edgepath $e_2, \dots, e_{n-1}$ in place of $e_1, \dots, e_n$. Continuing in this way produces a contradiction, and proves that if x and y are the vertices of c , then the only vertices of G in the image of f are $f(x)$ and $f(y)$.

Now let $v_1, \dots, v_n$ be the vertices of F , listed so that v_i and v_{i+1} are vertices of the same edge and where $x = v_1$ and $y = v_n$. Then the face of C_k^* which contains the edge joining $f(v_i)$ and $f(v_{i+1})$ for $i = 1, \dots, n-1$ also contain two edges of G and therefore is bigger than F . This contradiction completes the proof. $\square$

Sometimes the following lemma is applicable when none of the previous four are.

Lemma 10.18. *Assume two faces of an innermost component C_k of a picture P meet in a disconnected set. Then there is a deformation of P which divides C_k into two nonempty components, at least one of which is innermost. Moreover, the number of vertices of P_k^* is unchanged.*

Proof. Assume F and G are faces of C_k having edges e and f in distinct components of $F \cap G$. At least one of F or G is a front-face, say F is. Clear F . Let α be a loop in S^2 which meets each of the edges e and f transversely in one point, and otherwise is contained in $\text{int}(E) \cup \text{int}(F)$. Let w be the word in the group W corresponding to α . Note that the two letters a and b , say, in w which correspond to the edges e and f may be assumed to be adjacent and that these are the only letters in w which have k in their subscripts.

Since w represents the trivial element of the group W , it follows that the subscripts of a and b are made up of just two letters. For, if for instance $a = w_{ki}^L$ and $b = w_{kj}^L$, then even on the level of the symmetric group, k would not be fixed. So $a = b^{-1}$ or if, if the subscripts




FIGURE 51.

of a and b are in opposite order, we may assume $a = b^{-1}$ by introducing a pair of canceling swirls on either e or f . So surgery in F as in figure 41 using e and f can be done. Its easy to check that if the basepoint $*$ of S^2 is in $F \cup G$, then the surgery can be done so that both of the created components are innermost. And if $*$ $\in S^2 - (F \cup G)$, then one of these components is innermost. $\square$

The next two lemmas are also used in the proof of Theorem 10.1.

Lemma 10.19. *If a picture P in $\widetilde{W}_n$ consists of disjoint circular edges, then P represents an element of E_n provided $n \geq 5$.*

Proof. If there are no swirls on P , then P represents the trivial element. Suppose P contains a circular swirl. Let P' be the subgraph of P consisting of this swirl, and assume that k is the third subscript of P' , with i and j the other two. Then either $P'_i \equiv P_i \cap P'$ or $P'_j \equiv P_j \cap P'$ contains a clear bigon, say P'_i does. Use lemma 10.6 to eliminate this bigon. After this, P'_j contains a clear bigon. Eliminate it using lemma 10.6 again. Then after moving the two swirls introduced with the two applications of 10.6, P' consists of one edge with three knot swirls. Thus we assume that all the swirls of P are knot swirls. These are even in number, and P equals, in $H_2(\tilde{X})$, a picture consisting of half as many disjoint circular edges with two knot swirls on each.

So we now assume that P contains exactly two knot swirls, S_1 and S_2 . Let ij and ji be the labels of the two edges of P which meet both S_1 and S_2 . The single edge of each swirl which meets both of these edges is called the long edge of the swirl. By Lemma 10.3, we may assume that the arrows on the long edges of S_1 and S_2 point inward, that is in the same direction as those in Figure 51. Then the classification of swirls shows that, up to permuting the subscripts i and j , each of the swirls S_1 and S_2 is one of the four in Figure 51.

As a first case, assume that S_1 and S_2 are the same swirl. Then either they cancel trivially at the chain level giving $P = 0$ in $H_2(\tilde{X})_{StN}$, or the positions of the subscripts i and j in one of S_1 or S_2 are different from those in the Figure. In the latter case, the broken circles

lemma 10.4 can be used to effectively move a crossing of one of the swirls, say S_1 , from one side of it to the other. Then, after using Lemma 10.3 if necessary to change the direction of the arrows of the edges of S_1 with labels containing the third subscript, we have $P = 0$ at the chain level giving $P \in E_n$.

For another case assume that S_1 and S_2 are either the two swirls in the top row of Figure 51 or the two in the bottom row. By a previous case, we may assume that the ij subscripts are just as in the figure, and, by Lemma 10.3, that the third subscripts of both swirls are the same. Two crossings of P then cancel leaving one of the generators of E shown on the left of Figure 40.

For the last case, assume that one of S_1 or S_2 , say S_1 , is in the top row of the figure and the other is in the bottom. By Lemma 10.3, we may assume that the third subscript of S_1 and S_2 are different. Then the three circles lemma 10.4 can be used to move a crossing of S_2 from one side of this swirl to the other. This changes the picture P so that the position of the third subscript of S_2 preceeds the other subscript in each of the labels in which the third subscript appears. Previous cases now apply to complete the proof. $\square$

Remark 10.20. Lemma 10.4, the Broken Circles Lemma, which was stated and proved before the conventions that edges of pictures could contain swirls, is still true with this more general notion of edges. The proof of 10.4, appended as follows, shows this.

At the start of the proof move all swirls near, but not on e away from e outside of the annulus A . The edge e itself may contain swirls. Now, with the help of the swirl moving lemma, the proof of Lemma 10.4 goes through till the last paragraph. So assume the circle C there contains swirls. Deform C into two circles, one of which contains all the swirls on C and bounds a disk with interior disjoint from the picture. The other circle should contain no swirls. The resulting picture equals $P_1 + P_2$ in $H_2(\tilde{X})$ where P_1 is represented by the former circle and P_2 is one of the pictures considered in the proof of Lemma 10.4 where it was shown to equal $P_2 - e$, modulo an element of $\langle \mathcal{O} \rangle$. This completes the proof since P_1 represents an element of E_n by lemma 10.19.

Proof of Theorem 10.1. Let P be any picture wt_n . Let C_k be an innermost component of P_k . We show that there exists a deformation of P which either decreases the number of vertices of P_k^* or disconnects C_k , without changing the number of vertices of P_k^* , and gives an innermost component whose vertices are a proper subset of those of C_k^* . This deformation does not enlarge the set of subscripts of those w_{ij} which are labels of the edges of P . Nor does it change the class of P in $H_2(\tilde{X}_n)$, except possibly by an element of E_n . Since the number of vertices of C_k^* is finite, iterating such deformations eventually gives one which reduces the number of vertices of C_k^* . Thus all the vertices of C_k^* can be eliminated, leaving only circular edges. These can be eliminated using lemma 10.4, except for an innermost circular edge with swirls. So the theorem will follow from the previous lemma.

If C_k^* has a face with an odd number of vertices, then it has, by lemma 10.15, at least two such faces. At least one of these must be a front-face. The Clearing Lemma 10.10, and then lemma 10.13, can be applied to any such front face. So we now assume that all faces of C_k^* have an even number of edges.

If C_k^* contains a bigon, then either C_k^* consists of two vertices and three edges, or C_k^* has two faces which meet in a disconnected set. In the first case C_k^* contains three bigons, at least one of which is a front-face. Thus the number of trivalent vertices can be reduced by




FIGURE 52.

first applying the clearing lemma, and then using lemma 10.11. In the second case, lemma 10.18 can be applied to disconnect C_k .

So now we assume that the number of edges of each face of C_k^* is even and at least four.

Choose a biggest face F (as defined just before lemma 10.17) of C_k^* . Assume that F is surrounded by squares. (The case in which F is not surrounded by squares is considered in the last part of the proof.) By lemma 10.16, we may assume that F is a front-face. Clear F . Then by lemma 10.14, we may assume that F has edges of just two types, say i - and j -edges.

If F has more than four edges, then there must be two i -edges which have either both inward- or both outward-pointing arrows. Surgery using these two edges produces a front-face which meets what was the other largest face of C_k in at least two components. Thus C_k can be disconnected using lemma 10.18.

So assume F has four edges. We first will deform P so that a disk neighborhood of C_k is as shown in figure 52, with the back-face of C_k^* corresponding to the unbounded face in the figure. We then will show that after adding a multiple of *exotic* to P , the number of 3-valent vertices of P_k can be decreased.

The face of C_k^* which does not meet the back-face is called the inner face of C_k^* (because of the position of the corresponding face in figure 52). Clear the inner face. Then, if necessary, use lemma 10.6 to deform P in neighborhoods of the four vertices of the inner face so that the unlabeled graph P agrees with the corresponding neighborhoods in the figure. Similarly, deform P in small neighborhoods of the four vertices of the backface so that P agrees with the corresponding neighborhoods in the figure.

The four edges joining the inner and back-face of C_k are called diagonal edges. Any crossing of P on the interior of a diagonal edge must be a broken k -crossing. These crossings can be pushed off the diagonal edges in the direction of the backface using lemmas T and 10.7. This may introduce swirls and crossings on the edges of the backface, but C_k^* is unchanged. Now the only crossings in P on the interiors of the edges of C_k^* are those on the edges of the backface. These are all broken k -crossings. So P can be deformed, using the broken circles lemma 10.4, so that, up to the labels of the edges, a disk containing C_k meets P in the graph shown in figure 52.

We assume that the edges of the inner square are of only two types, say i - and j -edges, and transversely oriented as in the figure. For otherwise, the number of vertices of C_k^* can be reduced as above using lemmas 10.12 and 10.13. With this assumption, each diagonal edge of C_k is either an i - or a j -edge, and the dotted edges in the figure are all $\{i, j\}$ -edges. So if the backface contains an l -edge, where l is different from i and j , then all the edges of the back-face are l -edges and any face meeting the backface, when cleared, contains just one l -edge. Because of lemma 10.11, we may assume, as above, that the edges of the back-face are all either i - or j -edges with tranverse arrows as in the figure. Otherwise surgery gives bigons.

So we now assume that a disk neighborhood of C_k is as in figure 52. Let v be any vertex of the back-face and let e be the $\{i, j\}$ -edge of P incident with v . If the other vertex w of e is part of a solid ij -crossing, then use lemma 10.4 to introduce a circle of edges (with all broken crossings) contained in the boundary of the disk neighborhood of C_k . Do this so that the vertex introduced on the edge e cancels with the crossing at w . By repeating this procedure as necessary, we may assume that the vertex w of e is part of a solid ab -crossing, where $\{a, b\} \not\subseteq \{i, j\}$. Now if $\{a, b\} \not\subseteq \{i, j, k\}$, with say $a \notin \{i, j, k\}$, then a deformation given by lemma 10.3 which changes all the letters k in figure 52 to a eliminates C_k with no other effect on P_k . And if $\{a, b\} \subseteq \{i, j, k\}$, then after possibly introducing a pair of canceling swirls, we may assume that the vertex w of e is part of a solid ki - or kj -crossing. Also since the boundary of a disk neighborhood of C_k represents the word w_{ij}^8 (up to swirls), w is not a vertex of C_k . What's more, the above procedure may increase the number of vertices of C_k , but not of C_k^* .

Now let z be the cycle indicated in figure 62 below. By adding nz for suitable $n \in \{0, 1, \dots, 7\}$ to P , the graph C_k can effectively be rotated so that w cancels with a vertex of the backface of C_k . This decreases the number of vertices of P_k^* , but may change P by an element of the group E_n , as the proof of Theorem 12.1 below shows.

Now assume that F is a biggest face which is not surrounded by squares. Then by lemma 10.17, two edges of F , say e and f , are not edges of squares. If e and f are both edges of a face G different from F , then $F \cap G$ is not connected. So, by lemma 10.18, C_k can be disconnected.

So assume e and f are edges of distinct faces G and H , both different from F . At least one of G or H is a front-face, say G is. Clear G . By lemmata 10.12, 10.13 and 10.14, we may assume that G is a $4n$ -gon, where $n > 1$, with arrows on consecutive edges of the same type pointing in opposite directions.

Assuming e is an i -edge, let e' be another i -edge of G which points inward if e does, and outward otherwise. Surgery in G using e and e' produces a face F' of C_k^* with two more edges than F had. It's easy to check that C_k is still innermost. Now, if F' is not surrounded by squares, repeat the above argument with this new innermost component. Since arbitrarily large faces cannot be produced in this way, one of the previously considered cases must eventually occur. $\square$

11. THE ASSOCIATION P

Throughout this section stable versions of spaces defined above, such as $\tilde{X}$ and $\tilde{W}$, are used. However, all arguments go through with the corresponding unstable spaces, e.g. $\tilde{X}_n$ and $\tilde{W}_n$, in place of the stable ones provided $n \geq 5$. Theorem 11.7 is used for these n .

Let $f: (S^2, \mathcal{C}) \rightarrow \mathcal{T}^*$ be any cell-to-cell map. We next describe a way of associating with f a 2-cycle in $Z_2(X)$ along with a map $S^2 \rightarrow X$ representing it. Both the cycle and map are denoted by $p(f)$. Lifts of $p(f)$ from X to $\tilde{X}$ will be denoted by $P(f)$. This association is well-defined only in the sense that it induces a homomorphism from $H_2(\mathcal{T})$ to the cokernel of the composite $H_2(\tilde{W}) \rightarrow H_2(\tilde{X}) \rightarrow H_2(\tilde{X})_{StN}$, as lemma 11.6 below shows. Of course, since $\mathcal{T}$ is contractible, this is the trivial homomorphism. We will use this in conjunction with Theorem 11.7 below in the computation of $H_2(\tilde{X})_{StN}$. In this section, we only use the fact that $\mathcal{T}$ is simply-connected.

The map $P(f)$ is not a lift of f up the map $F|: \tilde{X} \rightarrow \mathcal{T}^*$ and, in general, no such lift exists. Here $F|$ is the composite $\tilde{X} \xrightarrow{\mathcal{F}} \mathcal{S} \xrightarrow{q} \mathcal{T}$ restricted to the 2-skeleton of $\tilde{X}$ followed by the map $\mathcal{T} \rightarrow \mathcal{T}^*$. From the point of view of pseudo-isotopy theory, $p(f)$ is essentially obtained from f by eliminating all dovetail singularities by joining these in pairs.

Let $\{\sigma_i\}_{i=1}^n$ be the set of closed 2-cells of $\mathcal{C}$ labeled so that, for $1 < k \leq n$, $\bigcup_{i < k} \sigma_i$ is a disk and $\sigma_k \cap \bigcup_{i < k} \sigma_i$ contains at least one edge. Such an ordering of the 2-cells is possible since each edge of $\mathcal{T}^*$ has distinct vertices. Orient each σ_i so that, if σ_i also denotes the generator of the cellular chain group $C_2(S^2)$ corresponding to the oriented 2-cell σ_i , then $\sum_{i=1}^n \sigma_i$ represents a generator of $H_2(S^2)$.

Any σ_i which f takes to a degenerate simplex of $\mathcal{T}$, as defined in §8.1, is itself referred to as degenerate. Corresponding to each nondegenerate σ_i , we will define an element $\sigma'_i = \sigma_i^c + \sigma_i^w$ of the cellular chain group $C_2(X)$. In case σ_i is degenerate, $\sigma'_i = 0$ by definition. The cycle $P(f)$ will be defined as a lift of the cycle $\sum_{i=1}^n \sigma'_i$. We will also define sides of each σ'_i as certain edgepaths in the boundary of the set σ'_i , with one side corresponding to each 1-cell of the boundary of σ_i . One term of each side will be defined as the visible term of the side. The other terms will be referred to as invisible.

We first define σ'_1 (or, if σ'_1 is degenerate, σ'_i , where i is the least integer such that σ'_i is not degenerate). If the 2-cell $f(\sigma_1)$ of $\mathcal{T}^*$ corresponds to a marked partitioned graph, G say, choose any coloring of G consistent with the partition. Let σ'_1 be the 2-cell of X corresponding to the resulting marked colored graph. The three or four sides of σ'_1 are defined as just the three or four edges of this 2-cell.

If $f(\sigma_1)$ is a w -triangle of $\mathcal{T}^*$, let G be the corresponding marked graph with edges a , b , and c such that G/a , G/b , and G/c are the marked roses corresponding to the vertices of $f(\sigma_1)$. Choose a vertex of $f(\sigma_1)$, say the one corresponding to G/a , and an edge of $f(\sigma_1)$ containing this vertex, say the one corresponding to the partition $\{a, b\}$ of G . (Different choices here lead to exotic elements of $K_3(\mathbb{Z})$.) If more than three edges attach to the nonbasepoint vertex of G , let G^d be the marked graph shown in figure 53(i) with the requirement that $G^d/d = G$. Note that G^d depends on the choice of the vertex G/a of $f(\sigma_1)$ made above since the edge a of G is subdivided to obtain G^d . We define σ'_1 to be the three 2-cells of X indicated in figure 53(ii). Here each 2-cell corresponds to the marked graph G^d partitioned as indicated. Any consistent choice of colorings of the three partitioned graphs may be used. One side of σ'_1 , as indicated in (ii), is the sequence $b \wedge a, d \wedge b, b \wedge c, c \wedge d$ with $b \wedge a$ designated as the visible term. The other two sides are the two one-term sequences $a \wedge c$ and $c \wedge b$. The visible term of each of these last two sequences is (necessarily) defined as the only term of each. If the choice of the edge made above had been $a \wedge c$ rather than $a \wedge b$, then a figure like 53(ii), but reflected about the vertical line bisecting it, would define σ'_1 . The sides of σ'_1 would be defined as the two one-term sequences $b \wedge a$ and $b \wedge c$, and the four-term sequence $d \wedge b, b \wedge c, c \wedge d$, and $a \wedge c$, with $a \wedge c$ designated as the visible term of this last side.




FIGURE 53.

If G has just three edges, then $\sigma'_1 = 0$, by definition. The sides of σ'_1 are still defined though and, given the first choice of edge and vertex of $f(\sigma_1)$ made above, are the sides of the triangle in figure 53(iii). Here each edge in X of a side corresponds to the marked graph G with the indicated partition. Any consistent choice of colorings may be used. The first term of the side $b \wedge a, a \wedge b, b \wedge c, c \wedge a$ is defined as the visible term of that side. This completes the definition of σ'_1 . We set $\sigma_1^c = \sigma'_1$ and $\sigma_1^w = 0$.

Now assume that σ'_i , along with its sides and one visible term per side, has been defined for each $i < k$. Also assume that the following hold for $i < k$:

- (a) Forgetting colorings defines a bijection from the visible terms of the sides of σ'_i to the edges of $f(\sigma_i)$.
- (b) If $i, j < k$, and σ_i and σ_j share an edge, then the sides of σ'_i and σ'_j corresponding to this edge are the same (up to a possible change in the orientation of the edgepath constituting the side) and have the same visible term.
- (c) Each of the two subsequences of each side of σ'_i consisting of the invisible terms coming before and after the single visible one corresponds to a (possible empty) word in the generators of the group W .

To define σ'_k , for $1 < k < n$, so as to satisfy (a), (b), and (c), first assume that the edge e of σ_k is not contained in $\cup_{i < k} \sigma_i$. Let s_j be the sides of the σ'_i , for $i < k$, which correspond to the edges of $\sigma_k \cap \cup_{i < k} \sigma_i$. If all the terms of s_j are visible, then σ'_k is defined as σ'_1 was, but using σ_k in place of σ_1 . In case $f(\sigma_k)$ is a w -triangle, let e be the chosen edge of σ_k so that the side of σ'_k corresponding to e contains all the invisible terms of the sides of σ'_k . Also if $\tau \in \sigma_i$ is any edge of $\sigma_k \cap \cup_{i < k} \sigma_i$, then the colorings of the marked graphs associated with σ'_k are defined to be those consistent with the coloring of the graph associated with the side of σ'_i corresponding to τ .

If any of the s_j contain invisible terms, then σ'_k is defined as $\sigma_k^c + \sigma_k^w$ where the term σ_k^w corresponds to 2-cells of X representing standard w -crossings. The marked partitioned graph or graphs corresponding to σ_k^c are, by definition, the same as those of σ'_k as defined above in the case where the s_j contain no invisible terms. The side or sides of σ'_k different from the s_j contain one triple of terms corresponding to a generator of W for each such triple in the s_j . In case σ_k is a w -triangle, there is one additional such triple. As usual, colorings are determined by those of graphs of any σ'_i where $i < k$ and $\sigma_i \cap \sigma_k \neq \emptyset$.

Figure 54 shows how the cells of σ_k^w and σ_k^c are arranged in two examples and so should clarify the definition of σ'_k . The outer square in (i) and the outer triangle in (ii) represent two




FIGURE 54. In each figure the inner rectangle or triangle is σ_i^c .

possible boundaries of σ_k . We assume, in each case, that only the bottom edge of σ_k is not one of the s_j . Additional marks in the figure are superimposed on the σ_k and indicate the cells in X corresponding to σ_k as follows. The dotted lines are w -lines. Where these cross the edges of σ_k indicate the positions of the corresponding triples of terms in the sides of σ'_k . The solid lines which cross the edges of the σ_k indicate the position of the visible terms in the sides of σ_k . Crossings of these solid lines with the dotted w -lines indicate standard w -crossings. Finally, the 2-cell of σ_k^c is indicated in each of (i) and (ii) by its edges, lightly drawn. Thus (ii) treats the case where $f(\sigma_k)$ is a w -triangle.

Finally we define σ'_n . The s_j , now defined as the sides in $\sigma_k \cap \cup_{i < n} \sigma_i$, form a loop (representing a relation in the group $\text{Aut}(F_n)$ for some n) in X which will bound σ'_n . Again σ'_n is defined as $\sigma_n^c + \sigma_n^w$ with σ_n^c and σ_n^w as above, except that now σ_n^w may also contain 2-cells of X representing standard ww -crossings. Parts (iii) and (iv) of figure 54, drawn with similar conventions as used in parts (i) and (ii), should clarify this. The circles in each indicate a loop in X corresponding to a word in the generators of the group W which represent the trivial element of W . The 2-cells of σ_n^w contained in these circles correspond to the defining relations of W used to trivialize this word. This completes the definition of $p(f)$ except that we allow σ_1^w to be nonzero provided the cells of σ_1^w make up standard w -crossings.

Recall that $P(f): S^2 \rightarrow \tilde{X}$ is either lift of $p(f): S^2 \rightarrow X$ and also denotes the corresponding spherical 2-cycle in $\tilde{X}$.

Parts (i) through (iv) of figure 54 suggest a natural subdivision of the CW -structure $\mathcal{C}$ of S^2 which makes the map $P(f): S^2 \rightarrow \tilde{X}$ cellular. This subdivision has 2-cells, called core 2-cells, corresponding to the σ_i^c . The remaining 2-cells make up w - and ww -crossings. The map $P(f)$ is not, in general, cell-to-cell with respect to this subdivision since it takes core 2-cells σ_i^c with $f(\sigma_i)$ corresponding to graphs with three edges to the 1-skeleton of X . But otherwise $P(f)$ restricts to a homeomorphism on each closed 2-cell of $\mathcal{C}$.

The map $P(f)$ is sometimes specified by just giving the core cells in S^2 and the colorings of the corresponding marked graphs, along with solid lines and dotted lines as in (i) through (iv) of figure 54. A solid line joins two sides of two core cells if these two sides correspond to the same edge of $\mathcal{C}$. The dotted w -lines cross the solid lines and each other indicating w - and ww -crossings, and begin and end in core cells corresponding to w -triangles.

Definition 11.1. A 2-cycle z in a CW -complex Y is called a simple spherical cycle if there is a cell-to-cell map $f: (S^2, \mathcal{C}) \rightarrow Y$ such that $z = \sum_i \text{gen}(f(\sigma_i))$. Here the σ_i vary over the 2-cells of $\mathcal{C}$ and the terms $\text{gen}(f(\sigma_i))$ are the generators of the cellular chain group $C_2(Y)$

corresponding to the 2-cells $f(\sigma_i)$, suitably oriented. The 2-cycle z is called a spherical cycle if it is a sum of simple spherical cycles.

The following technical lemma will be used often.

Lemma 11.2. *If Y is any simply-connected simplicial complex, then, using the simplices of Y as cells, each element of $H_2(Y)$ is represented by a spherical cycle.*

Proof. Let $z \in Z_2(Y)$ be a simplicial 2-cycle. Assume $f': S^2 \rightarrow Y$ is any map representing the homology class $[z] \in H_2(Y)$, meaning that $f'_*(1) = [z]$ for one of the generators 1 of $H_2(S^2)$. Approximate f' with a simplicial map $f: S^2 \rightarrow Y$, and then identify points in each simplex of S^2 which have the same image under f . The resulting quotient space $S^2/\sim$ has a natural simplicial structure and, if f' is not a constant map, consists of 2-spheres and 1-simplices which meet in discrete points. Moreover, the map $g: S^2/\sim \rightarrow Y$ induced by f is simplicial and is, in fact, cell-to-cell using the simplices of $S^2/\sim$ and Y as cells. So g restricted to each 2-sphere of $S^2/\sim$ is also cell-to-cell. Thus, if the 2-simplices of $S^2/\sim$ are denoted by e_i and oriented so that the quotient map $q: S^2 \rightarrow S^2/\sim$ preserves orientation (with S^2 oriented by the generator $1 \in H_2(S^2)$ mentioned above), then the 2-cycle $z' \equiv \sum_i g(e_i)$ is spherical and $[z'] = g_*(q_*(1)) = f_*(1) = [z]$ in $H_2(Y)$. $\square$

Now let $z \in Z_2(\mathcal{T})$ be any spherical 2-cycle. Choose cell-to-cell maps $f_j: (S^2, \mathcal{C}_j) \rightarrow \mathcal{T}^*$ such that $z = \sum_{i,j} \text{gen}(f_j(\sigma_{ij}))$, where, for fixed j , the σ_{ij} are the 2-cells of $\mathcal{C}_j$. Here the $\text{gen}(f_j(\sigma_{ij}))$ denote generators of $C_2(\mathcal{T}^*)$ as in Def. 11.1. Going from z to f_j can be done by first using Def. 11.1 to pick appropriate cell-to-cell maps $S^2 \rightarrow \mathcal{T}$ and then subdividing the domains of these to get the f_j . Let $P(z) = \sum_j P(f_j)$.

Definition 11.3. The group $H_2(\tilde{X}, \tilde{W})_{StN}$ is the cokernel of the composite $H_2(\tilde{W}) \rightarrow H_2(\tilde{X}) \rightarrow H_2(\tilde{X})_{StN}$.

Lemma 11.4. *The class of $P(z)$ in $H_2(\tilde{X}, \tilde{W})_{StN}$ is independent of the choices made in the definition of $P(z)$.*

Because of this, we often denote the class of $P(z)$ in $H_2(\tilde{X}, \tilde{W})_{StN}$ by $P_H(z)$.

Proof of 11.4. Let z and f_j be as above and let $g_k: (S^2, \mathcal{D}_k) \rightarrow \mathcal{T}^*$ be another choice of cell-to-cell maps such that $z = \sum_{k,l} g_k(\tau_{kl})$ where the τ_{kl} are the 2-cells of $\mathcal{D}_k$. Let

$$P_1(z) = \sum_j P(f_j) = \sum_{i,j} (\sigma_{ij}^c + \sigma_{ij}^w) \text{ and } P_2(z) = \sum_k P(g_k) = \sum_{k,l} (\tau_{kl}^c + \tau_{kl}^w).$$

Here the σ and τ with superscripts and subscripts denote consistent lifts of the like denoted cells in the definition of $p(f)$. So, for instance, if the 2-cell $g_k(\tau_{kl})$ of $\mathcal{T}^*$ is not a w -triangle, then the marked colored graph associated with the 2-cell τ_{kl}^c is, after forgetting colorings, the marked partitioned graph associated with $g_k(\sigma_{kl})$. Thus by using lemma 9.1 to surround core cells with dotted circles, we may assume that, if the 2-cells $f_j(\sigma_{ij})$ and $g_k(\tau_{kl})$ of $\mathcal{T}^*$ are the same and are not w -triangles, then the corresponding core cells σ_{ij}^c and τ_{kl}^c of $\tilde{X}$ are the same. Note that applications of lemma 9.1 may change the element of $H_2(\tilde{X})_{StN}$ which $P_1(z) - P_2(z)$ represents, but not the element of $H_2(\tilde{X}, \tilde{W})_{StN}$. If $f_j(\sigma_{ij}) = g_k(\tau_{kl})$ is a w -triangle, then a deformation like that indicated in figure 55 may also be needed. One can show that the two pictures in this figure differ by the trivial element of $H_2(\tilde{X})$ provided that




FIGURE 55.

the upper portion of the dotted curve on the right is a suitably chosen swirl. So we may now assume that $P_1(z) - P_2(z)$ consists entirely of cells making up swirls, and solid and dotted lines with standard w -crossings. Following any solid line gives a loop in X which can be eliminated (cf. the proof of the broken circles lemma 10.4). Thus $P_1(z) - P_2(z)$ is homologous to a 2-cycle consisting of w -lines and standard ww -crossings. Such a cycle represents the trivial element of $H_2(\tilde{X}, \tilde{W})_{StN}$. $\square$

The definition of P given above applies nearly verbatim to cell-to-cell maps $g: S^2 \rightarrow T_M(G)$ (where G is a fixed marked graph) and gives a 2-cycle in $\tilde{X}$ along with a map $S^2 \rightarrow \tilde{X}$, both of which we denote by $P'(g)$. Since the map G of section 8.1 takes degenerate simplices of the 2-skeleton of $\mathcal{T}$ to the 1-skeleton of its range, it turns out to be convenient to also define $P(g)$ for maps $g: S^2 \rightarrow T_M(G)$ which restrict to cell-to-cell maps on the 1-skeleton of the domain and which take each closed 2-cell either homeomorphically to a closed 2-cell or to the 1-skeleton. Such maps are referred to as almost cell-to-cell. The only additional remark required to define $P'(g)$ for an almost cell-to-cell map g is that σ'_i is defined to be 0 for each 2-cell σ_i of S^2 on which g is not injective. Note that the previous lemma, with a simpler proof, holds with $P'(g)$ in place of $P(z)$. We denote the class of $P'(g)$ in $H_2(\tilde{X}, \tilde{W})_{StN}$ by $P'_H(g)$.

The rather lengthy proof of the next lemma is given at the end of this section.

Lemma 11.5. *Let $f: S^2 \rightarrow T_M(G)$ be an almost cell-to-cell map where G is a fixed graph. Then $P'_H(f)$ is the trivial element of $H_2(\tilde{X}, \tilde{W})_{StN}$ regardless of choices.*

Lemma 11.6. *The association P induces a homomorphism $P_H: H_2(\mathcal{T}) \rightarrow H_2(\tilde{X}, \tilde{W})_{StN}$.*

Proof. Let $x \in H_2(\mathcal{T})$. By lemma 11.2, there is a simplicial, spherical 2-cycle z with $[z] = x$ in $H_2(\mathcal{T})$. We show that setting $P_H(x) = P_H(z)$ gives a well-defined function $P_H: H_2(\mathcal{T}) \rightarrow H_2(\tilde{X}, \tilde{W})_{StN}$.

Assume z' is another simplicial, spherical cycle with $[z'] = x$. Then $z - z' = \sum_i d\tau_i$ where the τ_i are generators of the group of simplicial 3-chains of $\mathcal{T}$. (As usual, we also denote the corresponding 3-simplices by τ_i .) Subdivide appropriate 2-simplices so that z , z' , and the $d\tau_i$ become spherical cycles in $\mathcal{T}^*$. Then, by lemma 11.4, we have $P_H(z) = P_H(z') + \sum_i P_H(d\tau_i)$.

With $\partial\tau_i$ denoting the topological boundary of the 3-simplex τ_i , let $h_i: \partial\tau_i \hookrightarrow \mathcal{T}$ be the inclusion map. Note that $P(d\tau_i) = P(h_i)$ in $H_2(\tilde{X}, \tilde{W})_{StN}$ by the definition of P on cycles and the cell-to-cell maps representing them. Let g_i be the composite $\partial\tau_i \hookrightarrow \text{image of } h_i \rightarrow T_M(G_{\tau_i})$. Here the first arrow is h_i with range restricted, and the second is the restriction of the map G (from the 2-skeleton of $\mathcal{T}^*$ to $\bigsqcup_{\sigma \in \mathcal{T}} T_M(G_\sigma)/\sim$) discussed at the end of section 8.1. Then, with suitable choices, we have $P(h_i) = P'(g_i)$ in $Z_2(\tilde{X})$. So, by lemma 11.4, equality holds in $H_2(\tilde{X}, \tilde{W})_{StN}$ regardless of choices. Since $P'(g_i) = 0$ by the previous lemma, each $P_H(d\tau_i) = 0$, and so P_H is well-defined. It is also a homomorphism by lemma 11.4. $\square$

The following theorem provides a crucial part of the computation of $H_2(\tilde{X})$. Let F be the composite $\tilde{X} \xrightarrow{\mathcal{F}} \mathcal{S} \xrightarrow{q} \mathcal{T}$, where $\mathcal{F}$ is the forgetful function and q the quotient map, both defined near the beginning of §8.

Theorem 11.7. *The composites $H_2(\tilde{X}) \xrightarrow{F_*} H_2(\mathcal{T}) \xrightarrow{P_H} H_2(\tilde{X}, \tilde{W})_{StN}$ and $H_2(\tilde{X}) \rightarrow H_2(\tilde{X})_{StN} \rightarrow H_2(\tilde{X}, \tilde{W})_{StN}$ agree.*

Proof. We first show that $H_2(\tilde{X})$ is generated by classes which are represented by simple spherical cycles, and then show that the two homomorphisms $H_2(\tilde{X}) \rightarrow H_2(\tilde{X}, \tilde{W})_{StN}$ indicated in the statement of the theorem agree on such generators.

Let $x \in H_2(\tilde{X})$. Since $\tilde{X}$ is simply-connected, there is a map $S^2 \rightarrow \tilde{X}$ representing x with image contained in the 2-skeleton $\tilde{X}^{(2)}$ of $\tilde{X}$. Make $\tilde{X}^{(2)}$ into a simplicial complex $\tilde{X}_*^{(2)}$ by adding one (of either) diagonal to each square 2-cell of $\tilde{X}^{(2)}$. Then, by lemma 11.2, $x = \sum_i [x_i]$ where each x_i is represented by a composite $S^2 \xrightarrow{f_i} \tilde{X}_*^{(2)} \hookrightarrow \tilde{X}$ with f_i cell-to-cell.

Fix a value of the subscript i . Let $\{c_l\}$ be the set of 1-simplices which f_i takes to the added diagonal edges of $\tilde{X}_*^{(2)}$. Write $\{c_l\} = \{d_j\} \sqcup \{e_k\}$ in such a way that there is a neighborhood of an interior point of c_l on which f_i is injective if and only if c_l is one of the e_k . For each j , the union of the two 2-simplices a_j and b_j which meet along d_j is a square. Subdivide each of a_j and b_j into two 2-simplices by adding the other diagonal d'_j to this square. Let S_*^2 be the resulting simplicial complex and let m_j be the vertex of $d_j \cap d'_j$ of S_*^2 . Define a simplicial map $f'_i: S_*^2 \rightarrow \tilde{X}_*^{(2)}$ which agrees with f_i on all the vertices of S_*^2 except the m_j . Define $f'_i(m_j)$ to be the common image under f_i of the two endpoints of d'_j . Note that f_i and f'_i are homotopic.

Next delete the 1-simplices e_k from S_*^2 so as to give a CW -complex $\mathcal{C}$ with one square 2-cell corresponding to each e_k . Note that f'_i , defined on $\mathcal{C}$ in the obvious way, followed by the map $\tilde{X}_*^{(2)} \rightarrow \tilde{X}^{(2)}$, takes each of these square cells homeomorphically to a square 2-cell of $\tilde{X}$. Thus, identifying points in cells of $\mathcal{C}$ with the same image under f'_i shows, as in the proof of 11.2, that $[x_i] = \sum_j [z_j]$ where each z_j is a cycle represented by a cell-to-cell map $S^2 \rightarrow \tilde{X}$. It follows that $H_2(\tilde{X})$ is generated by classes represented by simple spherical cycles.

For the remainder of the proof, assume that $x \in H_2(\tilde{X})$ equals $[z]$ where z is represented by the cell-to-cell map $f: S^2 \rightarrow \tilde{X}$. Subdivide the cells of S^2 giving S_*^2 so that $f: S_*^2 \rightarrow \tilde{X}$ is cell-to-cell using the simplices of $\tilde{X}$ as cells in the range. Then the composite $\mathcal{F} \circ f: S_*^2 \rightarrow \mathcal{T}$ is simplicial. As usual, let $q': S^2 \rightarrow S^2/\sim$ be the quotient map defined by identifying points in each simplex of S_*^2 with the same image under $\mathcal{F} \circ f$, and let $g: S^2/\sim \rightarrow \tilde{X}$ be the map induced by $\mathcal{F} \circ f$. At most one of the two 1-simplices making up each 1-cell of S^2 is contracted to a point by q' . Also, as one can check, $q'(e_i)$ is a disk for each 2-cell e_i of S^2 . It follows that $S^2/\sim$ is homeomorphic to a 2-sphere. So $P(g)$ is defined and $P(g) = P_H(F_*(x))$ since the cell-to-cell map g represents $F_*(x)$.

For each 2-cell e_i of S^2 , let G_i^{par} be the marked partitioned graph, with coloring forgotten, corresponding to the 2-cell $f(e_i)$ of $\tilde{X}$, and let G_i be the same, but with the partition also forgotten. Denote the 2-cell of $T_M(G_i)$ corresponding to G_i^{par} by $cell(G_i^{par})$. Orient this cell so that the map from the 2-cell $f(e_i)$ to it, defined by forgetting coloring, preserves orientation. Here we are assuming that the cell $f(e_i)$ is oriented so that f preserves orientation, with




FIGURE 56.

S^2 , in turn, oriented by the generator 1 of $H_2(S^2)$ such that $f_*(1) = x$. Let $gen(G_i^{par})$ be the generator of the cellular chain group $C_2(T_M(G_i))$ corresponding to the oriented 2-cell $cell(G_i^{par})$.

Let τ_{ij} be the 2-cells of $T_M(G_i)$ in the image of the composite $h \circ F \circ f|_{e_i}$, where h is the map, with domain and range suitably restricted, from $\mathcal{T}$ to $\mathcal{T}_*$ followed by the map G defined at the end of section 8.1. Let $gen(\tau_{ij})$ be the corresponding generators of $C_2(T_M(G_i))$ with orientations chosen, in the case where F is not injective on any of the three or four 1-cells of the boundary of $f(e_i)$, so that $y_i = gen(G_i^{par}) + \sum_j gen(\tau_{ij})$ is a cycle. In case F is injective on any of the edges of $f(e_i)$, let $y_i = gen(G_i^{par}) + \sum_j gen(\tau_{ij}) + \sum_k w_i^k$, where the w_i^k are generators corresponding to w -triangles, with one for each 1-cell of $f(e_i)$ on which F is injective, and again with orientations such that y_i is a cycle.

Each y_i is represented by a nearly cell-to-cell map $f_i: S^2 \rightarrow T_M(G_i)$, and we define $P'(y_i)$ as $P(f_i)$. It is not difficult to check that $\sum_i P'(y_i) = x - P(g)$ in $H_2(\tilde{X}, \tilde{W})_{StN}$. This follows, for instance, by an argument like that used to prove lemma 11.4. Since each $P'(y_i) = 0$ by lemma 11.5 and $P(g) = P_H(F_*(x))$, the theorem is proved. $\square$

Proof of 11.5. We first prove the lemma in the special case where $f: \partial\tau \rightarrow T(G)$ is the cell-to-cell inclusion of the boundary of a 3-cell τ of $T(G)$. If τ corresponds to a partitioned graph Γ , then the 2-cycle $P'(f)$ bounds a 3-cell in $\tilde{X}$, the one corresponding to Γ with marking and coloring those chosen in the definition of P' . So, in this case, $P'(f)$ is trivial in $H_2(\tilde{X})$ regardless of choices.

If τ corresponds to a w -tetrahedron, then the graph Γ associated with τ has two vertices, the basepoint $*$, and v say. Let a, b, c , and d be the four edges making up the four maximal trees corresponding to the vertices of τ . With suitable choices $P(f)$ consists of the 2-cells, suitably oriented, shown in figure 56. The graphs associated with these 2-cells are drawn near them. The heavily drawn dangling edge in these graphs represent all the halfedges incident with v , other than a, b, c , and d .

We next show that we may assume that Γ has only the four edges a, b, c , and d . More specifically, let G' be any graph having as some blowdown a graph Γ' with just two vertices and four edges, each edge of which meets both vertices. Also let τ' be the w -tetrahedron




FIGURE 57.

of $T(G')$ corresponding to Γ' . With $f': \partial\tau' \rightarrow T(G')$ the inclusion, we will show that, with suitable choices, $P(f') = P(f)$ in $H_2(\tilde{X})$.

Using 3-cells in $\tilde{X}$ corresponding to the graph G in figure 57 on the right, suitably marked and with various colorings, one can see that the 2-chain c_1 in the figure, oriented so that $dc_i = d\sigma'_1$, is homologous to σ'_1 . More specifically, $c_1 - \sigma'_1$ is the boundary of a 3-chain with six terms. These correspond to the graph G with each of the six partitions $\{\{c, d, d'\}, \{a, y\}\}$, $\{\{b, c\}, \{d, d'\}, \{a, y\}\}$, $\{\{b, d, d'\}, \{a, y\}\}$, $\{\{a, d, y\}, \{c, d'\}\}$, $\{\{a, d, y\}, \{b, c\}\}$, and $\{\{a, d, y\}, \{b, d'\}\}$. Colorings are determined by the coloring at any vertex common to σ'_1 and c_1 . Similarly the 2-chain c_2 indicated in the left of figure 57 is homologous to σ'_2 by way of the three 3-cells of $\tilde{X}$ corresponding to the graph on the left in the figure, suitably marked, with the three partitions $\{\{a, a', c, x\}\}$, $\{\{a, a', x\}, \{b, c\}\}$, and $\{\{a, a', b, x\}\}$. Also define 2-chains c_3 and c_4 similar to c_2 in such a way that c_3 and c_4 are homologous to σ'_3 and σ'_4 , respectively, by way of graphs like those in Fig. 56 corresponding to σ'_3 and σ'_4 , but with the edges labeled a subdivided by a vertex and with the dangling halfedges attached to this vertex. Then with choices like those giving figure 56, the cycle $P(f') = \sum_{i=1}^4 c_i + \sigma_4^w$ is homologous to $P(f)$.

So we now assume that Γ has just four edges. Consider figure 58 which shows three 3-cells of $\tilde{X}$ corresponding to the colored graphs shown near them. The 3-disk these three cells form is denoted by D . By the back face of D , we mean the three squares in the boundary of D with one edge $d \wedge a$ and the other edge one of $d' \wedge b$, $b \wedge c$, or $c \wedge d'$. With suitable markings, the nine faces in the boundary of D other than the three back faces are homologous with the standard w -crossing in σ_4^w . Thus this w -crossing is also homologous to the three back faces of D . With the w -crossing represented in this last way, the portion of $P(f)$ other than the ww -crossing in σ_4^w is the negative of this ww -crossing. Therefore, with the choices above, $P'(f) = 0$ in $H_2(\tilde{X})$. So, by lemma 11.4, $P'(f) = 0$ in $H_2(\tilde{X}, \tilde{W})_{StN}$ regardless of the choices made.

Now assume that $f: \partial\tau \rightarrow T(G)$ is the cell-to-cell inclusion of the boundary of a w -prism τ of $T(G)$. Let G' be the graph corresponding to τ . As a first case, assume G' has a subgraph $\{a, c, d, x, y\}$ as in figure 59(i) where the triangular faces of τ correspond to G'/x and G'/y . Possible additional edges of G' are indicated by the dangling half edges in the figure. There




FIGURE 58.




FIGURE 59.

may be still more half edges incident with $*$. But these remain so throughout the discussion, so are not shown. Fig. 60 illustrates $P(f)$ with certain choices. Colored graphs are drawn within the cells they correspond to. The three cells of $P(f)$ corresponding to G'/x are not shown, but we assume choices are such that these are the same as those corresponding to G'/y except for the markings of the colored graphs. Let c be the cube of $\tilde{X}$ corresponding to the colored graph in figure 59(iii) marked so that the vertex v in figure 60A is the vertex of c corresponding to the maximal tree consisting of the edges b , d , and y . We will show that $P(f)$ is homologous to the boundary of c . Let p_d , p_y , and p_c be the O-prisms shown in figure 60B. These correspond to the colored graphs drawn near them and are marked so that the vertices v , v' , w , and w' in this figure are the same as those with like labels in figure 60A. Let p_x be the O-prism corresponding to the same colored graph as p_y , but marked so that the edge analogous to the one labeled $d \wedge c$ in p_y joins v' and w' . Note that for $i = d, c, x$, and y , the only square face of p_i which is not contained in $P(f)$ is the face of the cube c corresponding to the graph in figure 59(iii) with the edge i blown down.

Let t_v and t_w be the O-tetrahedra shown in figure 60B, with t_v the one incident with v . Let t'_v and t'_w be the O-tetrahedra corresponding to the same colored graphs as t_v and t_w , respectively, but marked so that they are incident with v' and w' , respectively. Then, with suitable orientations, we have

$$P(f) + d(p_d + p_y + p_c + q_y) + d(t_v + t_w + t'_v + t'_w) = dc.$$

So, again by lemma 11.4, $P(f) = 0$ in $H_2(\tilde{X}, \tilde{W})_{StN}$ regardless of choices.

For another case, assume G' is as in figure 59(i), but with basepoint the vertex on the the upper right. With choices like those made in the previous case, $P(f)$ is as in figure 60A except for the position of the basepoints and for the colored graph in the middle of figure 60A. This should now be the colored graph Γ/y where Γ is the colored graph in figure 59(ii) with suitable dangling halfedges. Let c be the cube in $\tilde{X}$ corresponding to Γ , marked so that $\Gamma/\{a', d, y\}$ is the vertex of $P(f)$ analogous to the vertex v in figure 60A. We show that




FIGURE 60.

$P(f)$ is homologous to dc . Let γ be the graph in figure 59(ii) and let p_d be the O-prism in $\tilde{X}$ corresponding to γ with the partition $\{\{a, a', d\}, \{x, y\}\}$ and marked so that $\gamma/\{a', d, y\}$ is the vertex analogous to v . Also let p_c be the O-prism corresponding to γ partitioned by $\{\{a, a', c\}, \{x, y\}\}$ and marked so that $\gamma/\{a', c, y\}$ is the vertex analogous to w . Then, in $H_2(\tilde{X})$, we have, with suitable orientations, $P(f) + d(p_c + p_d) = dc$ which completes this case.

The case in which the basepoint of G' is the upper left vertex of the graph in figure 59(i) is similar to the case just considered unless the vertex incident with the edges a, c , and d is incident with no other edges. Then choices can be made so that $P(f)$ is trivial as a 2-cycle in $\tilde{X}$. The remaining cases in which τ is a w -prism are those where the edges a, c , and d are all incident with the same two vertices. These cases are easier than those already considered and are omitted.

Also omitted are the cases in which τ is a pillow. These are also easier than those above, especially if choices are made in defining $P(f)$ so that there is a single w -line with no crossings. Though in some cases this w -line will pass through the cells which realize the relation $w_{ij}^L = w_{ij}^R$.

Now assume that $f: (S^2, \mathcal{C}) \rightarrow T(G)$ is any cell-to-cell map. We will show, with the help of frequent implicit applications of lemma 11.4, that if f is transformed to g using any one of the operations (a) through (d) above, then $P(f) = P(g)$ in $H_2(\tilde{X}, \tilde{W})_{StN}$. This, along with lemma 7.1, will complete the proof. First assume that operation (a) is used to transform f to g , and let $D = \cup_i \sigma_i$ and τ be as in the statement of (a). Also let $i: \partial\tau \rightarrow T(G)$ be the inclusion of the boundary of τ . By, for instance, choosing the σ_i to come first in the orderings of the 2-simplices of $\mathcal{C}$ and $\partial\tau$ used to define $P(f)$ and $P(i)$, we may assume that these agree when restricted to D . Thus, since $P(f) - P(i) = P(g)$, the first part of the proof implies that $P(f) = P(g)$.




FIGURE 61.

Next assume that g is obtained from f by an operation of type either (b), (c), or (d), and let σ and τ be as in the statement of (b), (c), or (d), as appropriate. In the case of (b) or (d), we may assume, with suitable choices, that $\sigma' = -\tau'$. Here, and in the next few lines, the primes have the same meaning as in the definition of P in section 11. So it follows that $P(f) = P(g)$. In the case of (c), we may assume, after introducing w -crossings and then ww -crossings as needed, that $\sigma' + \tau'$ is a cycle which, as in the proof of 11.4, represents the trivial element of $H_2(\tilde{X}, \tilde{W})_{StN}$. So again, $P(f) = P(g)$. Here, $P(g) = P(g_1) + P(g_2)$ where g_1 and g_2 are the two cell-to-cell maps which f is transformed to by (c). $\square$

12. THE COMPUTATION OF $H_2(\tilde{X}_n)_{StN_n}$

Let $\mathbb{Z}^n$ be the free abelian group of rank n .

Theorem 12.1. *If $n \geq 5$, then $H_2(\tilde{X}_n)_{StN_n} \approx \mathbb{Z}_{24} \oplus \mathbb{Z}^n$.*

Proof. As remarked after theorem 10.1, we have $H_2(\tilde{X}_n)_{StN_n} = E_n$. We begin by finding a convenient set of generators for this group. The element $w_{14}^L \cdot I_1^{-3,-4}$ of $H_2(\tilde{X}_n)_{StN_n}$ is represented by the picture $I_1^{-3,-4}$ surrounded by a circle labeled w_{14}^L and with transverse orientation pointing inward. By the three circles lemma 9.2, this circle can be moved past all the crossings in the picture, except for those corresponding to the relation $w_{ij}^L = w_{ij}^R$. There are five of these, but because of their signs, four cancel. So, in $H_2(\tilde{X}_n)_{StN_n}$, we have

$$I_1^{-3,-4} = w_{14}^L \cdot I_1^{-3,-4} = I_4^{3,-2} - C_{14}^2.$$

Adding another surrounding circle gives

$$I_1^{-3,-4} = (w_{14}^L)^2 \cdot I_1^{-3,-4} = I_1^{3,-2} - (C_{14}^2 + C_{14}^4).$$

Now consider the picture g_{StN} shown in Figure 61. Cutting it horizontally through the middle and adding the crossing 3 to the top component and -3 to the bottom one gives the picture $-I_k^{-3,-4} + w_{kj}^L \cdot (-I_k^{3,-2})$. Thus

$$g_{StN} = -(I_k^{-3,-4} + I_k^{3,-2})$$

in $H_2(\tilde{X}_n)_{StN_n}$.

It follows from the three displayed equations above and Theorem 10.1 that the group $H_2(\tilde{X}_n)_{StN_n}$ is generated by the classes of $I_i^{-3,-4}$, $i \in \mathbf{n}$ and g_{StN} . The map $\Theta: H_2(\tilde{X}_n)_{StN_n} \rightarrow H_2(\tilde{X}_n/StN_n)$ takes the n classes $I_i^{-3,-4}$, $i \in \mathbf{n}$ onto the free abelian group $H_2(\tilde{X}_n/StN_n)$ of rank n . Thus these generate a free abelian subgroup of $H_2(\tilde{X}_n)_{StN_n}$ of rank n . The remaining generator g_{StN} is in the kernel of Θ and so generates it. One can show that $I_1^{-3,-4}$ and $I_1^{3,-2}$ are sent by the map $H_2(\tilde{X}_n)_{StN_n} \rightarrow H_2(\tilde{V})$ to the generator of $K_3(\mathbb{Z}) \approx \mathbb{Z}_{48}$ exhibited by Igusa in Fig. 7.6 of [6]. Thus the class of g_{StN} in $H_2(\tilde{X}_n)_{StN_n}$ has order at least 24. (Alternatively, Hatcher observed in [3] that Waldhausen's splitting theorem implies that the homology of the group $\text{AUT} = \lim_{n \rightarrow \infty} \text{Aut}(F_n)$ contains that of the symmetric group $\Sigma = \lim_{n \rightarrow \infty} \Sigma_n$ as a direct summand. Similarly for homotopy groups. In particular, $H_3(StN) = \pi_3(BStN^+) = \pi_3(BAUT^+)$ contains $\pi_3(B\Sigma^+) = \pi_3^S = \mathbb{Z}_{24}$ as a direct summand. As in the proof of Theorem 12.2 below, the kernel of Θ is $H_3(StN)$.) So to conclude the proof we show that the class of g_{StN} in $H_2(\tilde{X}_n)_{StN_n}$ has order at most 24.

We will use the fact that $g_{StN} = U$ in $H_2(\tilde{X}_n)_{StN_n}$ where U is as shown in Figure 61. To see this, note that the upper left portions of both of these pictures are the same. So they cancel in the difference $g_{StN} - U$ leaving a circular edge containing only the two swirls $-A4$, shown in Figure 51 and $A2$, which is the knot-like swirl at the bottom of the picture at the right of Figure 42. This circle is trivial in $H_2(\tilde{X}_n)_{StN_n}$ as in the first case of the proof Theorem 10.19.

Consider the picture S shown in Figure 62. The four swirls in the picture are all $A1$. Two have positive orientations and two negative ones as shown. We first show that $S = 6g_{StN}$, and then show that $4S = 0$, which will complete the proof.

Note that the middle triangle of the picture U in Figure 61, but with i and j permuted, is the same as the upper right triangle in S . Similarly, the triangle in the lower right of S is the same as the inner one of U if the transverse orientations of the i -edges of U are changed. The other two triangles in the right-hand component of S_k are the same as the inner triangle of two other translates of U . Since $\Theta(U) = 0$, these four translates of U are represented by pictures agreeing with U 's if labels are forgotten. So the class of $S - 4U$ in $H_2(\tilde{X}_n)_{StN_n}$ is represented by a picture $(S - 4U)'$ looking like the picture S of Figure 62 except all 12 crossings on the right are changed so that the broken edges are solid. Using Lemma 10.5 each of the four swirls in $(S - 4U)'$ can be moved (in either direction) past the first broken edge of $(S - 4U)'_k$ encountered. Crossings then cancel in pairs leaving the picture $S - 4U$ of Figure 63. Moving the swirls does not change the class of the picture in $H_2(\tilde{X}_n)_{StN_n}$ since all elements of $\langle \emptyset \rangle$ introduced in moving the swirls cancel.

Now the lower half of the picture H_{A2} in Figure 64 agrees with the lower half of the picture $S - 4U$, except for the swirl $A2$. Note that the two crossings labeled -2 and -4 which are part of one of the bigons in H_{A2} cancel with two of the crossings of the swirl $A2$ in H_{A2} . Similarly, the two crossings which are part of the other bigon cancel with crossings of $A1$, leaving the picture $-I_k^{3,-2}$. Thus $H_{A2} = -I_k^{3,-2}$ at the chain level.

Let H denote H_{A2} with the swirl $A2$ changed to $-A1$ so as to agree with more of $S - 4U$. Then, in $H_2(\tilde{X}_n)_{StN_n}$, we have

$$H = H_{A2} - (A1 + A2) = H_{A2} - I_k^{-2,3} = -(I_k^{3,-2} + I_k^{-3,-4}) = g_{StN} = U.$$






FIGURE 62. The cycle S

The picture $(w_{il}^L)^2 \cdot H$ is the same as H except for the transverse orientations of its i -edges. So a portion of it agrees with the upper half of the picture $S - 4U$, except that the two swirls of $(w_{il}^L)^2 \cdot H$ are $A4$ and $-A4$ rather than $A1$ and $-A1$. But since the two swirls of $(w_{il}^L)^2 \cdot H$ have opposite signs, they can be changed to $A1$ and $-A1$ without changing the class of $(w_{il}^L)^2 \cdot H$. Let $H_{\Delta i}$ denote the picture $(w_{il}^L)^2 \cdot H$ with its swirls changed in this way. Thus, $H = H_{\Delta i}$ in $H_2(\tilde{X}_n)_{StN_n}$. Now $H + H_{\Delta i} = S - 4U$ at the chain level and so $2U = 2H = S - 4U$ or $S = 6U = 6g_{StN}$ in $H_2(\tilde{X}_n)_{StN_n}$.

A picture representing nS in $H_2(\tilde{X}_n)_{StN_n}$, where $n \in \mathbb{Z}$, looks, up to swirls, like the picture S , except that if a dotted edge e_i of S joins the vertex v_i to the vertex w_{i+1} , then in nS there is an edge e'_i joining v_i to w_{i+n} . Here $w_{i+n} = w_m$ if $i + n \equiv m \pmod{8}$, and the swirls on e'_i are those on the edges $e_i, e_{i+1}, \dots, e_{i+n-1}$. So, for example, $8S = 0$ in $H_2(\tilde{X}_n)_{StN_n}$ even if $n = 3$. (The four swirls on each of the eight edges which join the two components of $(8S)_k$ cancel in pairs, leaving the trivial cycle.)

To see that $4S = 0$ if $n \geq 5$, use Lemma 10.3 to change the direction of the transverse orientations of all of the k -edges in one of the two components of $(4S)_k$. Then swirls again cancel leaving the trivial cycle. $\square$

The next theorem brings together a number of the main results. It also incorporates Lee and Szczarba's result in [8] that $K_3(\mathbb{Z}) \approx \mathbb{Z}_{48}$.

FIGURE 63. The cycle $S - 4U$ FIGURE 64. The cycle H_{A2}

Theorem 12.2. *If $n \geq 7$, there is a commutative diagram*

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \mathbb{Z}_{24} & \longrightarrow & \mathbb{Z}_{24} \oplus \mathbb{Z}^n & \longrightarrow & \mathbb{Z}^n \longrightarrow 0 \\
 & & \downarrow \approx & & \downarrow \approx & & \downarrow \Theta^{-1} \approx \\
 0 & \longrightarrow & H_3(StN_n) & \longrightarrow & H_2(\tilde{X}_n)_{StN_n} & \longrightarrow & H_2(\tilde{X}_n/StN_n) \longrightarrow 0 \\
 & & \downarrow \approx & & \downarrow f & & \downarrow \\
 0 & \longrightarrow & \mathbb{Z}_{24} & \longrightarrow & K_3(\mathbb{Z}) & \longrightarrow & \mathbb{Z}_2 \longrightarrow 0
 \end{array}$$

where f is induced by abelianization. All rows are exact, and four of the six vertical arrows are isomorphisms as indicated. The remaining two are surjective.

Proof. In the spectral sequence of the covering $\tilde{X}_n \rightarrow \tilde{X}_n/StN_n$ we have $E_{0,2}^2 = E_{0,2}^3 = H_2(\tilde{X}_n)$ and $E_{3,0}^2 = E_{3,0}^3 = H_3(StN_n)$ since $H_1(StN_n) = H_2(StN_n) = 0$. Also, since $H_3(\tilde{X}_n/StN_n) = 0$, the differential $d^3 : H_3(StN_n) \rightarrow H_2(\tilde{X}_n)$ is injective giving exactness of the middle row. The other assertions follow from Theorem 12.1 and its proof and Theorem 6.1. $\square$

Exotic elements of $K_3(\mathbb{Z})$ appear, in some sense, as nontrivial elements of the homology of triangular subgroups. Using the notation of remark 6.4, where this was discussed, we have

Theorem 12.3. *The class of the cycle t^t in $H_2(T)$ is nontrivial.*

Proof. The map

$$\begin{array}{ccc} \tilde{X} & \xrightarrow{\tilde{\mu}} & \tilde{X}^{alg} \\ \downarrow & & \downarrow \\ \tilde{X}/StN & \longrightarrow & \tilde{X}^{alg}/StN \end{array}$$

of covering spaces defined in §2.10 induces a map of the associated spectral sequences. This in turn gives the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & H_3(StN_n) & \longrightarrow & H_2(\tilde{X}_n)_{StN_n} & \xrightarrow{\Theta} & H_2(\tilde{X}_n/StN_n) \longrightarrow 0 \\ & & \downarrow = & & \downarrow \tilde{\mu}_* & & \downarrow \tilde{\mu}_* \\ & & H_3(StN_n) & \xrightarrow{j} & H_2(\tilde{X}_n^{alg})_{StN_n} & \longrightarrow & H_2(\tilde{X}_n^{alg}/StN_n) \longrightarrow 0 \\ & & & \searrow d & \downarrow a & & \uparrow i_* \\ & & & & K_3(\mathbb{Z}) & & H_2(ET/T) \end{array}$$

with exact rows. By the choice of t^t we have $i_*[t^t] = \tilde{\mu}_*[t]$. We claim this last class is nontrivial. Indeed, if $e \in H_2(\tilde{X}_n)_{StN_n}$ satisfies $\Theta(e) = [t]$, then $a(\tilde{\mu}_*(e))$ is not in the image of the injection d . Thus $\tilde{\mu}_*(e)$ is not in the image of j , which gives $\tilde{\mu}_*[t] \neq 0$. So $[t^t] \neq 0$. $\square$

REFERENCES

- [1] M. Culler, K. Vogtmann, Moduli of graphs and automorphisms of free groups, *Invent. math.* **84** (1986), 91–119.
- [2] S. Gersten, A presentation for the special automorphism group of a free group, *J. Pure and Appl. Algebra* **33** (1984), 296–279.
- [3] A. Hatcher, Homological stability for automorphism groups of free groups, *Comment. Math. Helv.* **70** (1995), 39–62.
- [4] A. Hatcher, W. Thurston, A presentation for the mapping class group of a closed orientable surface, *Topology* **19** (1980), 221–237.
- [5] A. Hatcher, K. Vogtmann, Cerf theory for graphs, *J. London Math. Soc.* **58** (1998), 633–655.
- [6] K. Igusa, The Borel regulator map on pictures. I. A dilogarithm formula, *K-Theory* **7** (1993), 201–224.
- [7] J. Kiralis, A non-abelian K-theory and pseudoisotopies of 3-manifolds, *K-Theory* **10** (1996), 135–174.
- [8] R. Lee, R. H. Szczarba, The group $K_3(\mathbb{Z})$ is cyclic of order forty-eight, *Ann. of Math.* **104** (1976), 31–60.

- [9] J.-L. Loday, Cyclic Homology, *Grundlehren Math. Wiss.* 301, Springer, 1998.
- [10] S. Mac Lane, Homology, *Grundlehren Math. Wiss.* 114, Springer, 1963.
- [11] A. A. Suslin, On the equivalence of K -theories, *Comm. in Alg.* 9 **15** (1981), 1559-1566.
- [12] I. A. Volodin, Algebraic K -theory as an extraordinary homology theory on the category of associative rings with a unit, *Izv. Akad. Nauk SSSR Ser. Mat.* **35** (1971), 844-873.
- [13] F. Waldhausen, *Algebraic K-theory of spaces*, Springer lecture notes in Math. 1126 (1984), 318-419.